

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Bird Control Seminars Proceedings

Wildlife Damage Management, Internet Center  
for

11-1-1979

## DISPERSAL PATTERNS AND HABITAT RELATIONSHIPS OF ROOSTING AND FLOCKING RED-WINGED BLACKBIRDS

Ronnie J. Johnson

University of Nebraska-Lincoln, ronj@clmson.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/icwdmbirdcontrol>



Part of the [Environmental Sciences Commons](#)

---

Johnson, Ronnie J., "DISPERSAL PATTERNS AND HABITAT RELATIONSHIPS OF ROOSTING AND FLOCKING RED-WINGED BLACKBIRDS" (1979). *Bird Control Seminars Proceedings*. 35.

<https://digitalcommons.unl.edu/icwdmbirdcontrol/35>

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Bird Control Seminars Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# DISPERSAL PATTERNS AND HABITAT RELATIONSHIPS OF ROOSTING AND FLOCKING RED-WINGED BLACKBIRDS

Ronnie J. Johnson★  
Department of Natural Resources  
Cornell University, Ithaca

Red-winged blackbird (*Agelaius phoeniceus*) dispersal patterns from a communal roost and associated habitat relationships were studied in central New York State during summer and fall, 1976 and 1977. Density of foraging birds in a 10° sector out to 48 km from a roost was highest at 10 km. The greatest density of after-hatching-year (AHY) males occurred at approximately 10 km; of hatching-year (HY) birds, males and females, at approximately 18 km. HY females generally foraged further than did HY males, as well as further than AHY males.

Relative proportions of each habitat type and arrangement pattern of habitats were mapped in the 10° sector. Corn covered 36% of the total area mapped, nearly twice as much as the next most abundant category, hay, which covered 19%. When dispersed from the roost, redwings used cornfields far more than any other habitat type. Data suggest that for the premigration period (before September 30) redwing habitat use averaged 77% in cornfields. During migration (after September 30) habitat use averaged 47% in cornfields and 48% in trees. Overall corn arrangement pattern did not correlate with overall redwing dispersal patterns. However, on a local level at various distances from a roost, redwings foraged primarily in cornfields.

---

\*Present address: Department of Forestry, Fisheries and Wildlife, University of Nebraska, Lincoln.