

10-15-2015

Diet-responsive MicroRNAs Are Likely Exogenous

Janos Zempleni

University of Nebraska-Lincoln, jzempleni2@unl.edu

Scott R. Baier

University of Nebraska-Lincoln, srbaier@hotmail.com

Kendal Hirschi

Baylor College of Medicine, kendalh@bcm.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/nutritionfacpub>

Zempleni, Janos; Baier, Scott R.; and Hirschi, Kendal, "Diet-responsive MicroRNAs Are Likely Exogenous" (2015). *Nutrition and Health Sciences -- Faculty Publications*. 37.

<http://digitalcommons.unl.edu/nutritionfacpub/37>

This Article is brought to you for free and open access by the Nutrition and Health Sciences, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nutrition and Health Sciences -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

[Letter]

Diet-responsive MicroRNAs Are Likely Exogenous

Janos Zempleni,¹ Scott R. Baier,¹ and Kendal Hirschi²

¹ Department of Nutrition and Health Sciences, University of Nebraska, Lincoln, Nebraska 68583–0806

² Department of Pediatrics, Baylor College of Medicine and USDA/ARS Children's Nutrition Research Center, Houston, Texas 77030

Correspondence — Emails: jzempleni2@unl.edu, srbaier@hotmail.com, or kendalh@bcm.edu

In a recent report Title *et al.* (1) fostered miRNA-375 and miR-200c knock-out pups to wild-type dams and arrived at the conclusion that milk microRNAs are bioavailable in trace amounts at best and that postprandial concentrations of microRNAs are too low to elicit biological effects. Their take home message is flawed. First, the majority of microRNAs in bovine milk are encapsulated in exosomes, thereby conferring protection against degradation and a mechanism for intestinal transport (2). Second, numerous dietary microRNAs have been cataloged among human samples (3). Third, the mere absence of a postprandial increase does not equate with zero absorption. Abundant microRNA targets in the intestinal mucosa or liver could be promoting rapid microRNA degradation, the classical concept of first-pass elimination. Artificial microRNA targets (“sponges”) have been used successfully to decrease the abundance of microRNAs in experimental settings. Fourth, the authors spaced sample collections days apart, which risks missing increases in microRNAs following milk consumption.

Title *et al.* (1) proposed that low concentrations of microRNAs lack biological activity. This notion is unsound as recent work suggests microRNAs may work at concentrations in the femtomolar to picomolar range (4, 5). We do agree with their proposal that other exosome cargos may contribute to the biological effects of milk intake and that studies of dietary microRNAs need to take advantage of protocols that allow distinction between microRNAs derived from dietary sources *versus* those derived from endogenous synthesis.

1. Title, A. C., Denzler, R., and Stoffel, M. (2015) Uptake and function studies of maternal milk-derived microRNAs. *J. Biol. Chem.* **290**, 23680–23691
2. Wolf, T., Baier, S. R., and Zempleni, J. (2015) The intestinal transport of bovine milk exosomes is mediated by endocytosis in human colon carcinoma Caco-2 cells and rat small intestinal IEC-6 cells. *J. Nutr.* 10.3945/jn.115.218586
3. Lukasik, A., and Zielenkiewicz, P. (2014) In silico identification of plant miRNAs in mammalian breast milk exosomes— a small step forward? *PLoS ONE* **9**, e99963
4. Fabbri, M., Paone, A., Calore, F., Galli, R., Gaudio, E., Santhanam, R., Lovat, F., Fadda, P., Mao, C., Nuovo, G. J., Zanesi, N., Crawford, M., Ozer, G. H., Wernicke, D., Alder, H., Caligiuri, M. A., Nana-Sinkam, P., Perrotti, D., and Croce, C. M. (2012) MicroRNAs bind to Toll-like receptors to induce prometastatic inflammatory response. *Proc. Natl. Acad. Sci. U.S.A.* **109**, E2110–E2116
5. Bryniarski, K., Ptak, W., Martin, E., Nazimek, K., Szczepanik, M., Sanak, M., and Askenase, P. W. (2015) Free extracellular miRNA functionally targets cells by transfecting exosomes from their companion cells. *PLoS ONE* **10**, e0122991