University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Robert Powers Publications

Published Research - Department of Chemistry

2014

Development of cyclobutene- and cyclobutane-functionalized fatty acids with inhibitory activity against *Mycobacterium tuberculosis*

Wantanee Sittiwong
University of Nebraska - Lincoln

Denise Zinniel
University of Nebraska-Lincoln, dzinniel2@unl.edu

Robert J. Fenton
University of Nebraska - Lincoln, rfenton2@unl.edu

Darrel Marshall
University of Nebraska - Lincoln

Courtney B. Story
University of Nebraska-Lincoln

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/chemistrypowers>

Sittiwong, Wantanee; Zinniel, Denise; Fenton, Robert J.; Marshall, Darrel; Story, Courtney B.; Kim, Bohkyung; Lee, Ji-Young; Powers, Robert; Barletta, Raul G.; and Dussault, Patrick, "Development of cyclobutene- and cyclobutane-functionalized fatty acids with inhibitory activity against *Mycobacterium tuberculosis*" (2014). *Robert Powers Publications*. 38.
<https://digitalcommons.unl.edu/chemistrypowers/38>

This Article is brought to you for free and open access by the Published Research - Department of Chemistry at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Robert Powers Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Wantanee Sittiwong, Denise Zinniel, Robert J. Fenton, Darrel Marshall, Courtney B. Story, Bohkyung Kim, Ji-Young Lee, Robert Powers, Raul G. Barletta, and Patrick Dussault

Published in final edited form as:

ChemMedChem. 2014 August ; 9(8): 1838–1849. doi:10.1002/cmdc.201402067.

© 2014 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

Development of cyclobutene- and cyclobutane-functionalized fatty acids with inhibitory activity against *Mycobacterium tuberculosis*

Wantanee Sittiwong^[a], Denise K. Zinniel^[b], Robert J. Fenton^[b], Darrel Marshall^[a], Courtney B. Story^[a], Dr. Bohkyung Kim^[c], Prof. Ji-Young Lee^[c], Prof. Robert Powers^[a], Prof. Raúl G. Barletta^[b], and Prof. Patrick H. Dussault^[a]

Patrick H. Dussault: pdussault1@unl.edu

^[a]Department of Chemistry, University of Nebraska–Lincoln, Lincoln, NE 68588-0304

^[b]School of Veterinary Medicine and Biomedical Sciences, University of Nebraska–Lincoln, Lincoln, NE 68583-0905

^[c]Department of Nutritional Sciences, University of Connecticut, Storrs, CT 06269-4017

Abstract

Eleven fatty acid analogs incorporating four-membered carbocycles (cyclobutenes, cyclobutanes, cyclobutanones, and cyclobutanols) were investigated for the ability to inhibit growth of *Mycobacterium smegmatis* (*Msm*) and *Mycobacterium tuberculosis* (*Mtb*). A number of the analogs displayed inhibitory activity against both mycobacterial species in minimal media. Several of the molecules displayed potent levels of inhibition against *Mtb* with MIC values equal to or below those obtained with the anti-tuberculosis drugs D-cycloserine and isoniazid. In contrast, two of the analogs displaying the greatest activity against *Mtb* failed to inhibit *E. coli* growth under either set of conditions. Thus, the active molecules identified here (**1**, **2**, **6**, and **8**) may provide the basis for development of anti-mycobacterial agents against *Mtb*.

Keywords

cyclobutane; cyclobutene; fatty acid; mycobacteria; tuberculosis

Introduction

Tuberculosis resulting from *Mycobacterium tuberculosis* (*Mtb*) infection remains one of mankind's most significant disease threats. Based on skin test reactivity, it is estimated that one-third of the world's population has been exposed, resulting annually in approximately nine million incident cases and 1.4 million deaths (2010 data).^[1] A number of therapeutic agents have been developed, but current treatment regimens require patients to take multiple drugs over a period of months and are associated with significant side effects.^[2] The result is frequent patient noncompliance, leading to relapses and the emergence of drug resistance,

Correspondence to: Patrick H. Dussault, pdussault1@unl.edu.

Supporting information for this article is available on the WWW under <http://www.chemmedchem.org> or from the author

with a high fraction of active cases now involving multi-drug resistant (MDR, XDR) strains.^[1] There is, therefore, a need for new classes of therapeutics for tuberculosis.^[3] Importantly, the lethal target and/or the mechanism of action should also be novel to avoid established mechanisms of resistance and provide synergy with existing treatments.

Much of the hardness and drug resistance of mycobacteria is due to an unusually thick lipid cell wall containing a significant proportion of mycolic acids, a unique class of C₅₄–C₆₃ branched-chain fatty acids.^[4] A number of existing treatments for *Mtb*, exemplified by isoniazid and ethionamide, disrupt mycolic acid biosynthesis by inhibiting important biosynthetic enzymes.^[5] Mycobacteria incorporate intact C₁₆ and C₁₈ fatty acid skeletons as biosynthetic feed stocks and given the prior evidence for the incorporation of modified fatty acids into mycolic acid biosynthesis,^[6] we hypothesized that hijacking this pathway could provide a unique approach for the development of potential therapeutics.^[4] Towards this end, we proposed to investigate specifically functionalized fatty acids for their ability to limit mycobacterial growth.^[7] As our first steps towards validating this hypothesis, we investigated two series of fatty acid analogs (Figure 1). One series (**1–4**) was based upon a decenoic acid (10:1) framework and a second series (**5–11**) based upon the frameworks of oleic or elaidic acids (18:1). The analogs preserve the approximate lengths and cross-sections of the “parent” fatty acids while incorporating four-membered carbocycles into the backbone. We now report that several of these analogs display significant anti-mycobacterial activity against *Mtb*.

Results and Discussion

Substrate Preparation

The only previous description of molecules included in this study was the preparation of a mixture of the methyl esters of **5** and **10** via a nonselective radical decarboxylation.^[8] In search of an efficient, general, and stereospecific approach to incorporation of four-membered rings onto a fatty acid backbone, we were drawn to methodology for modification of dichlorocyclobutanones.^[9] Our approach is illustrated in detail for the preparation of analogs **1–4** (Scheme 1). The benzyl ester of 9-decenoic acid underwent cycloaddition with dichloroketene to afford one major dichlorobutanone (**15**),^[10] which was only moderately stable towards purification, and was therefore directly reduced with sodium borohydride to furnish a mixture of stereoisomeric 2,2-dichloro-1-cyclobutanols. The choice of sodium borohydride was for convenience as a similar product mixture is available using other reducing reagents such as BH₃·THF and Na(OAc)₃BH. The alcohols were converted to the corresponding methanesulfonate (mesylate) esters (**16**). Reaction with sodium in ammonia resulted in simultaneous reductive cleavage of the benzyl ester and fragmentation/reduction of the β-chloro methanesulfonate to furnish cyclobutene **1**. Hydrogenation of **1** cleanly generated cyclobutane **2**. Alternatively, dehalogenation of initial cycloadduct **15** with zinc in acetic acid, followed by deprotection of the benzyl ester with Pd(C)/H₂, furnished cyclobutanone **3**. Hydride reduction of the intermediate ketone with hydride prior to hydrogenolysis generated cyclobutanol **4** as a mixture of diastereomers at the newly created alcohol stereocenter.

The preparation of analogs incorporating the four-membered ring carbocycles onto a *cis* C₁₈ framework is illustrated in Scheme 2. In this case, the initial cycloaddition with benzyl oleate produced an inseparable mixture of regioisomeric dichlorocyclobutanones (**19**). The cycloadducts, which were prone to decomposition during purification, were directly reduced to furnish a mixture of regioisomeric 2,2-dichloro-1-cyclobutanols. The corresponding methanesulfonate (mesylate) esters (**20**) underwent reduction/fragmentation as described above to furnish the cyclobutene **5**,^[8] which could be hydrogenated to cyclobutane **6**. The related cyclobutanone (**7**) and cyclobutanol (**8**) were prepared from the dichlorocyclobutanone in a similar manner as described earlier for the C₁₀ series. Alternatively, controlled dehalogenation of the dichlorocyclobutanone cycloadduct with stoichiometric Zn or Zn(Cu) in acetic acid furnished a mixture of regioisomeric monochlorocyclobutanones (**22**) contaminated by small amounts of dichloroketone (starting material) and cyclobutanone **3** (overreduction).^[11] Hydrogenolysis of the monochlorocyclobutanone as before furnished analog **9**.

The ready availability of elaidic acid, a C₁₈ fatty acid containing a *trans*-9,10 alkene, made it possible to investigate the influence of stereoisomers (Scheme 3). Beginning with elaidic acid (**14**) we were able to synthesize cyclobutene **10** and monochlorocyclobutanone **11** in a manner analogous to that described previously for oleate-derived substrates **5** and **9**. The only major difference from the route employed in Scheme 2 was the need to employ activated Zn or Zn/Cu for the initial ketene generation/cycloaddition.^[12]

Inhibitor stability and solubility

Analog **1** – **11** were each tested for thermal stability using differential scanning calorimetry (DSC) in tandem with thermal gravimetric analysis (TGA); details are provided in the Supporting Information. Although we originally had concerns for the stability of the cyclobutene-containing analogs (**1**, **5**, and **10**) towards thermally induced cycloreversion,^[13] we observed no decomposition of any analog below 100 °C; most of the new analogs were stable to 150 °C.

One-dimensional (1D) ¹H NMR spectroscopy was used to verify the aqueous stability and solubility for each of the eleven analogs. An example series is illustrated in Figure 2; more detailed data is provided in Supporting Information. The results demonstrate the analogs to be stable in aqueous buffer. All of the analogs based upon a C₁₈ scaffold (**5**–**11**) display evidence of aggregation at all concentrations tested (down to 100 mM). In contrast, the analogs based upon the C₁₀ backbone (**1**–**4**) were not observed to aggregate at even mM concentrations (Table 1).

MIC determination

MICs were determined against the following bacteria: *Escherichia coli* (*E. coli*) wild type strain G58-1; *M. smegmatis* (*Msm*) strain mc²155, a non-pathogenic mycobacteria used as a model for *Mtb* to analyze processes that are likely to be conserved in the genus; and two *Mtb* strains (CDC1551 and H37Rv). Kanamycin (Kan) was used as a positive inhibition control for *E. coli*. Isoniazid (Inh), one of the main first-line drugs used to treat TB, and D-cycloserine (DCS), a second line clinically used TB drug, were employed as controls that

would allow benchmarking against other inhibitors. Results for the fatty acid analogs are shown in Table 2.

Neither of the two analogs (**1**, a C₁₀ cyclobutene and **6**, a C₁₈ cyclobutane) tested against *E. coli* were inhibitory in rich (LB broth) or minimal media. Interestingly, the susceptibility of *E. coli* towards the control drugs (DCS, Inh and Kan) was greater in minimal media as indicated by the significantly lower MIC values. Preliminary studies indicated that *Msm* was not susceptible to compounds **7** and **10** in complete Middlebrook 7H9 broth supplemented with either Tween or Tyloxapol (data not shown). In this context, a potential interfering compound is bovine serum albumin that has been shown to bind to fatty acids.^[14] Indeed, preliminary tests using *Msm* and compound **7** indicate that increasing concentrations of BSA significantly increase the MIC value (unpublished results). Thus, the entire set of compounds synthesized was tested in minimal media against *Msm* and the two *Mtb* strains. Under these conditions, none of the compounds demonstrated significant activity against *Msm*. However, four (**1**, **2**, **6**, and **8**) of the eleven compounds yielded MIC values < 100 µM for both *Mtb* strains, with two of the compounds (**2** and **8**) proving superior to DCS on a molar basis. Importantly, compound **2** yielded MIC values equal (CDC1551 strain) or superior (H37Rv strain) to those obtained with Inh. In summary, these compounds were poor inhibitors of *Msm* but quite effective against *Mtb* grown in minimal media.

Nonspecific toxicity

A sampling of six analogs, along with decanoic acid (**12**) and the *Mtb* therapeutics isoniazid and DCS, were investigated for toxicity in RAW 264.7 macrophages. SDS was employed as a positive control for maximum toxic effect. As illustrated in Table S1 (see Supporting Information), three of the tested analogs (**4**, **5**, and **6**), as well as DCS and isoniazid showed little or no toxicity at concentrations below 250 µM;^[15] toxicity for **12** was only slightly greater. Modest or significant toxicity was observed for analogs **1**, **2**, and **8**

Discussion

The MIC results demonstrate promising levels of inhibition with a variety of four-membered ring carbocycles (cyclobutanol, cyclobutanone, cyclobutene, and cyclobutane). A critically important component of evaluating new and potentially promising therapeutic entities is validation of a drug-like mode of action. Unfortunately, there are numerous undesirable mechanisms resulting from poor physical behavior of a compound that will result in a false positive in a biological assay. These poor properties may include insolubility, reactivity, micelle formation, impurities, aggregation, instability and non-specific binding. This has resulted in numerous compounds erroneously reported as chemical leads that are not acceptable drug candidates because of these undesirable physiochemical properties.^[16] The fatty acid analogs discussed here are thermally stable. All of the analogs based upon a C₁₈ backbone aggregate at concentrations overlapping the MIC values observed for *Mtb* or *Msm*. Correspondingly, the biological mechanism of action for analogs **5** to **11** is suspect and these analogs are unlikely to be viable drug-leads. The propensity for these compounds to form micelles or aggregate at concentrations lower than their MIC values must also be considered in any attempt to extract structure-activity relationships. In contrast, the analogs based upon a C₁₀ backbone (**1–4**) displayed no sign of aggregation at concentrations up to 1000 µM,

levels well above the measured MICs (Figure 3). The observation of significant and superior inhibition of *Mtb* with compounds from the shorter chain series provides strong evidence that the inhibition is specific and not simply due to the general toxicity often observed with very hydrophobic substrates.^[16] Although some short-chain fatty acids have also demonstrated limited antimicrobial activity,^[17] *Mtb* is also reported to use decanoic acid (10:0) as a carbon source^[17] Moreover, decenoic acid (**12**), the parent 10:1 framework of our shorter chain analogs, demonstrated very modest levels of inhibition in *Mtb*. Finally, we note that the extremely hydrophobic oleic acid (**13**), the parent 18:1 framework of analogs **5–9**, is used as a standard nutrient for *Mtb* at 0.05 g/L (177 μ M), a concentration higher than the MICs of the best inhibitors. Nonetheless, oleic acid is always used in media containing bovine serum albumin that may ameliorate its toxicity.^[14]

The nature of the functionality has a clear impact on activity. Within the less hydrophobic short-chain series (compounds **1–4**), the measured MICs vary by more than sixty fold against *Mtb* H37Rv and more than thirty fold against strain CDC1551. In contrast, less than a ten-fold difference is observed for MICs measured against *Msm*. Although the potential for micelle formation is cause for caution in making structure-activity comparisons involving the long chain series, the different influences of a given functional group within the C₁₈ and C₁₀ series are interesting. For example, the short-chain cyclobutanone (**3**) and cyclobutanol (**4**) are much less potent toward *Mtb* than the corresponding long-chain analogs (**7** and **8**), while the values for the cyclobutenes (**1** vs. **5** or **10**) are similar. The influence of stereochemistry varies by analog. The MICs for the *cis* and *trans* isomers of the C₁₈ monochloroketones (**9** and **11**) differ by more than ten-fold for *Mtb* (H37Rv); in contrast, no difference is seen between the *cis*- and *trans*-cyclobutenes (**5** and **10**) with either *Mtb* strain. Finally, we note that in both the C₁₈ and C₁₀ series, the saturated analogs (cyclobutanes **6** and **2**) are considerably more potent than the corresponding cyclobutenes (**5** and **1**).

Comparing MIC values obtained against *Mtb* vs. *Msm*, ten of eleven analogs demonstrate lower MIC values in at least one strain of *Mtb* compared with *Msm*; for six of the eleven, lower MIC values are obtained against both strains of *Mtb*. The ratio of MIC values for *Mtb* vs. *Msm* ranges from less than one to over six hundred; viewed across the eleven analogs, the average ratio of MICs for *Mtb* vs *Msm* is sixty eight (CDC1551) and nearly ninety (H37Rv). The different levels of inhibition produced in *Msm* and *Mtb* may result from a mechanism of action specific to *Mtb*. Isoniazid, for example, demonstrates 100-fold more potent inhibition of *Mtb* compared with *Msm* and possesses low toxicity toward other mycobacteria and prokaryotic pathogens.^[21] The lack of inhibition observed in *E. coli* using two of the analogs (**1** and **6**) most potent against *Mtb* could indicate specific inhibition of a metabolic pathway unique to the latter.

The initial hypothesis behind the design of the molecules reported here was that uptake of a fatty acid analog bearing a reactive functional group (for example, cyclobutene, cyclobutanone, chloroketone) might disrupt an enzymatic process related to cell wall synthesis. We imagined that one such target might be methyl transferases, which are common in mycobacteria, but relatively uncommon in other organisms.^[18] In an earlier example of a related approach, the inhibition of growth in *Mtb* observed in the presence of alkyne-containing fatty acids (MICs of 20–25 μ M) was attributed to irreversible

modification of an enoyl reductase within the mycobacterial FASII system.^[4,7] However, the significant inhibition produced by cyclobutanes **2** and **6**, which lack a chemically reactive group, is more suggestive of a noncovalent mode of inhibition. The relatively low MIC values indicate our fatty acid analogs have promise as anti-mycobacterial agents, but there is still a need to confirm cellular uptake and to establish the *in vivo* target.

A variety of carbocyclic fatty acids including cyclopropyl, cyclopentyl, cyclohexyl and cycloheptyl fatty acids are found in bacteria and plants.^[19] Their roles, while not always completely understood, may be related to control of membrane fluidity.^[20] In the case of mycobacteria, the presence of cyclopropane subunits in mycolic acids is known to be associated with the structural integrity of the cell wall and the ability of the tubercle bacillus to resist oxidative stress inside macrophages. In contrast, cyclobutane fatty acids are rare. An unusual fatty acid incorporating a ladder-type structure of fused cyclobutanes, recently isolated from anammox bacteria found near undersea vents, is believed to contribute to the formation of dense and impermeable lipid membranes that protect the bacteria from harsh chemical environments.^[19] Thus, as an alternative to disrupting mycolic acids biosynthesis, it is also plausible that our four-membered ring analogs are converted into analogs of mycolic acids and that the physical properties of these unnatural cell wall constituents results in cell death. However, it must be noted that while mycobacteria are known to directly incorporate structurally modified fatty acid feedstocks into mycolic acid biosynthesis,^[6] there is as yet no evidence that the same is true for the specific fatty acid analogs described here.^[7] Indeed, further studies will be required in order to determine an *in vivo* mechanism of inhibition for our compounds.

In this context, it is worth noting that our initial choice of molecular scaffold was based upon the availability of low-cost precursors. If the modified fatty acid chains are indeed taken up into the mycobacterial biosynthetic pathways, greater activity or different specificities might be identified through the use of different molecular backbones or alternative positioning of the four-membered ring carbocycles.

Conclusion

Analogues of fatty acids incorporating four-membered ring carbocycles may provide a starting point for the development of new antimycobacterials. Four of eleven analogs tested yielded MIC values < 100 μ M for both *Mtb* strains, with two of the analogs proving more active than DCS and one analog proving equally potent or superior to Inh, one of the main first-line drug used for treatment of tuberculosis.

Experimental Section

General procedures

All reagents and solvents were used as purchased except for pyridine and CH₂Cl₂ (distilled from CaH₂ and kept under N₂) and THF (distilled from Na/benzophenone under N₂). Reactions were performed under a N₂ atmosphere unless stated otherwise. Thin-layer chromatography (TLC) was performed on 0.25 mm hard-layer silica G plates; developed plates were visualized with a hand-held UV lamp or by staining: 1% Ce(SO₄)₂ and 10%

(NH₄)₂MoO₄ in 10% aq. H₂SO₄ (general stain, after heating); 1% aq. KMnO₄ (for unsaturated compounds); 3% vanillin in 3% H₂SO₄ in EtOH (general stain, after heating). Unless otherwise noted, NMR spectra were recorded at 400 MHz (¹H) or 100 MHz (¹³C) in CDCl₃; peaks are reported as: chemical shift (multiplicity, *J* couplings in Hz, number of protons); “app” and “br” refer to apparent and broad signals, respectively. IR spectra were recorded as neat films (ZnSe, ATR mode) with selected absorbances reported in wavenumbers (cm⁻¹). Flash chromatography was performed on 32–60 μm silica gel. Preparative HPLC was performed on a 21 × 250 mm normal phase Si (8 micron) column at 10 mL/min of the indicated solvent unless otherwise noted. Analytical purity of compounds was checked using an analytical column (250 mm × 4.6 mm; Microsorb) at 1 mL/min of 20% EtOAc/Hexane; detection was accomplished using a differential refractometer interfaced with a data module. All compounds tested for biological activity showed >97% purity by HPLC analysis except for **5** and **7**. Melting points are uncorrected.

3-(Octanoic acid, 8-yl, benzyl ester) 2,2-dichlorocyclobutanone (15) was prepared similarly to compound **23** and afforded (13.8 g, 80%) as a brown oil from benzyl 9-decenoate (12.4 g, 48 mmol), trichloroacetyl chloride (10.7 mL, 95 mmol), and Zn(Cu) (15.55 g, 238 mmol) in ether 0.1 M (340 mL). Consistent with a previous report, only one isomer was observed.^[22] The assignment was confirmed by observation of the pair of geminal hydrogens adjacent to the ketone at δ 3.33 and 2.94 by ¹H-NMR and by the observation of eleven rather than twelve carbons (symmetry) upon dechlorination to cyclobutanone **3**: *R*_f = 0.34 (10% EtOAc/Hex); IR: 2935, 2852, 1814, 1730, 905, 725 cm⁻¹; ¹H NMR: δ 7.39–7.29 (m 5H), 5.12 (s, 2H), 3.33 (dd, *J* = 9.4, 17.2, Hz, 1H), 2.94 (dd, *J* = 9.0, 17.2 Hz, 1H), 2.88–2.81 (m, 1H), 2.37/2.36 (overlapping t, *J* = 7.5 Hz, 2H), 1.94–1.84 (m, 1H), 1.70–1.72 (m, 3H), 1.46–1.28 (m, 8H); ¹³C NMR: δ 192.8, 173.4, 136.0, 128.4, 128.0, 88.8, 65.9, 47.7, 45.8, 34.1, 31.2, 29.0, 28.9, 28.8, 27.2, 24.7; HREI-MS calcd. for C₁₉H₂₅O₃Cl₂: [M+H]⁺ 371.1181; found: 371.1188.

3-(Octanoic acid, 8-yl, benzyl ester) 2,2-dichlorocyclobutanol, methanesulfonate ester (16) was prepared similarly to compound **20** from the previous dichloroketone **15** (10.0 g, 26.9 mmol), NaBH₄ (1.53 g, 40.4 mmol), and *i*PrOH (350 mL). The crude product was purified by flash column chromatography with 10% EtOAc/Hex to furnish the dichlorobutanol (5.76 g, 57%) as a colorless oil: *R*_f = 0.39 (20% EtOAc/Hex); IR: 3440 (br), 2927, 2854, 1733, 1164, 960, 697 cm⁻¹; ¹H NMR: δ 7.40–7.30 (m 5H), 5.12 (s, 2H), 4.73–4.64 (m, 0.1H) 4.34–4.27 (m, 0.9H), 2.97 (d, *J* = 8.2 Hz, 0.1H, *OH* is *trans* to backbone), 2.87 (d, *J* = 10.3 Hz, 0.9H, *OH* is *cis* to backbone), 2.48–2.34 (m, 1H), 2.36 (t, *J* = 7.6 Hz, 2H), 1.78–1.60 (m, 3H), 1.53–1.28 (m, 11H); ¹³C NMR: δ 173.7, 136.0, 128.5, 128.1, 92.9, 75.4, 66.1, 45.6, 34.5, 34.2, 29.8, 29.2, 29.0, 28.9, 26.5, 24.8; HRESI-MS calcd. for C₁₉H₂₃O₃Cl₂ [M+Na]⁺: 395.1157; found: 395.1151.

Conversion to the methanesulfonate was performed as described for compound **20**, using methanesulfonyl chloride (20 mL, 29.4 mmol), the dichlorobutanol (5.50 g, 14.7 mmol), and triethylamine (9.20 mL, 66.3 mmol) in CH₂Cl₂ (70 mL). The crude product was purified by flash chromatography (10% EtOAc/Hex) to afford **16** as a light yellow oil (6.13 g, 92%): *R*_f = 0.36 (20% EtOAc/Hex); IR: 2927, 2858, 1731, 1364, 1179, 952 cm⁻¹; ¹H NMR: δ 7.36–7.31 (m 5H), 5.11 (s, 2H), 5.13–5.09 (m, 1H), 3.18 (s, 3H), 2.56–2.50 (m, 2H), 2.35 (t, *J*

= 7.6 Hz, 2H), 1.93-1.28 (m, 13H); ^{13}C NMR: δ 173.5, 136.0, 128.4, 128.1, 88.4, 78.3, 66.0, 45.8, 39.3, 34.1, 31.6, 29.8, 29.1, 28.87, 28.85, 26.2, 24.8; HRESI-MS calcd. for $\text{C}_{20}\text{H}_{28}\text{O}_5\text{Cl}_2\text{SNa}$ $[\text{M}+\text{Na}]^+$: 473.0932; found: 473.0930.

8-(2-Cyclobuten-1-yl)-octanoic acid (1) was prepared by a similar procedure as employed for **5** via reaction of Na metal (2.80 g, 120 mmol) with a solution of 5.40 g (12.0 mmol) of the mixture of methanesulfonate **16** in THF (30 mL) and NH_3 (~ 250 mL). The crude product was purified by flash column chromatography (2% EtOAc/Hex) to afford 9,10-ethenooctadecanoic acid **1** (1.64 g, 52 %) as a low-melting white solid: R_f = 0.47 (6:4:0.05 of Hex:EtOAc:AcOH); m.p. = 20–21 °C; IR: 3112 (br), 3043, 2920, 2853, 1705, 697 cm^{-1} ; ^1H NMR: δ 10.27 (br s, 1H, COOH), 6.11 (br d, J = 2.4 Hz, 1H), 6.04 (br dd, J = 0.8, 2.0 Hz, 1H), 2.77 (m, 1H), 2.66 (ddd, J = 0.8, 4, 13.2 Hz, 1H), 2.35 (t, J = 7.2 Hz, 2H), 2.04 (d, J = 13.2 Hz, 1H), 1.64 (quintet, J = 7.2 Hz, 2H), 1.51-1.38 (br m, 2H), 1.38-1.23 (br m, 8H); ^{13}C NMR: δ 180.0, 141.1, 135.1, 44.2, 36.9, 34.6, 34.0, 29.5, 29.2, 29.0, 27.9, 24.7; HRESI-MS calcd. for $\text{C}_{12}\text{H}_{20}\text{O}_2\text{Na}$ $[\text{M}+\text{Na}]^+$: 219.1361; found: 219.1355. Purity 100% by HPLC (retention time, 5.01 min).

8-Cyclobutyloctanoic acid (2) was prepared by a similar procedure as for **6** from the hydrogenation of cyclobutene **1** (0.034, 0.170 mmol), over 10% Pd/C (0.034 g, 0.030 mmol) in EtOAc (1.7 mL) to afford the unprotected acid **2** (0.030 g, 89%) as a low-melting white solid: R_f = 0.47 (6:4:0.05 of Hex:EtOAc:AcOH); m.p. = 28–30 °C; IR: 3035 (br), 2916, 2849, 1695, 909, 733 cm^{-1} ; ^1H NMR: δ 11.34 (br s, 1H, COOH), 2.37 (t, J = 7.2, 2H), 2.25 (septet, J = 7.2, 1H), 2.10-1.98 (m, 2H), 1.90-1.75 (m, 2H), 1.70-1.52 (m, 2H), 1.40-1.12 (br m, 10H); ^{13}C NMR: δ 180.3, 37.0, 36.2, 34.1, 29.4, 29.2, 29.0, 28.4, 27.1, 24.7, 18.5; HRCI-MS calcd. for $\text{C}_{12}\text{H}_{22}\text{O}_2$: $[\text{M}+\text{H}]^+$ 199.1698; found: 199.1695. Purity > 99% by HPLC (retention time, 4.92 min).

3-(Octanoic acid benzyl ester, 8-yl) cyclobutanone (17) was prepared in a similar manner as **21**, via reaction of dichloroketone **15** (1.05 g, 2.69 mmol), 6 mL of glacial acetic acid, and Zn(Cu) (1.76 g, 26.9 mmol). The product was purified by flash chromatography (10% EtOAc/Hex) to afford a colorless oil (0.4859 g, 59%) which was a mixture of regioisomers: R_f = 0.55 (20% EtOAc/Hex). IR: 2923, 2854, 2780, 2732, 1163, 730, 697 cm^{-1} ; ^1H NMR: δ 7.35-7.28 (br m, 5H), 5.09 (s, 2H), 3.13-3.03 (m, 2H), 2.65-2.57 (m, 2H), 2.33 (t, J = 7.2, 2H), 2.33-2.26 (m, 1H), 1.68-1.58 (br quintet, J = 6.4, 2H), 1.58-1.48 (br q, J = 7.2, 2H), 1.29 (br s, 8H); ^{13}C NMR: δ 208.2, 173.3, 134.0, 128.3, 127.9, 65.8, 52.3, 36.1, 34.0, 28.9, 28.8, 28.0, 24.7, 23.6; HRESI-MS calcd. for $\text{C}_{19}\text{H}_{26}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$: 325.1780; found: 325.1775.

3-(Octanoic acid, 8-yl) cyclobutanone (3) was prepared by a similar procedure as for **7** through reaction of benzyl ester **17** (0.0346, 0.1 mmol), 10% Pd/C (0.020 g, 0.02 mmol) and EtOAc (1 mL). The product was purified by flash column chromatography on silica gel (step gradient of 20%/40% EtOAc/Hex) to afford acid **3** (0.0189 g, 90%) as a low-melting white solid: R_f = 0.3 (6:4:0.05 of Hex:EtOAc:AcOH); m.p. = 35–37 °C; IR: 3094 (br), 2923, 2853, 1780, 1705 cm^{-1} ; ^1H NMR: δ 10.61 (br s, 1H, COOH), 3.17-3.08 (m, 2H), 2.69-2.61 (m, 2H), 2.35 (br t, J = 7.6, 2H), 2.39-2.31 (br, 1H), 1.69-1.53 (br m, 4H), 1.32 (br s, 8H); ^{13}C NMR: δ 208.8, 179.9, 52.5, 36.3, 34.0, 29.2, 29.1, 28.9, 28.2, 24.6, 23.8; HRCI-MS calcd.

for $C_{12}H_{20}O_3Na$ $[M+Na]^+$: 213.1491; found: 213.1490. Purity > 97% by HPLC (retention time, 10.75 min).

3-(Octanoic acid benzyl ester) cyclobutanol (18) was prepared by a similar procedure as compound **8** via reaction of cyclobutanone **17** (0.0509 g, 0.17 mmol) and $NaBH_4$ (0.013 g, 0.34 mmol) in anhydrous MeOH (2.5 mL). **18** (after purification by flash column chromatography with 20% EtOAc/Hex) was isolated as a low-melting white solid (0.040 g, 76%). The product was a 9:1 mixture of *cis/trans* stereoisomers based on the previous assignments for cyclobutanol **8**. R_f = 0.29 (20% EtOAc/Hex); m.p. = 30–31 °C; IR: 3356 (br), 2922, 1734, 1154, 735, 696 cm^{-1} ; 1H NMR: δ 7.40–7.30 (m, 5H), 5.11 (s, 2H), 4.42–4.34 (br m, 0.1H), 4.14–4.03 (br m, 0.9H), 2.48–2.40 (m, 0.18H), 2.35 (t, J = 7.2, 0.18H), 2.16–2.04 (m, 0.2H), 1.95 (t, J = 6.6, 0.2H), 1.70–1.56 (br m, 3H), 1.48–1.13 (br m, 12H); ^{13}C NMR: δ 173.6, 136.0, 128.4, 128.1, 66.0, 63.7, 39.7, 37.0, 34.2, 29.2, 29.1, 29.0, 27.3, 25.4, 24.82; HRCI-MS calcd. for $C_{19}H_{29}O_3Na$ $[M+Na]^+$: 305.2117; found: 305.2116.

3-(Octanoic acid) cyclobutanol (4) was prepared by a similar procedure as employed for **8** through reaction of benzyl ester **18** (0.034g, 0.110 mmol), 10% Pd/C (0.024 g, 0.022 mmol) and EtOAc (1 mL). The crude product was purified by flash column chromatography (40% EtOAc/Hex) to afford acid **4** (0.013 g, 54%) as a white solid: R_f = 0.24 (6:4:0.05 of Hex:EtOAc:AcOH); m.p. = 74–75 °C; IR: 3341 (br), 2916, 2849, 1695, 1064 cm^{-1} ; 1H NMR (MeOD): δ 4.30 (quintet, J = 6.8, 0.1H), 4.01 (quintet, J = 6.4, 0.9H), 2.45–2.35 (br m, 2H), 2.29 (t, J = 7.2, 2H), 1.75–1.55 (br m, 3H), 1.50–1.15 (br m, 12H); ^{13}C NMR (MeOD): δ 176.3, 62.8, 38.9, 36.9, 36.8, 33.6, 29.1, 29.0, 28.8, 27.1, 25.4, 24.7; HRESI-MS calcd. for $C_{12}H_{22}O_3Na$ $[M+Na]^+$: 237.1467 found: 237.1460. Purity > 97% by HPLC (retention time, 9.65 and 11.36 min).

2,2-Dichloro-4-(octanoic acid benzyl ester, 8-yl) 3-octylcyclobutanone-; 2,2-Dichloro-3-(octanoic acid benzyl ester, 8-yl) 4-octylcyclobutanone (19) was prepared by a modification of a known procedure.^[22–24] To a mixture of zinc dust (0.879 g, 13.4 mmol) and benzyl oleate **19** (1.02 g, 2.73 mmol) in 20 mL of anhydrous ether was slowly added a solution of trichloroacetyl chloride (0.75 mL, 6.8 mmol) in 7 mL of diethyl ether via dropping funnel over a period of 45 min. The reaction was stirred until judged complete by the absence of starting material (4 h, TLC). The reaction mixture was filtered through a plug of Celite and the residue was washed with ether (2 × 75 mL). The organic solution was stirred a minimum with 50% aq. $NaHCO_3$ and the separated aqueous layer back-extracted with ether (2 × 20 mL). The combined organic layers were dried over Na_2SO_4 and concentrated under vacuum to give g of the product (1.30 g, 99%) as a yellow oil. The unpurified material was generally used directly for the next reaction because of a tendency to decompose during chromatography. However, small quantities could be purified by rapid flash chromatography (5% EtOAc/Hex) to afford a 1:1 mixture of regioisomeric dichlorocyclobutanones **19** as a colorless oil: R_f = 0.53 (10% EtOAc/Hex); IR: 2926, 2855, 1800, 1735, 1456, 1162, 696 cm^{-1} ; 1H NMR: δ 7.36–7.32 (5H), 5.120/5.116 (overlapping s, 2H), 3.55–3.49 (m, 1H), 2.97–2.93 (m, 1H), 2.37/2.35 (overlapping t, J = 7.6 Hz, 2H), 1.74–1.62 (m, 4H), 1.66–1.48 (m, 4H), 1.45–1.27 (br m, 18H), 0.89/0.88 (overlapping triplets, J = 6.9 Hz, total 3H); ^{13}C NMR: δ 197.5, 173.5, 136.1, 128.5, 128.1, 88.1, 66.0, 58.0, 57.9, 49.4, 49.4, 34.2, 31.8, 31.80, 29.60, 29.3, 29.3, 29.2, 29.18, 29.15, 29.12, 29.0,

28.9, 28.9, 28.04, 27.96, 26.72, 26.68, 26.3; HRFAB-MS (3-NBA+Li matrix: calcd. for $C_{27}H_{40}O_3Cl_2Li$ $[M+Li]^+$: 489.2515; found: 489.2499.

2,2-Dichloro-4-(octanoic acid benzyl ester, 8-yl) 3-octylcyclobutanol, methanesulfonate ester; 2,2-Dichloro-3-(octanoic acid benzyl ester, 8-yl) 4-octylcyclobutanol, methanesulfonate ester (20). Reduction of the dichlorocyclobutanone **19** was achieved using a modification of a known procedure.^[9,24] To a 0 °C solution of dichlorocyclobutanones **19** (1.5 g, 3.1 mmol) in *i*PrOH (45 mL) was added $NaBH_4$ (0.176 g, 4.65 mmol). The reaction was allowed to slowly warm to rt and stirred until starting material had disappeared (TLC, ~ 2.5h; byproducts tended to accumulate upon prolonged reaction). The reaction was then cooled to 0 °C and quenched with 1N HCl (40 mL). The solution was stirred for an additional 30 min, concentration to one third the original volume, and then extracted with EtOAc (3 × 100 mL). The organic layer was washed sequentially with water/sat. $NaHCO_3$ /brine and then dried over Na_2SO_4 . Evaporation of the organic solvent and flash chromatography of the residue on silica gel, using a step gradient of 5%/10% EtOAc/Hex, gave (0.821 g, 55% yield) of a 1:1 mixture of regioisomeric cyclobutanols as a colorless oil. The products were predominantly a regioisomeric mixture of the *cis,cis* diastereomers^[22] based on 1H NMR, HSQC, 2D-COSY and 2D-NOESY analysis:^[25] R_f (10% EtOAc/Hex) = 0.33 (major, OH is *cis* to backbone) 0.28 (minor, OH is *trans* to backbone), 0.2 (major, OH is *cis* to backbone) and 0.15 (minor, OH is *trans* to backbone); IR: 3473, 3036, 2924, 2854, 1733, 1456, 1168, 696 cm^{-1} ; 1H NMR: δ 7.36-7.32 (m, 5H), 5.116/5.112 (app s, total 2H), 4.55 (dd, J = 10.8, 8.8 Hz, 0.92H, CH-OH, *cis* to the backbone assigned based on 2D-NOESY analysis and $^4J_{diagonal\ H} = 0$), 3.99 (br t, J = 10 Hz, 0.08H), 2.75-2.69 (m, 1H), 2.64/2.62 (overlapping d, J = 10.8 Hz, total 1H, -OH proton assigned based on the disappearance of this peak upon D_2O addition), 2.60-2.55 (m, 1H), 2.36/2.35 (overlapping t, J = 7.6 Hz, total 2H), 1.67-1.62 (bm, 3H), 1.52-1.19 (24H), 0.89/0.88 (overlapping t, J = 6.8 Hz, total 3H). Small quantities of the individual isomers could be isolated by semipreparative HPLC (10% EtOAc/Hex, RI detection):

First-eluting *cis*— R_f = 0.33; IR: 3457, 2924, 2854, 1736, 1456, 1170, 697 cm^{-1} ; 1H NMR: δ 7.37-7.32 (m, 5H), 5.12 (s, 2H), 4.55 (dd, J = 10.8, 8.8 Hz, 1H, CH-OH, *cis* to the backbone assigned based on 2D-NOESY analysis and $^4J_{diagonal\ H} = 0$), 2.72 (dt, J = 9.2, 5.6 Hz, 1H), 2.59 (d, J = 10.8 Hz, 1H, -OH), 2.55 (app dq, J = 9.2, 5.6 Hz, 1H), 2.36 (t, J = 7.6 Hz, 2H), 1.67-1.51 (br m, 3H), 1.52-1.13 (m, 24H), 0.88 (app t, J = 6.8 Hz, 3H); ^{13}C NMR: δ 173.6, 136.1, 128.5 (two overlapping signals), 128.2 (two overlapping signals), 93.5, 77.2, 66.1, 49.5, 40.9, 34.3, 31.8, 29.9, 29.4, 29.4, 29.3, 29.2, 29.0, 27.0, 25.7, 24.9, 23.2, 22.7, 14.1; HRFAB-MS (3-NBA matrix): calcd. for $C_{27}H_{43}O_3Cl_2$ $[M+H]^+$: 485.2589; found: 485.2572.

First-eluting *trans*— R_f = 0.28; IR: 3449, 3028, 2920, 2850, 1736, 1456, 1157, 696 cm^{-1} ; 1H NMR: δ 7.37-7.32 (5H), 5.12 (s, 2H), 3.99 (ddd, J = 10.0, 8.4, 1.2 Hz, 1H, CH-OH, *trans* to the backbone assigned based on 2D-NOESY analysis and $^4J_{diagonal\ H} = 1.2$ Hz), 2.68-2.64 (m, 1H), 2.52 (d, J = 10.0 Hz, 1H, -OH), 2.34 (t, J = 7.6 Hz, 1H), 2.29-2.20 (m, 1H), 1.67-1.26 (br m, 27H), 0.88 (app t, J = 6.8 Hz, 3H); ^{13}C NMR: δ 173.6, 136.1, 128.5 (two overlapping signals), 128.2 (two overlapping signals), 91.1, 82.6, 66.1, 50.3,

44.4, 34.3, 31.9, 29.7, 29.5, 29.2, 29.04, 29.02, 28.4, 28.23, 28.15, 27.7, 24.9, 22.7, 14.1; ESI-MS calcd. for $C_{27}H_{42}O_3Cl_2Na$ $[M+Na]^+$: 507.2409; found: 507.2422.

Second-eluting cis— R_f = 0.20; IR: 3464, 2924, 2854, 1736, 1456, 1169, 696 cm^{-1} ; 1H NMR: δ 7.37–7.32 (m, 5H), 5.11 (s, 2H), 4.55 (dd, J = 10.8, 8.8 Hz, 1H CH-OH, *cis* to the backbone assigned based on 2D-NOESY analysis and $^4J_{diagonal\ H} = 0$), 2.72 (dt, J = 9.2, 5.6 Hz, 1H), 2.65 (d, J = 10.8 Hz, 1H, -OH), 2.54 (app dq, J = 9.2, 5.6 Hz, 1H), 2.35 (t, J = 7.2 Hz, 2H), 1.68–1.60 (bm, 3H), 1.52–1.13 (m, 24H), 0.89 (app t, J = 6.8 Hz, 3H); ^{13}C NMR: δ 173.7, 136.1, 128.5 (two overlapping signals), 128.2 (two overlapping signals), 93.5, 77.2, 66.1, 49.5, 40.9, 34.3, 31.8, 29.6, 29.4, 29.2, 29.1, 29.04, 29.00, 27.0, 25.8, 24.8, 23.2, 22.7, 14.1; HRFAB-MS (3-NBA matrix): calcd. for $C_{27}H_{43}O_3Cl_2$ $[M+H]^+$: 485.2589; found: 485.2581

Second-eluting trans— R_f = 0.15; IR: 3464, 3033, 2925, 2854, 1738, 1498, 1160, 697 cm^{-1} ; 1H NMR: δ 7.37–7.32 (m, 5H), 5.11 (s, 2H), 3.99 (ddd, J = 10.0, 8.4, 1.2 Hz, 1H, CH-OH, *trans* to the backbone assigned based on 2D-NOESY analysis and $^4J_{diagonal\ H} = 1.2$ Hz), 2.69–2.64 (m, 1H), 2.56 (d, J = 10.0 Hz, 1H, -OH), 2.35 (t, J = 7.6 Hz, 2H), 2.26–2.22 (m, 1H), 1.66–1.26 (m, 27H), 0.88 (app t, J = 6.8 Hz, 3H); ^{13}C NMR: δ 173.7, 136.1, 128.5 (two overlapping signals), 128.2 (two overlapping signals), 91.1, 82.6, 66.1, 50.3, 44.3, 34.3, 31.8, 29.7, 29.4, 29.2, 29.04, 28.97, 28.5, 28.21, 28.18, 27.5, 24.8, 22.7, 14.1; HRESI-MS calcd. for $C_{27}H_{42}O_3Cl_2Na$ $[M+Na]^+$: 507.2409; found: 507.2426.

The mixture of regioisomeric cyclobutanols converted to the methanesulfonate using a variant of published procedures.^[9,24] Methanesulfonyl chloride (62 mL, 0.80 mmol) was slowly added to a stirred 0 °C solution of the cyclobutanols (0.193 g, 0.398 mmol) and triethylamine (0.25 mL, 8.0 mmol) in CH_2Cl_2 (2 mL) over 15 min. The reaction was allowed to slowly warm to rt and was stirred for 17 h or until no starting material remained (TLC). The reaction was diluted with CH_2Cl_2 (75 mL) and washed sequentially with 1N HCl (2 × 25 mL), sat aq. $NaHCO_3$ (2 × 25 mL) and water (2 × 25 mL). The organic solution was then dried with Na_2SO_4 and concentrated under reduced pressure. The residue was purified by flash chromatography using 10% EtOAc/Hex to give **20** (0.206 g, 92%) as a light yellow oil consisting of a 1:1 mixture of regioisomers, each a 92:8 mixture of *cis/trans* stereoisomers based upon 1H NMR: R_f (10% EtOAc/Hex) = 0.31 (first *trans*), 0.26 (first *cis*/second *trans*), 0.21 (second *cis*); IR: 3032, 2925, 2855, 1735, 1458, 1180, 696 cm^{-1} ; 1H NMR: δ 7.36–7.33 (m, 5H), 5.33 (d, J = 9.6 Hz, 0.92H), 5.114/5.109 (two s, total 2H), 4.84 (d, J = 7.6 Hz, 0.08H), 3.208/3.205 (two s, total 3H), 2.76–2.74 (m, 2H), 2.36/2.34 (two overlapping t, J = 7.6 Hz, total 2H), 1.66–1.60 (m, 4H), 1.46–1.26 (m, 22H), 0.88/0.87 (overlapping t, J = 6.8 Hz, total 3H). Small quantities of individual isomers could be separated for analysis by HPLC using 10% EtOAc/Hex. The stereochemical assignments were confirmed by 2D-NMR experiments and by preparation of individual methanesulfonates from individual stereoisomers of alcohol **19**, prepared as described in the previous step.

First eluting trans— R_f = 0.31; IR: 3032, 2926, 2855, 1737, 1733, 1456, 1180, 697 cm^{-1} ; 1H NMR: δ 7.37–7.32 (5H), 5.12 (s, 2H), 4.84 (d, J = 7.6 Hz, 1H), 3.20 (s, 3H),

2.74-2.68 (m, 2H), 2.36 (t, $J = 7.6$ Hz, 2H), 1.67-1.32 (br m, 26H), 0.88 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR: δ 173.6, 136.1, 128.5 (two overlapping signals), 128.2 (two overlapping signals), 86.4, 85.8, 66.1, 50.5, 41.1, 39.5, 34.3, 31.8, 29.5, 29.4, 29.3, 29.2, 29.01, 28.98, 28.3, 28.1, 27.6, 27.1, 24.9, 22.6, 14.1; HRESI-MS calcd. for $\text{C}_{28}\text{H}_{44}\text{O}_5\text{Cl}_2\text{S}$ $[\text{M}+\text{Na}]^+$: 585.2184; found: 585.2188.

Second-eluting trans— $R_f = 0.26$; IR: 3040, 2927, 2855, 1734, 1456, 1179, 697 cm^{-1} ; ^1H NMR: δ 7.37-7.32 (m, 5H), 5.11 (s, 2H), 4.84 (dd, $J = 8.4, 0.8$ Hz, 1H), 3.19 (s, 3H), 2.76-2.65 (m, 2H), 2.35 (t, $J = 7.6$ Hz, 2H), 1.67-1.32 (26H), 0.89 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR δ 173.6, 136.1, 128.5 (two overlapping signal), 128.2 (two overlapping signal), 86.4, 85.7, 66.1, 50.5, 41.1, 39.5, 34.3, 31.8, 29.6, 29.31, 29.25, 29.2, 29.0, 28.3, 28.1, 27.6, 27.0, 24.9, 22.7, 14.1; HRESI-MS; calcd. for $\text{C}_{28}\text{H}_{44}\text{O}_5\text{Cl}_2\text{S}$ $[\text{M}+\text{Na}]^+$: 585.2184; found: 585.2205.

First-eluting cis— $R_f = 0.26$; IR: 2926, 2855, 1736, 1456, 1182, 697 cm^{-1} ; ^1H NMR: δ 7.37-7.32 (m, 5H), 5.33 (app d, $J = 9.6$ Hz, 1H), 5.11 (s, 2H), 3.19 (s, 3H), 2.80-2.72 (bm, 2H), 2.36 (app t, $J = 7.6$ Hz, 2H), 1.70-1.63 (m, 4H), 0.87 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR: δ 173.6, 136.0, 128.5 (two overlapping signal), 128.1 (two overlapping signal), 88.9, 80.7, 66.0, 49.5, 40.2, 39.2, 34.2, 31.8, 29.7, 29.3, 29.3, 29.2, 29.0, 28.8, 26.7, 25.7, 24.8, 23.7, 22.6, 14.1; HRFAB-MS (3-NBA matrix): calcd. for $\text{C}_{28}\text{H}_{45}\text{O}_5\text{Cl}_2\text{S}$ $[\text{M}+\text{H}]^+$: 563.2365; found: 563.2349.

Second-eluting cis— $R_f = 0.21$; IR: 2925, 2855, 1735, 1457, 1181, 697 cm^{-1} ; ^1H NMR: δ 7.36-7.32 (m, 5H), 5.33 (app d, $J = 9.6$ Hz, 1H), 5.11 (s, 2H), 3.20 (s, 3H), 2.80-2.69 (bm, 2H), 2.34 (app t, $J = 7.6$ Hz, 2H), 1.74-1.60 (4H), 1.50-1.19 (22H); ^{13}C NMR: δ 173.6, 136.1, 128.5 (two overlapping signal), 128.1 (two overlapping signal), 89.0, 80.7, 66.0, 49.5, 40.2, 39.3, 34.2, 31.8, 29.5, 29.5, 29.3, 29.2, 28.9, 28.9, 28.7, 26.8, 25.7, 24.8, 23.7, 22.6, 14.1; HRFAB-MS (3-NBA matrix): calcd. for $\text{C}_{28}\text{H}_{45}\text{O}_5\text{Cl}_2\text{S}$ $[\text{M}+\text{H}]^+$: 563.2365; found: 563.2360.

1-(Octanoic acid, 8-yl)-2-octylcyclobutene (5) was prepared from the regioisomeric mixture of dichloromethanesulfonates **20** using a reported procedure.^[9,24] NH_3 (~ 15 mL, liquid) was condensed into a 100 mL three-necked round bottom flask fitted with a dry ice/acetone condenser. Sodium (0.113 g, 4.90 mmol, sliced under Hex) was added in small pieces. The resulting deep blue solution was stirred for 20 min. A solution of 0.276 g (0.49 mmol) of **20** in anhydrous THF (1.5 mL) was added over 10 min. The cooling bath was removed and the reaction was allowed to stir at ~ -35 $^{\circ}\text{C}$ (refluxing NH_3) 5–30 min (depending on the scale of preparation). The reaction was then recooled to -78 $^{\circ}\text{C}$ (dry ice/acetone) and stirred for 2 h or until the starting material was consumed by TLC. Saturated aq. NH_4Cl was slowly added until the blue color was no longer visible. The condenser was removed and the reaction mixture was slowly allowed to warm to 0 $^{\circ}\text{C}$ with evaporation of NH_3 . The remaining reaction mixture was diluted with 30 mL of water and the suspension was extracted with CH_2Cl_2 (3×50 mL). The organic layers were combined and washed sequentially with 1N aq. HCl /water/saturated aq. NaHCO_3 /water/brine. The organic solution was then dried with NaSO_4 , and concentrated under reduced pressure. The residue was

purified by flash column chromatography using 10% EtOAc/Hex to afford the cyclobutene carboxylic acid **5** (0.131 g, 87%) as a low-melting white solid: R_f = 0.50 (80:20:1 of Hex:EtOAc:AcOH); mp = 33–35 °C; IR: 3493, 3036, 2923, 2853, 1708, 1467, 1275, 728 cm^{-1} ; ^1H NMR: δ 10.94 (br s, 1H, COOH), 6.17–6.15 (br m, 2H), 2.83–2.75 (br s, 2H), 2.35 (t, J = 7.6 Hz, 2H), 1.68–1.60 (br m, 2H), 1.52–1.40 (br m, 2H), 1.35–1.20 (23H), 0.88 (t, J = 6.8 Hz, 3H); ^{13}C NMR (found 18 C) δ 180.6, 140.0, 139.9, 46.8, 46.7, 34.1, 31.9, 29.9, 29.8, 29.8, 29.7, 29.6, 29.5, 29.3, 29.3, 29.2, 29.0, 28.3, 28.2, 24.6, 22.7, 14.1; HRFAB-MS (Gly matrix) calcd. for $\text{C}_{20}\text{H}_{35}\text{O}_2$ $[\text{M}-\text{H}]^-$: 307.2637; found: 307.2636. Purity >99 % by HPLC (retention time, 4.63 min).

cis-9,10-Ethanoctadecanoic acid (6). A solution of cyclobutene **5** (0.031 g, 0.100 mmol) and 10% Pd/C (0.020 g, 0.020 mmol) in EtOAc (1 mL) was placed in a vial under an atmosphere of H_2 and stirred at rt overnight (16 h). The mixture was then filtered through a plug of Celite, which was washed with approximately 10 mL of EtOAc. The solution was then evaporated under vacuum to afford acid **6** (0.030 g, 96%) as a colorless oil which was sufficiently pure to use without chromatography: R_f = 0.50 (80:20:1 of Hex:EtOAc:AcOH); IR: 3051 (br), 2921, 2852, 1708 cm^{-1} ; ^1H NMR: δ 10.93 (br s, 1H, COOH), 2.35 (t, J = 7.3, 2H), 2.24 (br s, 2H), 2.00–1.90 (br m, 1H), 1.64 (quintet J = 7.3 Hz, 2H), 1.60–1.50 (m, 2H), 1.45–1.10 (br m, 24H), 0.88 (br t, J = 7.0 Hz, 3H); ^{13}C NMR: δ 180.2, 37.7, 37.6, 34.1, 31.9, 30.11, 30.05, 29.9, 29.7, 29.7, 29.4, 29.3, 29.1, 27.64, 27.55, 24.8, 24.7, 22.7, 14.1; HRCI-MS calcd. for $\text{C}_{20}\text{H}_{38}\text{O}_2$ $[\text{M}+\text{H}]^+$: 311.2950; found: 311.2949. Purity 100% by HPLC (retention time, 4.56 min).

2-(Octanoic acid benzyl ester, 8-yl) 3-octylcyclobutanone and 3-(Octanoic acid benzyl ester, 8-yl) 2-octylcyclobutanone (21) was prepared by a modification of a known procedure.^[22–24] To a solution of the regioisomeric dichloroketones **19** (0.670 g, 1.40 mmol) in glacial acetic acid (3 mL) was added Zn(Cu) (0.450 g, 6.90 mmol). The mixture was stirred at rt under N_2 until TLC showed complete disappearance of starting material (~5 h). The reaction mixture was then filtered through a plug of Celite and washed with ether. The filtrate was washed sequentially with saturated aq. NaHCO_3 and water, and dried over Na_2SO_4 . The residue obtained upon concentration was purified by flash chromatography on silica gel (5% EtOAc in Hex) to afford (0.388 g, 87%) of a mixture of cyclobutanones **21** as a colorless oil: R_f = 0.41 (10% EtOAc/Hex); IR: 3034, 2923, 2853, 1774, 1735, 1161, 696 cm^{-1} ; ^1H NMR: δ 7.36–7.26 (br m, 5H), 5.115 (s, 2H), 3.28–3.17 (br m, 1H), 3.15–3.05 (br m, 1H), 2.51–2.43 (m, 1H), 2.42–2.32 (br m, 1H), 2.35/2.34 (overlapping t, J = 7.53 Hz, 2H), 1.70–1.50 (br m, 4H), 1.45–1.14 (br m, 22H), 0.88/0.87 (overlapping t, J = 7.19 Hz, 3H); ^{13}C NMR: δ 212.0, 173.6, 136.1, 128.5, 128.1, 66.0 (br s), 61.9/61.8 (two app s), 50.1/50.1 (two app s), 34.2, 31.8, 30.1, 30.0, 29.6, 29.6, 29.5, 29.4, 29.4, 29.23, 29.17, 29.0, 28.2, 28.08, 28.05, 27.98, 27.6, 24.9, 24.3, 24.3, 22.6, 14.1; HREI-MS calcd. for $\text{C}_{27}\text{H}_{41}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$: 437.3032; found: 437.3029.

2-(Octanoic acid, 8-yl) 3-octylcyclobutanone and 3-(octanoic acid, 8-yl) 2-octylcyclobutanone (7). A solution of cyclobutanones **21** (0.039 g, 0.09 mmol) and 10% Pd/C (0.020 g, 0.18 mmol) in anhydrous MeOH (1 mL) in a vial was placed under an atmosphere of H_2 and stirred at rt overnight (16 h). The mixture was then concentrated under vacuum. The residue was resuspended in EtOAc (~ 20 mL) and filtered through a plug

of Celite. The residue obtained upon concentration was purified by flash column chromatography on silica gel (10%/25% EtOAc/Hex) to afford the regioisomeric mixture of carboxylic acids **7** (0.027 g, 87%) as a colorless oil: R_f = 0.16 (25% EtOAc/Hex); IR: 3322 (br), 2926, 2856, 1776, 1709, 913, 749 cm^{-1} ; ^1H NMR: δ 3.28-3.18 (br m, 1H), 3.13/3.09 (overlapping dd, J = 9.6, 3.1 and 9.0, 3.5, 0.56H/0.44H), 2.50/2.46 (two app abnormal-shape quintets, 0.55H/0.52H), 2.42 (br dd, J = 9.6, 4.5 Hz, 1H), 2.35/2.34 (two app t, J = 7.45 Hz, 2H), 1.63-1.54 (br m, 4H), 1.43-1.18 (br m, 22H), 0.88/0.87 (overlapping br t, J = 7.0 Hz, 3H); ^{13}C NMR: δ 212.1, 179.9, 62.0/61.9 (two app s), 50.20/50.17 (two app s), 34.0, 31.9, 30.1, 30.1, 29.7, 29.6, 29.6, 29.4, 29.3, 29.2, 29.02, 28.98, 28.2, 28.1, 28.0, 27.6, 24.6, 24.3, 24.3, 22.7, 14.1; HREI-MS calcd. for $\text{C}_{20}\text{H}_{36}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$: 347.2557; found: 347.2557. Purity >97% by HPLC (retention time, 6.82 min).

2-(Octanoic acid, 8-yl) 3-octylcyclobutanol and 3-(Octanoic acid, 8-yl) 2-octylcyclobutanol (8). To a -10°C (ice/ NaCl bath) solution of the cyclobutanone **21** (0.037 g, 0.070 mmol) in anhydrous MeOH (1.5 mL) was added NaBH_4 (0.005 g, 0.150 mmol) under N_2 . The reaction mixture was stirred at -10°C until no starting material remained on TLC (~ 20 min). The reaction was diluted with CH_2Cl_2 (about 5 mL) and quenched with sat. NaHCO_3 (2 mL). The mixture was stirred for 10 min. The organic layer was removed and the aqueous layer was extracted with CH_2Cl_2 (3×5 mL). The organic layers were combined, dried over with Na_2SO_4 , and evaporated to afford the alcohol as a colorless oil (0.029 g, 99%). Although the crude product appears pure by ^1H NMR, it is actually a mixture of two regioisomers, each a 70:30 mixture of the 1,2,3- *cis/cis* and the 1,2-*trans*-2,3-*cis* isomers (identities established through 2D-NMR). The individual isomers can be detected by TLC in 20% EtOAc/Hex: R_f = 0.50 (major, all *cis*), 0.45 (major, all *cis*), 0.39 (minor, OH is *trans* to backbone), and 0.37 (minor, OH is *trans* to backbone); IR: 3396 (br), 2922, 2852, 1737, 1156, 732, 696 cm^{-1} ; ^1H NMR: δ 7.39-7.28 (br m, 5H), 5.14 (s, 2H), 4.23 (br q, J = 7.3, 0.7H), 3.93 (br q, J = 7.3, 0.3H), 2.42-2.36 (overlapping signal, 1H), 2.37 (t, J = 7.6, 2H), 2.15-1.95 (br m, 1H), 1.90-1.78 (br m, 1H), 1.70-1.60 (br m, 2H), 1.60-1.48 (br m, 2H), 1.45-1.20 (br m, 22H), 0.90 (br t, J = 7.6, 2H); ^{13}C NMR: δ 173.68/173.65 (overlapping), 136.1, 128.5, 128.1, 72.2, 66.09/66.06 (two overlapping s), 66.0, 43.97/43.95 (overlapping), 37.3, 34.3, 31.9, 30.7, 30.6, 30.2, 29.94, 29.87, 29.8, 29.7, 29.62, 29.58, 29.5, 29.33, 29.29, 29.2, 29.14, 29.08, 29.0, 27.7, 27.6, 24.90, 24.88, 23.4, 23.3, 22.7, 14.1; HRESI-MS calcd. for $\text{C}_{27}\text{H}_{44}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$: 439.3188; found: 439.3174.

Using a procedure similar to that described for preparation of **6**, the cyclobutanol benzyl esters (0.025 g, 0.060 mmol) were subjected to hydrogenation in the presence of 10% Pd/C (0.013 g, 0.012 mmol) and EtOAc (0.6 mL) to afford cyclobutanol acid **8** (0.016 g, 81%) as a colorless oil. Although the product is assumed to consist of a similar mixture of regio- and stereoisomers as was present in the benzyl ester precursor, the individual isomers are not separable: R_f = 0.32 (4:6:0.05 of EtOAc:Hex:AcOH); IR: 3309 (br), 2922, 2853, 1709 cm^{-1} ; ^1H NMR: δ 4.22 (q, J = 7.6, 0.7H, assumed -OH is *cis* to backbone from the characterization of starting material), 3.92 (q, J = 7.6, 0.3H, assumed OH is *trans* to backbone), 2.43-2.33 (br m, 1H), 2.34 (t, J = 7.3, 2H), 2.18-1.92 (m), 1.84 (br septet, J = 8.0, 1H), 1.68-1.58 (br quintet, J = 7.0, 2H), 1.58-1.48 (br m, 2H), 1.43-1.18 (br m, 22H), 0.88 (br t, J = 7.0 Hz, 3H); ^{13}C NMR: δ 179.50/179.46 (two app s), 72.28, 66.16/66.12 (two

app s), 48.87/48.79 (two app s), 43.97, 37.26, 35.06, 33.98, 31.89, 30.69/30.62, 30.24, 30.18, 29.90, 29.87, 29.79, 29.74, 29.63, 29.60, 29.51, 29.46, 29.34, 29.31, 29.21, 29.11, 29.06, 29.01, 28.99, 28.97, 28.67, 28.60, 28.05, 27.95, 27.92, 27.73, 27.60, 24.66/24.63, 23.38/23.34, 22.67, 14.10; HRESI-MS calcd. for $C_{20}H_{38}O_3Na$ $[M+Na]^+$: 349.2719; Found 349.2715. Purity >99 % by HPLC (retention time, 11.8 and 15.9 min).

2-Chloro-4-(octanoic acid benzyl ester, 8-yl)-3-octylcyclobutanone; 2-Chloro-3-(octanoic acid benzyl ester, 8-yl) 4-octylcyclobutanone (22) was prepared based on a modification of a known procedure.^[9,22,24] To a solution of dichloroketone **19** (0.992 g, 2.100 mmol) in 4 mL of glacial acetic acid was added Zn dust (0.148 g, 2.30 mmol). [Note: The amount of Zn required depended upon the purity of the dichlorocyclobutanone. and it was generally necessary to add a second equivalent after approximately 12 h of reaction.] The mixture was stirred at rt until complete disappearance of the starting material was observed (16 h, TLC). The reaction mixture was then cooled in an ice bath and diluted with water (20 mL). The resulting solution was extracted with EtOAc (3 × 50 mL) and the combined organic layers were washed sequentially with water (2 × 100 mL) and saturated $NaHCO_3$ solution (2 × 50 mL). The resulting organic layer was dried over Na_2SO_4 and concentrated under reduced pressure to furnish a colorless oil (0.895 g, 97% crude yield) as a 1:1 mixture of regioisomers, each a 9:1 mixture of *cis/trans* stereoisomers (1H NMR); the mixture was used directly in the next step. Analysis of the 1H NMR indicated that the expected product was accompanied by recovered starting material **19** (approximately 14%) and the cyclobutanone **21** (overreduction product, approximately 10%). These byproducts could not be removed easily by flash chromatography. R_f = 0.58/0.53 (major), 0.48/0.43 (minor) (10% EtOAc in Hex); IR: 3036, 2925, 2854, 1789, 1735, 1456, 1162, 696 cm^{-1} ; 1H NMR: δ 7.36-7.25 (m, 5H), 5.12/5.11 (overlapping s, 2H), 4.97 (dd, J = 2.6, 9.5, 0.1H), 4.51 (dd, J = 2.8, 7.6 Hz, 0.9H), 3.35-3.20 (m, 1.1H), 2.75-2.65 (m, 0.1H), 2.53-2.40 (m, 0.9H), 2.36/2.35 (overlapping t, J = 7.2 Hz, 2H), 1.75-1.20 (m, 26H), 0.92-0.85 (m, 3H); ^{13}C NMR: δ 203.48/203.45, 173.5, 136.1, 128.5, 128.1, 66.0, 65.7, 58.3, 58.2, 40.74/40.72, 34.18, 31.77/31.75, 29.5, 29.3, 29.24, 29.16, 29.14, 29.13, 29.02, 28.97, 28.90, 28.86, 28.02, 27.96, 27.92, 27.87, 25.64/25.57, 24.80/24.78, 22.59/22.58, 14.0; HRESI-MS calcd. for $C_{27}H_{41}O_3ClNa$ $[M+Na]^+$: 471.2642; found 471.2635.

2-Chloro-4-(octanoic acid, 8-yl) 3-octylcyclobutanone-; 2-Chloro-3-(octanoic acid, 8-yl) 4-octylcyclobutanone (9) was prepared in a similar manner as compound **6** from reaction of benzyl ester **22** (0.085 g, 0.19 mmol), 10% Pd/C (0.040 g, 0.038 mmol) and EtOAc (1.9 mL). The product, acid **9** (0.048 g, 70%), was obtained as a colorless oil after purification by flash column chromatography (20% EtOAc/Hex). The product, which was predominantly the *cis,cis*-stereoisomer on the cyclobutanone, included approximately 5% of an inseparable *trans*-chlorocyclobutanone **11** (1H NMR signal at δ 4.37) probably arising from the epimerization of **9**: R_f = 0.19 (4:1 Hex:EtOAc); IR: 3036, 2924, 2854, 1788, 1706, 1461 cm^{-1} ; 1H NMR: δ 10.74 (br s, 1H, COOH), 4.96 (dd, J = 9.4, 2.6, 0.05H), 4.50 (dd, J = 7.6, 2.8, 0.95H), 3.35-3.20 (m, 1.05H), 2.46 (quintet, J = 8.1, 0.95H), 2.74-2.62 (m, 0.05H), 2.35/2.33 (overlapping t, J = 7.3, 2H), 1.80-1.15 (m, 22H), 0.88/0.87 (overlapping t, J = 6.9, 3H); ^{13}C NMR: δ 203.62/203.56, 180.1, 65.7, 58.30/58.26, 401.8, 34.0, 31.8, 29.5, 29.4,

29.3, 29.18/29.15, 29.04/29.00, 28.9, 28.04/27.98, 27.95/27.90, 25.67/25.59, 24.5, 22.6, 14.1; HRESI-MS calcd. for $C_{20}H_{35}O_3ClNa$ $[M+Na]^+$: 381.2172; found 381.2191.

2,2-Dichloro-4-(octanoic acid benzyl ester, 8-yl) 3-octylcyclobutanone-; 2,2-Dichloro-3-(octanoic acid benzyl ester, 8-yl) 4-octylcyclobutanone (23) was prepared based on a known procedure.^[22–24] To a mixture of Zn(Cu) (1.84 g, 28.0 mmol) and benzyl elaidate (2.07 g, 5.60 mmol) in 11 mL of anhydrous ether was added over 2 h (syringe pump) trichloroacetyl chloride (2.0 mL, 18 mmol). The reaction mixture was stirred at rt for 5 h (or until the starting material is consumed by TLC) and filtered through a plug of Celite. The residue was washed with ether (20 mL). The combined ether layers were transferred to a round bottom flask and cooled over ice. The black solution was then diluted with 40 mL of water and 40 mL of sat. $NaHCO_3$. The mixture was allowed to warm to rt and stirred overnight. The aqueous layer was extracted with ether (3×50 mL) and the combined organic layers were dried over Na_2SO_4 and evaporated to dryness. The residue obtained upon concentration could be used for the next step without further purification or, alternatively, could be purified by rapid flash column chromatography through a short plug of silica (2.5%/5% EtOAc/Hex) to afford a yellow oil (1.69 g, 63%) consisting of a 1:1 mixture of regioisomers: R_f = 0.63 (20% EtOAc/Hex); IR: 2926, 2855, 1802, 1735, 1456, 1162, 696 cm^{-1} ; 1H NMR: δ 7.47–7.23 (m, 5H), 5.120/5.117 (overlapping s, 2H), 3.158/3.126 (overlapping dt, J = 7.2, 1.5 Hz, 1H), 2.56–2.46 (m, 1H), 2.37/2.34 (overlapping t, J = 7.6 Hz, 2H), 1.96–1.18 (m, 26H), 0.90/0.87 (overlapping t, J = 7.2 Hz, 3H); ^{13}C NMR, 75 MHz: δ 196.3, 173.5, 136.0, 128.4, 128.1, 87.1, 66.0, 60.7, 60.6, 52.3, 34.2, 34.1, 31.8, 31.7, 31.37, 31.36, 29.42, 29.40, 29.3, 29.23, 29.20, 29.14, 29.12, 29.08, 28.93, 28.90, 28.85, 27.5, 27.4, 27.12, 27.05, 24.80, 24.78, 22.59, 22.57, 14.04, 14.03; HRESI-MS calcd. for $C_{27}H_{40}O_3Cl_2Na$ $[M+Na]^+$: 505.2252; found: 505.2248.

2,2-Dichloro-4-(octanoic acid benzyl ester, 8-yl) 3-octylcyclobutanol, methanesulfonate ester; 2,2-Dichloro-3-(octanoic acid benzyl ester, 8-yl) 4-octylcyclobutanol, methanesulfonate ester (24) was prepared by a similar procedure as **20** through reaction of dichlorocyclobutanone **23** (0.210 g, 0.400 mmol), $NaBH_4$ (0.055 g, 1.50 mmol), and *i*PrOH 6 mL. The 1:1 mixture of regioisomeric dichlorocyclobutanols (0.166 g, 85%) was obtained as a colorless oil following flash chromatography (5% EtOAc/Hex): R_f = 0.66(minor)/0.61(major)/0.56(minor)/0.51(major) (10% EtOAc/Hex). IR: 3459 (br), 2924, 2853, 1736, 1456, 1161, 734, 696 cm^{-1} ; 1H NMR: δ 7.36–7.31 (m, 5H), 5.12 (s, 2H), 4.37 (br d, J = 6.4 Hz, 0.4H), 3.91 (br d, J = 8.0 Hz, 0.6H), 2.80–2.45 (br m, 2H), 2.37/2.36 (overlapping t, J = 7.6 Hz, 2H), 1.80–1.05 (br m, 26H), 0.89/0.88 (overlapping t, J = 6.7 Hz, 3H). Small quantities of individual isomers could be separated for analysis by flash column chromatography using 10% EtOAc/Hex. Stereochemical assignments are based upon 2D-NMR experiments (COSY, NOESY, and HSQC).

First-eluting H_1 - H_3 trans— R_f = 0.66; IR: 3463 (br), 2927, 2855, 1738, 1453, 1158 cm^{-1} ; 1H NMR: δ 7.37–7.31 (m, 5H), 5.12 (s, 2H), 4.38 (ddd, $^3J_{H_1,H_4}$ = 6.8 Hz, $^3J_{H_1,OH}$ = 5.1 Hz, $^4J_{H_1,H_3(trans)}$ = 1.3 Hz, 1H, $CH-OH$), 2.62–2.55 (m, 1H, $CH-CCl_2$), 2.44 (d, $^3J_{H_1,OH}$ = 5.1 Hz, $-OH$), 2.36 (t, J = 7.6 Hz, 2H), 2.18–2.08 (m, 1H, $CH-CHOH$), 1.80–1.10 (br m, 26H), 0.89 (t, J = 6.7 Hz, 3H); ^{13}C NMR (NOESY): δ 173.6, 136.1, 128.5, 128.2, 90.2, 78.5,

66.1, 57.0, 42.0, 34.3, 31.9, 31.3, 29.7, 29.5, 29.4, 29.3, 29.0, 27.7, 27.5, 26.6, 24.9, 22.7, 14.1; HRESI-MS calcd. for $C_{27}H_{40}O_3Cl_2$ $[M+Na]^+$: 507.2409; found: 507.2409.

First-eluting H₁-H₃ cis— R_f =0.61; IR: 3448 (br), 2925, 2854, 1737, 1456, 1163, 734, 697 cm^{-1} ; 1H NMR: δ 7.37-7.31 (m, 5H), 5.12 (s, 2H), 3.91 (dd, $^3J_{H1,OH}$ = 10.8 Hz, $^3J_{H1,H4(trans)}$ = 8.0 Hz, $^4J_{H1,H3(cis)}$ = 0 Hz, 1H, CH-OH), 2.46 (d, J = 10.8 Hz, 2H, OH), 2.36 (t, J = 7.6 Hz, 2H), 2.11-2.04 (m, 1H, CH-CCl₂), 1.75-1.18 (m, 27H), 0.88 (t, J = 6.8 Hz, 3H); ^{13}C NMR, 75 MHz (NOESY): δ 173.6, 136.1, 128.5, 128.2, 91.0, 81.6, 66.1, 51.9, 48.1, 34.3, 33.0, 31.8, 30.0, 29.6, 29.41, 29.38, 29.2, 29.0, 27.3, 26.8, 24.9, 22.6, 14.1; HRESI-MS calcd. for $C_{27}H_{40}O_3Cl_2$ $[M+Na]^+$: 507.2409; found: 507.2415.

Second-eluting H₁-H₃ trans— R_f = 0.56; IR: 3463 (br), 2925, 2854, 1736, 1456, 1163, 696 cm^{-1} ; 1H NMR: δ 7.36-7.31 (m, 5H), 5.11 (s, 2H), 4.40-4.34 (m, 1H), 2.64-2.55 (m, 1H, CH-OH), 2.64-2.55 (m, 1H, CH-CCl₂), 2.46 (d, J = 4.8 Hz, 1H, OH), 2.35 (t, J = 7.6 Hz, 2H), 2.20-2.03 (m, 1H, CH-CHOH), 1.80-1.10 (br m, 26H), 0.88 (t, J = 6.4 Hz, 3H); ^{13}C NMR, 75 MHz: δ 173.7, 136.1, 128.5, 128.2, 90.2, 78.4, 66.1, 57.0, 42.0, 34.3, 31.9, 31.3, 29.6, 29.5, 29.41, 29.39, 29.2, 29.1, 29.0, 27.6, 27.5, 26.7, 24.9, 22.7, 14.1; HRESI-MS calcd. for $C_{27}H_{40}O_3Cl_2$ $[M+Na]^+$: 507.2409; found: 507.2398.

Second-eluting H₁-H₃ cis— R_f = 0.51; IR: 3444 (br), 2924, 2854, 1734, 1162, 734, 696 cm^{-1} ; 1H NMR: δ 7.37-7.31 (m, 5H), 5.11 (s, 2H), 3.90 (dd, $^3J_{H1,OH}$ = 10.8 Hz, $^3J_{H1,H4(trans)}$ = 8.0 Hz, $^4J_{H1,H3(cis)}$ = 0 Hz, 1H, CH-OH), 2.52 (OH, d, J = 10.8 Hz, 2H), 2.35 (t, J = 7.6 Hz, 2H), 2.13-2.02 (m, 1H, CH-CCl₂), 1.75-1.18 (m, 27H), 0.89 (t, J = 6.8 Hz, 3H); ^{13}C NMR: (75 MHz): δ 173.7, 136.1, 128.5, 128.1, 91.0, 81.6, 66.1, 51.9, 48.0, 34.2, 33.0, 31.8, 30.0, 29.6, 29.4, 29.3, 29.2, 29.02, 28.97, 28.9, 27.2, 26.8, 24.9, 24.8, 22.6, 14.1; HRESI-MS calcd. for $C_{27}H_{40}O_3Cl_2$ $[M+Na]^+$: 507.2409; found: 507.2409.

The mixture of dichlorocyclobutanols was converted to the methanesulfonates (mesylate) in a similar manner as for **20**. Reaction of methanesulfonyl chloride (0.05 mL, 0.70 mmol), the dichlorocyclobutanols (0.166 g, 0.300 mmol), and triethylamine (0.20 mL, 1.7 mmol) in CH_2Cl_2 (2 mL), followed by purification using flash chromatography (5% EtOAc/Hex) provided dichloromesylate **24** as a light yellow oil (0.146 g, 86 %) as a mixture of inseparable region- and stereoisomers: R_f = 0.40/0.38 (20% EtOAc/Hex); IR: 2925, 2851, 1734, 1368, 1180, 964, 697 cm^{-1} ; 1H NMR: δ 7.38-7.30 (m, 5H), 5.20 (dd, J = 7.1, 1.2 Hz, 0.4H), 5.115/5.112 (two overlapping s, total 2H), 4.84 (d, J = 8.8 Hz, 0.6H), 3.202/3.199 (two overlapping s, 1.8H), 3.166/3.163 (two overlapping s, total 1.2H), 2.66-2.58 (m, 0.4H), 2.40-2.30 (m, 2H), 2.21-2.12 (m, 0.6H), 2.10-1.98 (m 0.6H), 1.80-1.15 (m, 26.4H), 0.91-0.86 (m, 3H); ^{13}C NMR: 173.6, 136.10/136.08, 128.5/128.1, 86.45/86.38, 84.09, 83.86/83.82, 66.03/66.01, 57.18, 51.81/51.80, 44.51/44.50, 41.71, 39.5, 39.0, 34.2, 32.21/32.18, 31.78/31.77, 31.41/31.36, 30.1, 29.49 29.46, 29.42, 29.34, 29.32, 29.27, 29.22, 29.16, 28.95, 28.90, 28.2, 27.0, 26.9, 26.8, 26.7, 26.6, 26.5, 26.4, 26.3, 24.8, 22.6, 14.1; HRESI-MS $C_{28}H_{44}O_5Cl_2SNa$ $[M+Na]^+$: 585.2184; found: 585.2186.

1-(Octanoic acid, 8-yl)-2-octylcyclobutene (10) was prepared from reductive fragmentation and deprotection of dichloromethanesulfonate **24** (0.143 g, 0.250 mmol) in THF (1 mL), NH_3 (~ 10 mL) and Na (~ 0.090 g, 4.10 mmol) by a procedure similar to that

employed for **5**. The crude product was purified by flash chromatography (10% EtOAc/Hex) to afford the acid **10** (0.035 g, 46%) as a colorless oil: R_f = 0.35 (60:40 of Hex:EtOAc); IR: 3124, 2923, 2852, 1708, 910, 736 cm^{-1} ; ^1H NMR: δ 6.12 (br s, 2H), 2.35 (t, J = 7.2 Hz, 2H), 2.26 (t, J = 7.2 Hz, 2H), 1.64 (quintet, J = 7.2 Hz, 2H), 1.44 (br m, 4H), 1.35-1.20 (23H), 0.88 (t, J = 6.8 Hz, 3H); ^{13}C NMR: δ 179.7, 139.2, 139.1, 50.4, 50.3, 34.02, 33.97, 31.9, 29.9, 29.6, 29.3, 29.2, 29.0, 28.4, 28.3, 24.7, 22.7, 14.1; HRFAB-MS (3-NBA matrix): calcd. for $\text{C}_{20}\text{H}_{35}\text{O}_2\text{Li}_2$ $[\text{M}-\text{H}+2\text{Li}]^+$: 321.2957; found: 321.2969. Purity 100% by HPLC (retention time, 4.73 min).

2-Chloro-4-(octanoic acid, 8-yl) 3-octylcyclobutanone-; 2-Chloro-3-(octanoic acid, 8-yl) 4-octylcyclobutanone (11) was prepared by a similar procedure as described for compound **9**. Reaction of benzyl ester **23** (0.500 g, 1.000 mmol), Zn(Cu) (0.074 g, 1.10 mmol) and glacial acetic acid (2 mL) furnished, after flash chromatography (5% EtOAc/Hex), *trans*-monochlorocyclobutanone (0.313 g, 70%) as a colorless oil which included an equal mix of the 2-chloro-4-octyl and 2-chloro-3-octyl regioisomers, each of which included both epimers at the chloride-bearing carbon: R_f = 0.42/0.39 (minor) and 0.33 (major) (10% EtOAc/Hex). The amount of Zn dust varied from 1.1–2.2 equivalents; generally, a second equivalent of zinc dust was added to the incomplete reaction mixture after stirring for approximately 12 h. The residual dichlorocyclobutanone was easily removed by flash column chromatography. However, the cyclobutanone product of overreduction was formed in significant amounts (28% based upon ^1H NMR and was nearly inseparable by flash column chromatography: IR: 2924, 2854, 1788, 1735, 1457, 1161, 733, 696 cm^{-1} ; ^1H NMR: δ 7.40-7.32 (m, 5H), 5.113/5.110 (overlapping s, 2H), 4.95 (dd, J = 3.1, 9.4, 0.3H), 4.37 (dt, J = 2.7, 7.8 Hz, 0.7H), 2.97-2.90 (m, 0.3H), 2.84-2.74 (m, 0.7H), 2.40-2.30 (overlapping m, 2.3H), 2.08-1.97 (m, 0.7H), 1.85-1.10 (m, 26H), 0.92-0.84 (m, 3H); ^{13}C NMR: δ 204.6, 202.0, 173.5, 136.1, 128.5, 128.1, 66.0, 65.52/65.46, 63.98, 63.78/63.75, 63.57/63.52, 60.74/60.69, 49.8, 43.7, 37.4, 36.63/36.59, 35.12/35.10, 34.2, 31.8, 31.3, 30.3, 30.1, 29.5, 29.42, 29.38, 29.32, 29.25, 29.19, 29.03, 29.00, 28.96, 28.93, 28.88, 28.2, 27.67, 27.60, 27.55, 27.48, 27.38, 27.3, 27.2, 27.1, 24.83/24.80, 22.6, 14.0; HRESI-MS calcd. for $\text{C}_{27}\text{H}_{41}\text{O}_3\text{ClNa}$ $[\text{M}+\text{Na}]^+$: 471.2642; found: 471.2630.

The resulting mixture of monochloroketone mixtures was subjected to debenzylation described in the preparation of compound **6**. Reaction of the benzyl ester mixture (0.034 g, 0.11 mmol), with 10% Pd/C (0.024 g, 0.022 mmol) and EtOAc (1 mL) afforded the unprotected acid (0.013 g, 54%) as a white solid: R_f = 0.24 (6:4:0.05 of Hex:EtOAc:AcOH); mp = 74–75 °C; IR: 3062, 2924, 2854, 1789, 1708, 1464 cm^{-1} ; ^1H NMR: δ 8.70 (br s, 1H), 4.95 (dd, J = 2.9, 9.2, 0.3H), 4.37 (ddd, J = 7.8, 2.5, 1.3, 0.7H), 2.97-2.91 (m, 0.3H), 2.85-2.73 (m, 0.7H), 2.40-2.22 (m, 22H), 2.11-2.98 (m 0.7H), 1.85-1.10 (m 26H), 0.92-0.82 (m, 3H); ^{13}C NMR δ 211.7, 204.7, 202.2, 179.9, 65.6/65.5, 64.0, 63.85/63.82, 63.64/63.58, 60.82/60.77, 49.81, 43.8, 37.5, 36.69/36.65, 35.2, 34.0, 31.8, 31.3, 30.4, 30.2, 29.55, 29.48, 29.44, 29.38, 29.31, 29.25, 29.2, 29.06, 29.06, 28.9, 28.3, 28.2, 27.73, 27.67, 27.62, 27.44, 27.37, 27.26, 27.20, 27.17, 24.63/24.60, 22.7, 14.1; HRESI-MS calcd. for $\text{C}_{20}\text{H}_{35}\text{O}_3\text{ClNa}$ $[\text{M}+\text{Na}]^+$: 381.2172; found: 381.2170.

Determination of inhibitor aqueous stability and solubility

One-dimensional (1D) ^1H -NMR spectroscopy was used to verify the chemical purity, aqueous stability, and concentration dependent micelle formation of eleven fatty acid analogs. Each compound was dissolved in deuterated dimethyl sulfoxide ($\text{DMSO-}d_6$) to obtain a stock concentration of 20 mM. Six different concentrations were prepared from the stock solutions for NMR analysis: 1.00 mM, 750 μM , 500 μM , 100 μM , and 0 μM . All NMR samples consisted of 600 μL of a deuterated 50 mM potassium phosphate buffer at pH 7.2 with 50 μM of 3 (trimethylsilyl) propionic-2,2,3,3- d_4 acid sodium salt (TMSP). Each 600 μL NMR sample contains 30 μL (5%) of $\text{DMSO-}d_6$ and was transferred to a 5 mm NMR tube for analysis.

A Bruker Avance DRX 500-MHz spectrometer equipped with a 5 mm triple-resonance cryoprobe (^1H , ^{13}C , ^{15}N) with a z-axis gradient was utilized for all 1D ^1H NMR experiments. Acquisition of NMR spectra was automated using a BACS-120 sample changer and Icon NMR software. All spectra were acquired at 298.15 K with 16 dummy scans, 64 scans, 32K data points, a spectral width of 5482.46 Hz, and a relaxation delay of 1.5 s. The NMR spectra were processed and analyzed using ACD/1D NMR Manager (Advanced Chemistry Development). The resulting 1D ^1H NMR spectra were visually inspected for evidence of micelle formation (peak broadening), or chemical instability/impurities (additional peaks).

Measurement of nonspecific cytotoxicity

RAW 264.7 macrophages were incubated with 0–100 mM of oleic acid or the C18 cyclobutene fatty acid **2**, each delivered as bovine serum albumin complexes. After 24 h, cell viability was relative to untreated controls was assessed using an IN CTYOTOX-CVDE Crystal violet Dye Elution Kit. The values shown in Figure 2 are means $\pm$ SEM based upon measurement in triplicate.

Bacterial strains and culture conditions

Bacterial strains used in this study are the *E. coli* wild type strain G58-1, the genome sequencing *Msm* strain mc²155, and the *Mtb* strains CDC 1551 and H37Rv. Cells were grown shaking at 37 °C in LB broth for *E. coli* or complete Middlebrook 7H9 broth supplemented with 0.05% v/v Tween 80 to an O.D.₆₀₀ of 0.6–1.2 for mycobacteria. For MIC determinations, cells were inoculated into LB for *E. coli* or a modified previously reported minimal medium for both *E. coli* and mycobacteria.^[26] The minimal media components and final concentrations are as follows: 22 mM dibasic potassium phosphate, 16 mM monobasic potassium phosphate, 2.8×10^{-5} mM ferric chloride, 8.7×10^{-3} mM zinc sulfate, 8.4×10^{-4} mM cobalt(II) chloride, 1.0×10^{-2} mM manganese chloride, 6.8×10^{-2} mM calcium chloride, 2.4 mM magnesium sulfate, 5.0 mM ammonium chloride, 25 mM glycerol, and 0.02% v/v Tyloxapol.

Drug susceptibility assays

MICs were determined by a 96-well microplate method, as described by Chacon.^[26] Bacteria were harvested, washed 2 $\times$ with minimal media, and inoculated to an initial

concentration of approximately 1.0×10^5 colony forming units (CFU) per well. The initial inoculum was plated to verify retrospectively the desired CFU/ml for each strain. Stocks of fatty acid analog compounds were prepared in 100% DMSO- d_6 , as this solvent was chosen to provide consistency with the NMR studies. Appropriate doubling dilution gradients ($\mu\text{g/mL}$ of compound dissolved in the corresponding % v/v DMSO- d_6) were prepared in the following ranges for each compound tested: *E. coli*, 8 to 512; *Msm*, 16 to 1024; and *Mtb*, 0.25 to 256 (e.g., 64 $\mu\text{g/mL}$ of compound corresponded to 0.064% v/v of DMSO- d_6). Alternatively, the concentration of DMSO- d_6 was adjusted in all wells to its maximum concentration in the corresponding gradient (e.g., 64 $\mu\text{g/mL}$ of compound corresponded to 0.256% v/v of DMSO- d_6). We tested cells in the absence of compound and DMSO- d_6 to verify cell viability. The effect of DMSO- d_6 was also tested by growing cells in minimal media with DMSO- d_6 and no compound. We observed no significant effect of DMSO- d_6 on cell viability for *E. coli* and *Msm*. However for *Mtb*, we observed some variability with anomalous inhibitory results in a few technical replicate wells containing DMSO- d_6 alone at higher concentrations. Nonetheless, most of the replicate wells displayed no inhibition allowing us to eliminate these anomalous results. Moreover, the effect of the compound was observed at DMSO- d_6 concentrations having no inhibitory action. Plates were incubated at 37 °C for 2 days (*E. coli*), 4 days (*Msm*) or 7 weeks (*Mtb*). MIC values were determined by the consistent results of three biological and three technical replicates. The MIC was defined by taking the mode of three independent cultures where the MIC did not differ by more than one doubling dilution, discarding any results that are two doubling dilutions away from the mode.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The research was funded from the Life Sciences Interdisciplinary Research Program (University of Nebraska-Lincoln). Portions of this research were performed in facilities renovated with support from the NIH (RR016544). We wish to acknowledge useful discussions with Prof. Ken Nickerson and Prof. Martha Morton (University of Nebraska-Lincoln) and Dr. Chris Schwartz. We also acknowledge technical assistance from Ms. Wendy Austin, Ms. Sara Basaiga, and Mr. Shingo Ishihara (Univ. of Nebraska-Lincoln).

Abbreviations

ATR	Attenuated total reflection
COSY	Correlation spectroscopy
DCC	dicyclohexyl carbodiimide
DCS	D-cycloserine
DMF	<i>N,N'</i> -dimethyl formamide
DMSO-d_6	Hexadeutero dimethyl sulfoxide
DSC/TGA	Differential scanning calorimetry/thermal gravimetric analysis

EtOAc	Ethyl acetate
Hex	Hexane
HPLC	High-performance liquid chromatography
HRESI	High resolution mass spectrometry via electrospray ionization
HRFAB	High resolution mass spectrometry via fast atom bombardment
HSQC	Heteronuclear single quantum coherence spectroscopy
DMAP	4-dimethylaminopyridine
iPrOH	isopropyl alcohol
MDR	Multidrug resistant
MIC	Minimum inhibitory concentration
Msm	<i>Mycobacterium smegmatis</i>
Mtb	<i>Mycobacterium tuberculosis</i>
NOESY	Nuclear Overhauser effect spectroscopy
NMR	Nuclear magnetic resonance
THF	tetrahydrofuran
TLC	thin layer chromatography
UV	ultraviolet

References

1. Raviglione M, Marais B, Floyd K, Lönnroth K, Getahun H, Migliori GB, Harries AD, Nunn P, Lienhardt C, Graham S, Chakaya J, Weyer K, Cole S, Kaufmann SHE, Zumla A. Lancet. 2012; 379:1902–1913. [PubMed: 22608339]
2. a) Koul A, Arnoult E, Lounis N, Guillemont J, Andries K. Nature. 2011; 469:483–490. [PubMed: 21270886] b) Zhang Y. Annu Rev Pharmacol Toxicol. 2005; 45:529–564. [PubMed: 15822188]
3. Ballell L, Field RA, Duncan K, Young RJ. Antimicrob Agents Chemother. 2005; 49:2153–2163. (erratum, 2007, 51, 1888). [PubMed: 15917508] van den Boogard J, Kibiki GS, Kisanga ER, Boerjee MJ, Aarnoutse RE. Antimicrob Agents Chemother. 2009; 53:849–862. [PubMed: 19075046]
4. Takayama K, Wang C, Besra GS. Clin Microbiol Rev. 2005; 18:81–101. [PubMed: 15653820]
5. Kremer L, Dover LG, Morbidoni HR, Vilchèze C, Maughan WN, Baulard A, Tu SC, Honoré N, Deretic V, Sacchettini JC, Loch C, Jacobs WR Jr, Besra GS. J Biol Chem. 2003; 278:20547–20554. [PubMed: 12654922]
6. Several reports have described incorporation of labeled or modified fatty acids into mycolic acid biosynthesis: Morbidoni HR, Vilcheze C, Kremer L, Bittman R, Sacchettini JC, Jacobs WR. Chem Biol. 2006; 13:297–307. [PubMed: 16638535] Wheeler PR, Besra GS, Minnikin DE, Ratledge C. Biochim Biophys Acta. 1993; 1167:182–188. [PubMed: 8466947] Dreher RK, Poralla K, Kong WA. J Bacteriol. 1976; 127:1136–1140. [PubMed: 8428] Kaneda T. Microbiol Rev. 1991; 55:288–302. [PubMed: 1886522]
7. For an overview of application of fatty acids as antimycobacterial activity. see: Carballeira NM. Prog Lipid Res. 2008; 47:50–61. [PubMed: 18023422]
8. Mahindroo VK, Singh V, Dev S. Ind J Chem, Sect B. 1988; 27B:1080–1089.

9. Powers DC, Leber PA, Gallagher SS, Higgs AT, McCullough LA, Baldwin JE. *J Org Chem.* 2007; 72:187–194. [PubMed: 17194098]
10. a) Hyatt JA, Raynolds PW. *Organic Reactions.* 1994; 45:159–646.b) Darses B, Greene AE, Coote SC, Poisson JF. *Org Lett.* 2008; 10:821–824. [PubMed: 18269288] c) Darses B, Greene AE, Poisson J-F. *Org Lett.* 2010; 12:3994–3997. In our hands, the use of Zn and not the more active Zn/Cu was important in obtaining the cis-cyclobutene and not a cis/trans mixture. [PubMed: 20722381]
11. Blaszczyk K, Koenig H, Mel K, Paryzek Z. *Tetrahedron.* 2006; 62:1069–1078.
12. Hassner A, Krepski LR. *J Org Chem.* 1978; 43:2879–2882.
13. Frey HM, Marshall DC, Skinner RF. *Trans Faraday Soc.* 1965; 61:861–865.
14. Davis BD, Dubos RJ. *J Exp Med.* 1947; 86:215–228. [PubMed: 19871672]
15. Even simple fatty acids can display toxicity towards a macrophage line at concentrations beyond 50 – 100 μ M: Martins de Lima T, Cury-Boaventura TMF, Giannocco G, Nunes MT, Curi R. *Clin Sci(Lond).* 2006; 111:307–317. [PubMed: 16737441]
16. a) Rishton GM. *Drug Discov Today.* 1997; 2:382–384.b) McGovern SL, Caselli E, Grigorieff N, Shoichet BK. *J Med Chem.* 2002; 45:1712–1722. [PubMed: 11931626] c) McGovern SL, Helfand BT, Feng B, Shoichet BK. *J Med Chem.* 2003; 46:4265–4272. [PubMed: 13678405] d) Seidler J, McGovern SL, Doman TN, Shoichet BK. *J Med Chem.* 2003; 46:4477–4486. [PubMed: 14521410]
17. Camargo EE, Kertcher JA, Larson SM, Tepper BS, Wagner HN Jr. *Int J Lepr Other Mycobact Dis.* 1982; 50:200–204. [PubMed: 6811449]
18. Barkan D, Liu Z, Sacchettini JC, Glickman MS. *Chem Biol.* 2009; 16:499–509. [PubMed: 19477414]
19. a) Sinninghe Damsté JS, Strous M, Rijpstra WI, Hopnams EC, Geenevasen JA, van Duin AC, van Niftrik LA, Jetten MS. *Nature.* 2002; 419:708–712. [PubMed: 12384695] b) Sinninghe Damsté JS, Rijpstra WIC, Geenevasen JAJ, Strous M, Jetten MS. *FEBS J.* 2005; 272:4270–4283. [PubMed: 16098207]
20. Moore, BS.; Floss, HG. *Comprehensive Natural Products Chemistry.* Sankawa, U.; Barton, DHR.; Nakanishi, K.; Meth-Cohn, O., editors. Vol. 1. Elsevier; New York: 1999. 61 p.81 p.
21. Zhang Y, Heym B, Allen B, Young D, Cole S. *Nature.* 1992; 358:591–593. [PubMed: 1501713]
22. a) Deprés JP, Coelho F, Greene AE. *J Org Chem.* 1985; 50:1972–1973.b) Bak DA, Brady WT. *J Org Chem.* 1979; 44:107–110.c) Greene AE, Deprés JP. *J Am Chem Soc.* 1979; 101:4003–4005.d) Brady WT, Waters OH. *J Org Chem.* 1967; 32:3703–3705.
23. Zn dust was used instead of Zn(Cu) and 2.5 equiv of Cl_3CCOCl was employed
24. Baldwin JD, Belfield KD. *J Org Chem.* 1987; 52:4772–4776.
25. a) Seidl, PR.; Dias, JF. *The Chemistry of Cyclobutanes.* Rappoport, Z.; Liebman, JF., editors. John Wiley & Sons, Ltd; Chichester: 2005. p. 213-256.b) Pretsch, E.; Buhlmann, P.; Affolter, C. *Structure Determination of Organic Compounds: Tables of Spectral Data.* Springer-Verlag; Berlin: 2000. p. 203
26. Chacon O, Feng Z, Harris NB, Cáceres NE, Adams LG, Barletta RG. *Antimicrob Agents Chemother.* 2002; 46:47–54. [PubMed: 11751110]

Figure 1. Analogs investigated, along with structures of oleic (*cis*-18:1), elaidic (*trans*-18:1) and decenoic (10:1) acids. R = octyl.

Figure 2. 1D ^1H NMR spectra of compounds **5** (Figure 2a) and **1** (Figure 2b) were inspected for evidence of micelle formation (peak broadening) by comparing peak widths relative to TMS (standard). Peak widths for **5** demonstrated potential micelle formation or aggregation over the range of tested concentrations (highlighted region). No micelle formation was observed for compounds **1–4** between 100 – 1000 μM .

Scheme 1.

Preparation of analogs in the C₁₀ series (**1–4**): a) DMAP, DCC, BnOH; b) Zn dust (5 equiv), Cl₃CCOCl (2.5 equiv) rt, Et₂O; c) NaBH₄, 2-propanol; d) MsCl, Et₃N; e) Na/NH₃, –78 to –33 °C; f) H₂, Pd/C; g) Zn (5 equiv), AcOH; h) NaBH₄, MeOH.

Scheme 2.

Synthesis of analogs **5–9**: a) DMAP, DCC, BnOH; b) Zn dust (5 equiv), Cl₃CCOCl (2.5 equiv), ether; c) NaBH₄, 2-propanol; d) MsCl, Et₃N; e) Na/NH₃, –78 to –33 °C; f) H₂, Pd/C EtOAc; g) Zn (5 equiv), AcOH; h) NaBH₄, MeOH, –10 °C; i) Zn (1.1 equiv), AcOH. R = octyl.

Scheme 3.

Synthesis of analogs **10** and **11**: a) DMAP, DCC, BnOH; b) Zn(Cu) (5 equiv), Cl₃CCOCl (2.5 equiv), rt, ether; c) NaBH₄, 2-propanol; d) MsCl, Et₃N; e) Na/NH₃, -78 to -33 °C; f) Zn (1.1 equiv), AcOH; g) H₂, Pd/C EtOAc.

Table 1Critical micellar coefficient (CMC) of analogs under assay conditions^a

CMC	100 μ M	1000 μ M
Analog	5–11	1–4

^a Designated concentration or range indicates onset of aggregation observed by ¹H NMR of buffer solutions; see *Experimental Section*.

Table 2

MIC values against *E. coli*, *Msm* and *Mtb*^[a]

Compound	Scaffold	Functionality	<i>E. Coli</i> (G58-1) Day 2		<i>Msm</i> mc2155	<i>Mtb</i>	
			ug/mL (uM)		Day 4	CDC1551 ug/mL (uM)	H37Rv Week 7 ug/mL (uM)
			LB				
1	C ₁₀	alkene	>512 (>2608)	>512 (2608)	512 (2608)	16 (82)	16 (82)
2	C ₁₀	alkane	-	-	512 (2582)	4 (20)	4 (20)
3	C ₁₀	ketone	-	-	256 (1206)	128 (603)	256 (1206)
4	C ₁₀	alcohol	-	-	128 (597)	64 (299)	128 (597)
5	<i>cis</i> -C ₁₈	alkene	-	-	128 (415)	128 (415)	16 (52)
6	<i>cis</i> -C ₁₈	alkane	>512 (>1649)	>512 (1649)	256 (824)	8 (26)	16 (52)
7	<i>cis</i> -C ₁₈	ketone	-	-	128 (394)	64 (197)	32 (99)
8	<i>cis</i> -C ₁₈	alcohol	-	-	256 (784)	8 (24)	2 (6)
9	<i>cis</i> -C ₁₈	chloroketone	-	-	256 (713)	256 (713)	16 (45)
10	<i>trans</i> -C ₁₈	alkene	-	-	256 (830)	128 (415)	16 (52)
11	<i>trans</i> -C ₁₈	chloroketone	-	-	128 (357)	256 (713)	256 (713)
12	C ₁₀	alkene(acyclic)	-	-	512 (3007)	32 (188)	32 (188)
DCS			25 (2508)	16 (157)	64 (627)	4 (39)	8 (78)
Isoniazid			>512 (>3733)	16 (117)	32 (233)	4 (29)	8 (58)
Kanamycin			64 (110)	< 8 (14)	-	-	-

^[a] MIC's were determined as described in the experimental section.

Supporting Information

© Copyright Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, 2014

Development of Cyclobutene- and Cyclobutane- Functionalized Fatty Acids with Inhibitory Activity against *Mycobacterium tuberculosis*

Wantanee Sittiwong,^[a] Denise K. Zinniel,^[b] Robert J. Fenton,^[b] Darrell D. Marshall,^[a]
Courtney B. Story,^[a] Bohkyung Kim,^[c] Ji-Young Lee,^[c] Robert Powers,^[a] Raúl G. Barletta,^[b] and
Patrick H. Dussault^{*[a]}

cmdc_201402067_sm_miscellaneous_information.pdf

Description of Material	page numbers
Synthesis of known benzyl esters	2-3
Cytotoxicity data	4
Thermal stability data	5-10
Investigation of aggregation in aqueous buffer. Stacked ¹ H NMR plots for decenoic acid (12), oleic acid (13), and analogs 2 , 4 , 5 , and 7-11 .	11-20
References for experimental materials	21
¹ H-NMR and ¹³ C-NMR spectrum of 9-decenoic acid, benzyl ester	22
¹ H-NMR and ¹³ C-NMR spectrum of 15	23
¹ H-NMR and ¹³ C-NMR spectrum of dichlorocyclobutanols (C10 series)	24
¹ H-NMR and ¹³ C-NMR spectrum of 16	25
¹ H-NMR and ¹³ C-NMR spectrum of 1	26
¹ H-NMR and ¹³ C-NMR spectrum of 2	27
¹ H-NMR and ¹³ C-NMR spectrum of 17	28
¹ H-NMR and ¹³ C-NMR spectrum of 3	29
¹ H-NMR and ¹³ C-NMR spectrum of 18	30
¹ H-NMR and ¹³ C-NMR spectrum of 4	31
¹ H-NMR and ¹³ C-NMR spectrum of 19 (mixture and individual isomers)	32-37
¹ H-NMR and ¹³ C-NMR spectrum of 20 (mixture and individual isomers)	38-42
¹ H-NMR and ¹³ C-NMR spectrum of 5	43
¹ H-NMR and ¹³ C-NMR spectrum of 6	44
¹ H-NMR and ¹³ C-NMR spectrum of 21	45
¹ H-NMR and ¹³ C-NMR spectrum of 7	46
¹ H-NMR and ¹³ C-NMR spectrum of <i>cis</i> -18:1 cyclobutanols (benzyl esters)	47
¹ H-NMR and ¹³ C-NMR spectrum of 8	48
¹ H-NMR and ¹³ C-NMR spectrum of 22	49
¹ H-NMR and ¹³ C-NMR spectrum of 9	50
¹ H-NMR and ¹³ C-NMR spectrum of benzyl elaidate	51
¹ H-NMR and ¹³ C-NMR spectrum of 23	52
¹ H-NMR and ¹³ C-NMR spectrum of dichlorocyclobutanols (<i>trans</i> 18:1 series)	53-57
¹ H-NMR and ¹³ C-NMR spectrum of 24	58
¹ H-NMR and ¹³ C-NMR spectrum of 10	59
¹ H-NMR and ¹³ C-NMR spectrum of chloroketones (benzyl ester)	60
¹ H-NMR and ¹³ C-NMR spectrum of 11	61

Benzyl 9-decenoate was prepared by a known procedure.¹ Thionyl chloride (36.9 mL, 51 mmol) was added to a stirred solution of 9-decenoic acid **DA** (8.67 g, 51 mmol) and catalytic dimethylformamide (DMF, 0.12 mL, 1.5 mmol) in toluene (400 mL). The reaction was stirred at rt until the starting material was consumed (TLC, ~ 24 h). The reaction was concentrated and the residue redissolved in toluene (400 mL) and triethylamine (8.5 mL). The solution was stirred for 24 h and washed with water. The aqueous layer was then extracted with ether. The combined organic solution were combined, dried with Na₂SO₄, and concentrated under vacuum. The residue was purified by flash column chromatography on silica gel (5-10% EtOAc in Hex) to give benzyl 9-decenoate (13.1 g, 99%) as a colorless oil: R_f = 0.40 (10% EtOAc/ Hex); IR (ZnSe, cm⁻¹): 3072, 2927, 2855, 1734, 1161, 908, 730 cm⁻¹; ¹H NMR: δ 7.38-7.32 (m, 5H), 5.83 (ddt, J = 17, 10.2, 6.6 Hz, 1H), 5.14 (s, 2H), 5.02 (ddd, J = 7.6 Hz, 2H), 5.00 (dq, , J = 17.0, 1.7 Hz, 1H), 4.96 (d of quintet, J = 10.2, 1.2 Hz, 1H), 2.37 (t, J = 7.3 Hz, 2H), 2.10-2.02 (br q, J = 7.3 Hz, 2H), 1.70-1.63 (br m, 2H), 1.41-1.48 (m, 10H); ¹³C NMR: δ 173.5, 138.9, 136.1, 128.4, 128.1, 128.0, 114.1, 65.9, 34.2, 33.7, 28.99, 28.97, 28.8, 28.7, 24.8, 22.6, 14.0; HRESI-MS: calcd. for C₁₇H₂₄O₂Na: [M+Na]⁺ 283.1674; Found: 283.1671.

(Z)-9-Octadecenoate, benzyl ester (benzyl oleate) was prepared by a variation of a known procedure.² Benzyl alcohol (3.83 g, 35.4 mmol) was added to a stirred solution of oleic acid (5.136 g, 18.1 mmol) and DMAP (43.7 mg, 3.57 mmol) in CH₂Cl₂ (40 mL). The solution was cooled to 0 °C and DCC (4.211 g, 20.4 mmol) was added. The mixture was allowed to warm to rt and stirred for 2.5 h. The resulting suspension was filtered through a cotton plug and the precipitate was washed with CH₂Cl₂. The CH₂Cl₂ solution was washed with water and dried over Na₂SO₄. Evaporation of the organic solvent and flash chromatography of the residue over silica gel, using 5% EtOAc/Hex, gave benzyl oleate (5.73 g, 85 %) as a colorless oil: R_f = 0.83 (10%

EtOAc/Hex); IR (ZnSe, cm^{-1}) 2924, 2853, 1738, 1456, 1162, 696 cm^{-1} ; ^1H NMR: δ 7.38-7.31 (5H), 5.41-5.31 (2H), 5.13 (s, 2H), 2.37 (t, $J = 7.6$ Hz, 2H), 2.05-2.00 (4H), 1.70-1.62 (m, 2H), 1.40-1.29 (20H), 0.90 (t, $J = 6.8$ Hz, 3H); ^{13}C NMR δ 173.6, 136.12, 129.9, 129.7, 128.5 (two overlapping signals), 128.11, 128.09, 66.0, 34.3, 31.9, 29.7, 29.6, 29.50, 29.3, 29.1, 29.1, 27.2, 27.1, 24.9, 22.7, 14.1.

(*E*)-9-octadecenoic acid, benzyl ester (benzyl elaidate)³ was prepared from the reaction of elaidic acid (0.9836, 3.5 mmol), DCC (0.79 g, 3.8 mmol), DMAP (0.087 g, 0.7 mmol), and benzyl alcohol (0.73 mL, 7 mmol) in CH_2Cl_2 (10 mL) by a similar procedure as employed for benzyl oleate. The product was purified by flash column chromatography on silica gel (2.5% / 5% EtOAc/Hex) to afford 1.2280 g (95%) of benzyl elaidate as a colorless oil: $R_f = 0.57$ (10% EtOAc/Hex). IR (ZnSe, cm^{-1}) 2922, 2852, 2737, 1160, 966, 696 cm^{-1} ; ^1H NMR: δ 7.38-7.34 (m, 5H), 5.41-5.38 (m, 2H), 5.13 (s, 2H), 2.37 (t, $J = 7.2$ Hz, 2H), 1.98 (br s, 4H), 1.68-1.63 (m, 2H), 1.38-1.18 (br m, 20H), 0.90 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR δ 173.6, 136.1, 130.4, 130.2, 128.5, 128.1, 66.0, 34.3, 32.57, 32.52, 31.9, 29.6, 29.52, 29.46, 29.3, 29.2, 29.1, 29.0, 28.9, 24.9, 22.7, 14.1.

Data from cytotoxicity studies

Figure S1. Analogs investigated

Table S1. Toxicity of Fatty acid analogs in RAW 2647 macrophages

Compound	Average % viability at indicated concentration (μM)					
	0	25	50	100	250	Control ^a
1 (DA-CB)	100	67.8	58.5	48.1	0.9	0.2
2 (DA-satCB)	100	72.9	49.2	7.5	0.6	0.2
4 (DA-alcCB)	100	86.5	78.7	77.5	58.8	0.2
5 (OA-CB)	100	93.8	91.8	89.3	73.4	0.2
6 (OA-satCB)	100	87.8	84.6	82.5	78.9	0.2
8 (OA-alcCB)	100	5.5	4.4	4.4	3.7	0.2
DA	100	83.5	77.3	73	38.3	0.2
D-cycloserine	100	81.8	77.5	72.2	69.4	0.2
Isoniazid	100	76.5	75.1	71.7	64	0.2

a. Sodium dodecyl sulfate (SDS)

Thermal Stability (Figure S1-S11)

Differential scanning calorimetry (DSC) and thermal gravimetric analysis (TGA) were acquired simultaneously on 4-10 mg samples of the indicated molecules, using a Perkin Elmer STA 6000 analyzer operated at a temperature ramp of 2 °C/minute except for compound **5** at a 5 °C/minute

Figure S1. Thermal stability of **1**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S2. Thermal stability of **2**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S3. Thermal stability of **3**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S4. Thermal stability of **4**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S5. Thermal stability of **5**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S6. Thermal stability of **6**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S7. Thermal stability of **7**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S8. Thermal stability of **8**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S9. Thermal stability of **9**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S10. Thermal stability of **10**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Figure S11. Thermal stability of **11**. TGA-measured mass (red) and DSC-measured heat flow (blue).

Decenoic acid: aggregation assay in phosphate buffer

Analog 2 (C₁₀ cyclobutane); aggregation assay in phosphate buffer

Analog 4 (C₁₀ cyclobutanol); aggregation assay in phosphate buffer

Oleic acid: aggregation assay in phosphate buffer

Analog 5 (C_{18} *cis*-cyclobutene); aggregation assay in phosphate buffer

Analog 7 (C_{18} *cis*-cyclobutanone); aggregation assay in phosphate buffer

Analog 8 (C_{18} cyclobutanol); aggregation assay in phosphate buffer

Analog **9 (C₁₈ *cis*-chlorocyclobutanone); aggregation
assay in phosphate buffer**

Analog **10** (C_{18} *trans*-cyclobutene); aggregation assay in phosphate buffer

Analog **11** (C₁₈ *trans*-chloroketone); aggregation assay in phosphate buffer

References

¹ J. J. Li, C. Limberakis, D. A. Pflum, *Modern Organic Synthesis in the Laboratory: A Collection of Standard Experimental Procedures*, Oxford University Press, New York, **2007**, p 45.

² B. Neises, W. Steglich, *Angew. Chem. Int. Ed.* **1978**, *17*, 522–524. Previous report: E. G. Maleeva, *Zhurnal Obshchei Khimii* **1953**, *23*, 1662–1664; K. Murai, G. J. Akazome, *Jpn Oil Chem. Soc.* **1955**, *4*, 125–127. (no detail for preparation)

³ E. G. Maleeva, *Zhurnal Obshchei Khimii*. **1953**, *23*, 1662–1664 [*Chem. Abstr.* **1954**, *48*, 77506].

Figure S1.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum.

Figure S1.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum.

Figure S2.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **15**.

Figure S2.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **15**.

Figure S3.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum.

Figure S3.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum.

Figure S4.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **16**.

Figure S4.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **16**.

Figure S5.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **1**.

Figure S5.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **1**.

Figure S6.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **2**.

Figure S6.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **2**.

Figure S7.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **17**.

Figure S7.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **17**.

Figure S8.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **3**.

Figure S8.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **3**.

Figure S9.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **18**.

Figure S9.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **18**.

Figure S10.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of 4.

Figure S10.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of 4.

Figure S11.1. ^1H -NMR (400 MHz, chloroform-*d*) spectrum of **19**.

Figure S11.2. ^{13}C -NMR (100 MHz, chloroform-*d*) spectrum of **19**.

Figure S12.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of the crude.

Figure S12.2. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of first eluting *cis* isomer.

Figure S12.3. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of first eluting *cis* isomer.

Figure S12.4. ^1H -NMR (400 MHz, chloroform- d) spectrum of first eluting *trans* isomer.

Figure S12.5. ^{13}C -NMR (100 MHz, chloroform- d) spectrum of first eluting *trans* isomer.

Figure S12.6. ¹H-NMR(400 MHz, chloroform-*d*) spectrum of second eluting *cis* isomer.

Figure S12.7. ¹³C-NMR(100 MHz, chloroform-*d*) spectrum of second eluting *cis* isomer.

Figure S12.8. ¹H-NMR(400MHz, chloroform-*d*) spectrum of second eluting *trans* isomer.

Figure S12.9. ¹³C-NMR(100MHz, chloroform-*d*) spectrum of second eluting *trans* isomer.

Figure S13.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of crude **20**.

Figure S13.2. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of first eluting *trans* isomer of **20**.

Figure S13.3. ¹³C-NMR(100 MHz, chloroform-*d*) spectrum of first eluting *trans* isomer of **20**.

FigureS13.4. ¹H-NMR(400MHz,chloroform-*d*)spectrum of second eluting *trans* isomer of **20**.

FigureS13.5. ¹³C-NMR(100MHz,chloroform-*d*)spectrum of second eluting *trans* isomer of **20**.

Figure S13.6. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of first eluting *cis* isomer of 20.

Figure S13.7. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of first eluting *cis* isomer of 20.

Figure S13.8. ¹H-NMR(400 MHz, chloroform-*d*) spectrum of second eluting *cis* isomer of **20**.

Figure S13.9. ¹³C-NMR(100 MHz, chloroform-*d*) spectrum of second eluting *cis* isomer of **20**.

Figure S20.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of compound 5.

Figure S20.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of compound 5.

Figure S21.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of compound 6.

Figure S21.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of compound 6.

Figure S22.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of compound **21**.

Figure S22.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of compound **21**.

Figure S23.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of compound 7.

Figure S23.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of compound 7.

Figure S24.1. ^1H -NMR (400 MHz, chloroform- d) spectrum.

Figure S24.2. ^{13}C -NMR (100 MHz, chloroform- d) spectrum.

Figure S25.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of compound 8.

Figure S25.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of compound 8.

Figure S26.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of compound **22**.

Figure S26.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of compound **22**.

Figure S27.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of compound 9.

Figure S27.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of compound 9.

Figure S28.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum.

Figure S28.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum.

Figure S29.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **23**.

Figure S29.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **23**.

Figure S30.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum.

Figure S30.2. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of first eluting *trans* isomer.

Figure S30.3. ¹³C-NMR(100 MHz, chloroform-*d*) spectrum of first eluting *trans* isomer.

Figure S30.4. ^1H -NMR (400 MHz, chloroform- d) spectrum of first eluting *cis* isomer.

Figure S30.5. ^{13}C -NMR (100 MHz, chloroform- d) spectrum of first eluting *cis* isomer.

Figure S30.6. ^1H -NMR (400 MHz, chloroform-*d*) spectrum of second eluting *trans* isomer.

Figure S30.7. ^{13}C -NMR (100MHz, chloroform-*d*) spectrum of second eluting *trans* isomer.

Figure S30.8. ^1H -NMR (400 MHz, chloroform-*d*) spectrum of second eluting *cis* isomer.

Figure S30.9. ^{13}C -NMR (100 MHz, chloroform-*d*) spectrum of second eluting *cis* isomer.

Figure S31.1. ^1H -NMR (400 MHz, chloroform-*d*) spectrum of **24**.

Figure S31.2. ^{13}C -NMR (100 MHz, chloroform-*d*) spectrum of **24**.

Figure S32.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **10**.

Figure S32.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **10**.

Figure S33.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum.

Figure S33.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum.

Figure S34.1. ¹H-NMR (400 MHz, chloroform-*d*) spectrum of **11**.

Figure S34.2. ¹³C-NMR (100 MHz, chloroform-*d*) spectrum of **11**.