University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Mammalogy Papers: University of Nebraska
State Museum

Museum, University of Nebraska State

11-1-1971

Second Specimen of *Oryzomys dimidiatus*

Hugh H. Genoways

Texas Tech University, h.h.genoways@gmail.com

J. Knox Jones Jr.

Texas Tech University

Follow this and additional works at: <https://digitalcommons.unl.edu/museummammalogy>

Part of the [Zoology Commons](#)

Genoways, Hugh H. and Jones, J. Knox Jr., "Second Specimen of *Oryzomys dimidiatus*" (1971).
Mammalogy Papers: University of Nebraska State Museum. 46.
<https://digitalcommons.unl.edu/museummammalogy/46>

This Article is brought to you for free and open access by the Museum, University of Nebraska State at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Mammalogy Papers: University of Nebraska State Museum by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

SECOND SPECIMEN OF *ORYZOMYS DIMIDIATUS*

The species *Oryzomys dimidiatus* was originally named and described by Thomas (Ann. Mag. Nat. Hist., ser. 7, 15:584–591, 1905) as a member of the genus *Nectomys* on the basis of a single specimen from the Río Escondido, 7 miles below Rama, Nicaragua (approximately 1 km. S and 8½ km. E Rama, 20 m, Zelaya), obtained by W. G. Palmer on 5 November 1904. Although there has been speculation on the relationships of this taxon (currently regarded as the only member of the subgenus *Micronectomys*, genus *Oryzomys*—see HersHKovitz, J. Mamm., 51:789–794, 1970, for review), the holotype has remained the only known specimen.

A second individual (University of Kansas Museum of Natural History no. 106607) assignable to this species was taken on 26 July 1966 by R. W. Turner at El Recreo, 25 m, Zelaya, Nicaragua, approximately 15 kilometers to the west of the type locality. This rat, a young adult male, was trapped in a stand of cane, 8 to 10 feet tall, along the south bank of the Río Mico. The cane was nearly impenetrable excepting for two paths through it to the river, and it was along these paths that traps were set, baited with rolled oats. In addition to *O. dimidiatus*, other small mammals taken at this place included *Oryzomys caliginosus*, *O. fulvescens*, *O. palustris*, *Sigmodon hispidus*, and one *Sylvilagus brasiliensis* that was shot along the edge of the cane.

Measurements (in millimeters) of our specimen followed by (in parentheses, when available) those of the holotype given by HersHKovitz (*op. cit.*) are: head and body, 118 (125); length of tail, 110 (115); length of hind foot, 28 (27, dry); length of ear, 15 (13); greatest length of skull, 29.0 (29.8); condylobasal length, 26.7 (27.2); zygomatic breadth, 15.0 (16.8); interorbital breadth, 4.5 (5.2); breadth of braincase, 11.9 (12.3); mastoid breadth, 11.5; interparietal, 2.5 × 6.9 (2.9 × 7.4); length of nasals, 10.7 (11.7); breadth of zygomatic plate, 2.7 (3.1); rostral width, 5.6 (6.1); rostral length, 10.1; length of incisive foramina, 5.9 (6.5); length of diastema, 7.4 (7.5); length of palatal bridge, 5.5; length of maxillary tooththrow, 4.4 (4.7); crown length of M1, 2.0 (2.2); breadth across upper molars, 5.4 (6.2); length of mandibular tooththrow, 4.5; weight (grams), 46.0. The testes were 11 millimeters long.

Of the species of *Oryzomys* occurring in eastern Nicaragua, *O. dimidiatus* closely resembles *O. palustris*, but differs from that species most conspicuously in smaller external and cranial size, less robust dentition, darker (more black evident) dorsal coloration, grayish as opposed to buffy venter, and proportionally shorter tail (clearly shorter than the length of head and

body in *dimidiatus*, whereas in *palustris* the tail and head and body are approximately the same length).

Field work in Nicaragua was supported by a contract (DA-49-193-MD-2215) from the U. S. Army Medical Research and Development Command. We are grateful to Philip Hershkovitz of the Field Museum of Natural History for confirming the identification of the rice rat here reported.—HUGH H. GENOWAYS AND J. KNOX JONES, JR., *The Museum, Texas Tech University, Lubbock, 79409. Accepted 20 June 1971.*

TWO RHINO MICE (*MUS MUSCULUS*) FROM MINNESOTA

A family of rhinoceros mice, so termed because of their hairless and wrinkled or corrugated skin, was described in 1856 by Gaskoin (Proc. Zool. Soc. London, 24:38–40, 1856). He cited a preserved specimen received by the Museum of the College of Surgeons, London, in 1820, which closely resembled his animals. Allen (Proc. Amer. Acad. Sci., 40:59–163, 1904) described a mouse similar to those reported by Gaskoin, which was caught in Boston about 1904. Howard (J. Hered., 31:467–470, 1940) first used the term “rhino” in designating a specific mutation different from the hairless mutant. She was working with brother-to-sister matings in a laboratory stock derived from a cross between two highly inbred lines. She determined that the mutant gene, rhino, is an allele of hairless and is recessive both to hairless and the normal allele. A rhino mouse may justifiably be considered rare in a natural population of house mice.

On 5 November 1969, Mrs. Agnes Court of St. Cloud, Minnesota, captured a female rhino mouse in her home. This rare mutant of the house mouse had been observed in the kitchen during the previous two months. In late evening, it often ate scraps of food left by the family dog.

FIG. 1.—Rhino mouse (*Mus musculus*) captured in St. Cloud, Minnesota.

The mouse had the typical rhino characteristics (Fig. 1): no hair, except whiskers; excessive growth and folding of the skin to the point where the eyes could no longer be opened and the nails of the right hind foot kept the flank skin constantly irritated; lack of pigment, except that the eyes and the tips of the ears and tail were dark gray. The mouse weighed 36.8 grams. The weights of house mice in Minnesota range from 10 to 25 grams. The rhino mouse was kept in the animal room of the biology department until it died on 2 February 1970. An attempt to breed the mouse to an albino laboratory mouse failed.