

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

The Prairie Naturalist

Great Plains Natural Science Society

---

12-2012

## Terrestrial and Semi-Aquatic Vertebrates in Diets of Largemouth Bass in Central Nebraska

Seth A. Lundgren

*University of Nebraska at Kearney*, [lundgrena@lopers.unk.edu](mailto:lundgrena@lopers.unk.edu)

Keith Geluso

*University of Nebraska at Kearney*

Casey W. Schoenebeck

*University of Nebraska at Kearney*

Follow this and additional works at: <https://digitalcommons.unl.edu/tpn>



Part of the [Biodiversity Commons](#), [Botany Commons](#), [Ecology and Evolutionary Biology Commons](#), [Natural Resources and Conservation Commons](#), [Systems Biology Commons](#), and the [Weed Science Commons](#)

---

Lundgren, Seth A.; Geluso, Keith; and Schoenebeck, Casey W., "Terrestrial and Semi-Aquatic Vertebrates in Diets of Largemouth Bass in Central Nebraska" (2012). *The Prairie Naturalist*. 45.

<https://digitalcommons.unl.edu/tpn/45>

This Article is brought to you for free and open access by the Great Plains Natural Science Society at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in The Prairie Naturalist by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## NOTES

**TERRESTRIAL AND SEMI-AQUATIC VERTEBRATES IN DIETS OF LARGEMOUTH BASS IN CENTRAL NEBRASKA**—Largemouth bass (*Micropterus salmoides*) are opportunistic predators often feeding in littoral areas during crepuscular periods (Snow 1971). Adult largemouth bass primarily are piscivorous, whereas juveniles generally consume zooplankton, insects, and small fish (Zweiacker and Summerfelt 1974, Carlander 1977, Post 2003). Age-0 largemouth bass generally consume zooplankton and macroinvertebrates (Keast and Webb 1966, Keast 1985, Phillips et al. 1995, Post 2003) and shift to piscivory at 50-70 mm in length (Phillips et al. 1995, Olson 1996, Post 2003). Few studies have documented detailed accounts of other vertebrates besides fish in diets of largemouth bass, which include amphibians (Hodgson and Kitchell 1987, Hodgson and Hansen 2005), reptiles (Hodgson and Kitchell 1987, Britson and Gutzke 1993), birds (Hodgson and Hansen 2005), and mammals (Clady 1974, Hodgson 1986, Hodgson and Kitchell 1987, Hodgson and Kinsella 1995). Depending on location and season, non-fish vertebrates can contribute substantially to diets of largemouth bass. For example, small mammals composed over 50% of the volume of bass diets in May-June in Cub Lake, Michigan (Clady 1974). To our knowledge, no study in the Great Plains has documented terrestrial or semi-aquatic vertebrates in largemouth bass diets to date. Herein we report on the presence of terrestrial and semi-aquatic vertebrates in largemouth bass diets in borrow-pit lakes in central Nebraska, representing several new species of prey items for largemouth bass.

We sampled eight borrow pits in Lincoln and Buffalo counties, Nebraska, for largemouth bass in July–October

2010 and May–June 2011, to survey all months with open water. Borrow pits were created in the 1960s when sand was excavated for use in construction of Interstate-80 (McCarraher et al. 1974). Many of these groundwater-filled lakes have been stocked with fish species for recreational purposes by the Nebraska Game and Parks Commission (McCarraher et al. 1974). Borrow pits consist of sandy bottoms, clear water, varying depths, and differing percentages of aquatic vegetative coverage. Fish communities in each lake generally consisted of largemouth bass, bluegill (*Lepomis macrochirus*), and channel catfish (*Ictalurus punctatus*). Additional fish species encountered with less frequency and varying densities included black crappie (*Pomoxis nigromaculatus*), rock bass (*Ambloplites rupestris*), yellow perch (*Perca flavescens*), redear sunfish (*L. microlophus*), northern pike (*Esox luciosus*), flathead catfish (*Pylodictis olivaris*), walleye (*Sander vitreus*), green sunfish (*L. cyanellus*), common carp (*Cyprinus carpio*), and gizzard shad (*Dorosoma cepedianum*).

Aquatic vegetation was sampled during September of 2010 to obtain percent cover by family of submergent and emergent macrophytes. We assessed submergent vegetation by random point sampling following the methods of Stukel (2003). We created random points in ArcView 3.2 (ESRI, Inc., Redlands, CA, USA) and determined sample size by lake surface area with a minimum number of 15 points for each lake. We assessed emergent vegetation using the latest aerial images from the United States Geological Service (USGS) and ArcView by dividing the area containing vegetation by the total surface area of the lake. Additionally, we took Secchi disk readings to assess water clarity (Table 1).

Table 1. Surface area (ha), maximum depth (m), Secchi depth (m), and percent coverage (%) of emergent and submergent vegetation in borrow pit lakes sampled in central Nebraska, 2010–2011.

Lake	Surface Area	Max Depth	Secchi Depth	Vegetation Coverage	
				Emergent	Submergent
Ft. Kearny #5	1.1	6.5	1.2	0.0	33.3
Windmill #2	1.2	5.5	1.2	0.0	46.7
Kea West	3.0	4.5	1.3	17.0	35.0
Crystal	3.0	3.5	2.2	2.8	20.0
West Gothenburg	4.7	5.5	2.4	15.6	20.0
Bufflehead	5.1	3.0	>3.0	0.0	92.0
Fremont Slough	9.9	5.0	0.9	2.9	12.0
Pawnee Slough	12.2	2.5	0.9	24.7	96.0

We captured largemouth bass at night via boat electrofishing with pulsed DC to increase catch per unit effort (Paragamian 1989). We sampled once monthly May–August, and intensively ( $n = 4\text{--}9$  sampling events per pit) during September and October while studying the impact of largemouth bass predation on stocked yellow perch survival in borrow pits. In each borrow pit we sampled until either the entire shoreline was sampled or 50 largemouth bass were captured. We collected prey of largemouth bass from their stomachs using pulsed gastric lavage (Seaburg 1957) and recorded total lengths (mm) and weights (g) of all largemouth bass. We preserved all stomach contents in 95% alcohol and identified vertebrate prey items to the lowest identifiable taxa. Representative individuals of terrestrial and semi-aquatic vertebrates were deposited in natural history collections at the University of Nebraska at Kearney. Research was approved by the Institutional Animal Care and Use Committee at the University of Nebraska at Kearney (Approval number 061010).

We observed that terrestrial and semi-aquatic vertebrates composed less than 1% volumetrically and numerically of all prey items in the diets of largemouth bass for both years. In all, we examined 3,657 stomach samples of largemouth bass and observed 20 terrestrial and 22 semi-aquatic vertebrates, including 6 species of mammals, 2 species of reptiles, 2 taxa of birds, and at least 1 species of amphibian (Table 2). Largemouth bass that consumed vertebrates were mostly adults ( $\geq$ age-3), with exception of one age-1 (TL = 173 mm) and one age-2 (TL = 222 mm) individual. Almost all vertebrates in stomachs were singletons; however, one largemouth bass contained two passerine birds and another contained two unidentifiable Anurans.

Our study documents mammalian species formerly not known as prey items for largemouth bass. To our knowledge, only six species of small mammals are known as prey items for largemouth bass in the United States. The meadow vole (*Microtus pennsylvanicus*), northern short-tailed shrew (*Blarina brevicauda*), woodland jumping mouse (*Napeozapus insignis*), and American water shrew (*Sorex palustris*) were documented in largemouth bass stomachs in the Upper Peninsula of Michigan (Clady 1974, Hodgson 1986). Hodgson and Kinsella (1995) observed two additional species of mammals in bass diets in kettle lakes in the Upper Peninsula of Michigan, including the eastern chipmunk (*Tamias striatus*) and star-nosed mole (*Condylura cristata*). The western harvest mouse (*Reithrodontomys megalotis*), least shrew (*Cryptotis parva*), masked shrew (*S. cinereus*), and prairie vole (*M.*

*ochrogaster*) have not been previously reported as prey items for largemouth bass.

Reptiles also are consumed by largemouth bass, including the painted turtle (*Chrysemys picta*; Hodgson and Hanson 2005); however, Britson and Gutzke (1993) suggested that largemouth bass have a learned avoidance for consuming hatchling painted turtles. We documented a single hatchling painted turtle (weight = 8.07 g, carapace length = 35 mm) consumed by a largemouth bass in central Nebraska (Table 2). The plains garter snake (*Thamnophis radix*) also appears to be the first reported occurrence in the diet of largemouth bass; however, Hodgson and Hansen (2005) documented a northern ribbon snake (*T. sauritus*) consumed in kettle lakes in the Upper Peninsula of Michigan.

Studies indicate that largemouth bass also prey upon frogs including the northern leopard frog (*Lithobates pipiens*), green frog (*L. clamitans*), and spring peeper (*Hyla crucifer*; Hodgson and Hansen 2005). Our documentation of adult plains leopard frogs (*L. blairi*) appears to be the first published record in the diet of largemouth bass. Birds also are prey items of largemouth bass (Hodgson and Hansen 2005). Hodgson and Hansen (2005) observed a red-winged blackbird (*Agelaius phoeniceus*) and a warbler species (*Dendroica* spp.) consumed by largemouth bass. Birds appear least frequently of all non-fish, vertebrates prey items consumed by largemouth bass.

This is the first study to document terrestrial and semi-aquatic vertebrates in the diets of largemouth bass in the Great Plains. Terrestrial and semi-aquatic vertebrates usually do not constitute a large portion of largemouth bass diets, and this study supports what others have reported in other regions of the United States. Surface disturbances, such as swimming, likely contribute to vertebrates being predated by largemouth bass. Our study demonstrates that largemouth bass will prey upon many terrestrial and semi-aquatic vertebrates if opportunities are presented and prey items are of appropriate size.

We thank the Nebraska Game and Parks Commission for use of their electrofishing boats and the University of Nebraska at Kearney for laboratory equipment and technical support. We thank the Federal Aid for Sport Fish Restoration grant F-190-R-2 for financial support. Jared Lorenson, Caleb Huber, Keith Koupal, Ron Albert, Josh Kreitman, and Nate Munter assisted in the field and in the laboratory. We thank Mark Kaemingk for suggestions on an earlier version of this manuscript. — Seth A. Lundgren<sup>1</sup>, Keith Geluso, Casey W. Schoenebeck. Department of Biology, University of Nebraska at Kearney, Kearney NE 68849, USA. <sup>1</sup>Corresponding author email address: [lundgrensa@lopers.unk.edu](mailto:lundgrensa@lopers.unk.edu).

Table 2. Species, date, and location of terrestrial and semi-aquatic vertebrates observed in diets of largemouth bass (*Micropterus salmoides*; LMB) in central Nebraska, 2010–2011. We also reported length (mm) and weight (g) of each LMB sampled.

Species	Date	Lake <sup>a</sup>	LMB	
			Length	Weight
<b>Amphibians</b>				
Anura spp.	31 Aug 2010	Windmill #2	285	355
<i>Lithobates</i> spp.	3 Sep 2010	Windmill #2	260	258
<i>Lithobates</i> spp.	3 Sep 2010	Kea West	222	146
Anura spp.	3 Sep 2010	Ft. Kearny #5	264	239
Anura spp.	6 Sep 2010	Kea West	395	953
<i>Lithobates</i> spp.	15 Sep 2010	Ft. Kearny #5	274	260
<i>Lithobates blairi</i>	20 Sep 2010	Ft. Kearny #5	312	380
1 <i>Lithobates</i> spp. + 1 Anura spp.	20 Sep 2010	Ft. Kearny #5	475	1740
<i>Lithobates</i> spp.	20 Sep 2010	Windmill #2	285	340
<i>Lithobates</i> spp.	28 Sep 2010	Ft. Kearny #5	294	277
<i>Lithobates blairi</i>	28 Sep 2010	Ft. Kearny #5	251	182
<i>Lithobates</i> spp.	28 Sep 2010	Kea West	418	1040
<i>Lithobates</i> spp.	5 Oct 2010	Windmill #2	294	388
Anura spp.	5 Oct 2010	Windmill #2	306	424
<i>Lithobates</i> spp.	5 Oct 2010	Ft. Kearny #5	173	60
Anura spp.	5 Oct 2010	Ft. Kearny #5	287	317
Anura spp.	12 Oct 2010	Ft. Kearny #5	299	375
Anura spp.	12 Oct 2010	Windmill #2	364	641
<i>Lithobates</i> spp.	18 Oct 2010	Kea West	366	768
Anura spp.	18 Oct 2010	Windmill #2	286	360
<b>Small Mammals</b>				
<i>Microtus pennsylvanicus</i>	1 Sep 2010	West Gothenburg	284	300
<i>Blarina brevicauda</i>	7 Sep 2010	West Gothenburg	281	197
<i>Microtus pennsylvanicus</i>	9 Sep 2010	Pawnee Slough	335	488
<i>Reithrodontomys megalotis</i>	9 Sep 2010	Pawnee Slough	285	322
<i>Microtus</i> spp.	9 Sep 2010	Pawnee Slough	377	758
<i>Cryptotis parva</i>	23 Sep 2010	West Gothenburg	302	330
<i>Cryptotis parva</i>	27 Sep 2010	West Gothenburg	290	320
<i>Sorex cinereus</i>	4 Oct 2010	West Gothenburg	282	300
<i>Cryptotis parva</i>	4 Oct 2010	West Gothenburg	291	333
<i>Microtus ochrogaster</i>	26 May 2011	Crystal	320	446
<i>Microtus ochrogaster</i>	26 May 2011	Crystal	288	284
<i>Microtus ochrogaster</i>	26 May 2011	Crystal	315	424
<i>Microtus pennsylvanicus</i>	22 Jun 2011	Crystal	291	316
<b>Reptiles</b>				
<i>Thamnophis radix</i>	12 Oct 2010	Windmill #2	279	300
<i>Thamnophis radix</i>	15 Jun 2011	Pawnee Slough	406	902
<i>Thamnophis radix</i>	15 Jun 2011	Pawnee Slough	291	352
<i>Chrysemys picta</i>	22 Jun 2011	Fremont Slough	369	718
<b>Birds</b>				
2 Passerine spp.	12 Jul 2010	Pawnee Slough	234	186
Passerine spp.	7 Sep 2010	Fremont Slough	404	1045
Tyrannidae spp.	15 Sep 2010	Ft. Kearny #5	290	299

<sup>a</sup> Windmill Lake #2 (40.7059° N, 98.8422° W), Kea West (40.6715° N, 99.1069° W), and Ft. Kearny Lake #5 (40.6537° N, 98.9951° W) were in Buffalo County. West Gothenburg (40.9864° N, 100.3029° W), Pawnee Slough (41.0815° N, 100.54° W), Crystal Lake (41.0834° N, 100.618° W), and Fremont Slough (41.0972° N, 100.6670° W) were in Lincoln County.

## LITERATURE CITED

- Britson, C. A., and W. H. N. Gutzke. 1993. Antipredator mechanisms of hatchling freshwater turtles. *Copeia* 1993:435–440.
- Carlander, K. D. 1977. Handbook of freshwater fishery biology, Volume 2. Iowa State University, Ames, USA.
- Clady, M. D. 1974. Food habits of yellow perch, smallmouth bass, and largemouth bass in two unproductive lakes in northern Michigan. *American Midland Naturalist* 91:453–459.
- Hodgson, J. R. 1986. The occurrence of small mammals in the diets of largemouth bass (*Micropterus salmoides*). *The Jack-Pine Warbler* 64:39–40.
- Hodgson, J. R., and E. M. Hansen. 2005. Terrestrial prey items in the diet of largemouth bass, *Micropterus salmoides*, in a small north temperate lake. *Journal of Freshwater Ecology* 20:793–794.
- Hodgson, J. R., and M. J. Kinsella. 1995. Small mammals in the diet of largemouth bass, revisited. *Journal of Freshwater Ecology* 10:433–435.
- Hodgson, J. R., and J. F. Kitchell. 1987. Opportunistic foraging by largemouth bass (*Micropterus salmoides*). *American Midland Naturalist* 118:323–336.
- Keast, A. 1985. The piscivore feeding guild of fishes in small freshwater ecosystems. *Environmental Biology of Fishes* 12:119–129.
- Keast, A., and D. Webb. 1966. Mouth and body form relative to feeding ecology in the fish fauna of a small lake, Lake Opinicon, Ontario. *Journal of the Fisheries Research Board of Canada* 23:1845–1874.
- McCarragher, D. B., R. A. McDonald, and G. L. Adrian. 1974. Hydrobiological characteristics of Interstate-80 highway lakes in Nebraska. *Transactions of the Kansas Academy of Science* 77:93–102.
- Olson, M. H. 1996. Ontogenetic niche shifts in largemouth bass: variability and consequences for first-year growth. *Ecology* 77:179–190.
- Paragamian, V. L. 1989. A comparison of day and night electrofishing: size structure and catch per unit effort for smallmouth bass. *North American Journal of Fisheries Management* 9:500–503.
- Phillips, J. M., J. R. Jackson, and R. L. Noble. 1995. Hatching date influence on age-specific diet and growth of age-0 largemouth bass. *Transactions of the American Fisheries Society* 124:370–379.
- Post, D. M. 2003. Individual variation in the timing of ontogenetic niche shifts in largemouth bass. *Ecology* 84:1298–1310.
- Seaburg, K. G. 1957. A stomach sampler for live fish. *The Progressive Fish Culturist* 19:137–139.
- Snow, H. E. 1971. Harvest and feeding habits of largemouth bass in Murphy Flowage, Wisconsin. Wisconsin Department of Natural Resources Technical Bulletin 50. Madison, USA.
- Stukel, S. M. 2003. Assessing the sustainability of fish communities in glacial lakes: habitat inventories and relationships between lake attributes and fish communities. Thesis, South Dakota State University, Brookings, USA.
- Zweiacker, P. L., and R. C. Summerfelt. 1974. Seasonal variation in food and diet periodicity in feeding of northern largemouth bass, *Micropterus salmoides salmoides* (Lacepede), in an Oklahoma Reservoir. *Proceedings of the Southeastern Association of Fish and Wildlife Agencies* 27:579–591.

Submitted 4 April 2012. Accepted 29 September 2012.

Associate Editor was M. Wuellner.