

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications, etc. -- Nebraska Forest Service

Nebraska Forest Service

2009

Insect Pests of Evergreen Trees

Mark Harrell

Nebraska Forest Service, University of Nebraska

Rachel Allison

Nebraska Forest Service, University of Nebraska, rachel.allison@unl.edu

Laurie Stepanek

Nebraska Forest Service, University of Nebraska, lstepanek2@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/nebforestpubs>



Part of the [Forest Sciences Commons](#)

Harrell, Mark; Allison, Rachel; and Stepanek, Laurie, "Insect Pests of Evergreen Trees" (2009). *Publications, etc. -- Nebraska Forest Service*. 48.

<https://digitalcommons.unl.edu/nebforestpubs/48>

This Article is brought to you for free and open access by the Nebraska Forest Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications, etc. -- Nebraska Forest Service by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Zimmerman pine moth



Identification

- On pines—mostly ponderosa, Austrian and Scotch.
- Branches are broken or dead.
- Top of the tree may be broken or dead.
- Reddish or yellowish sticky masses of pitch (resin) are present on the trunk or branches.

Control

- Spray bark with a drenching spray of permethrin (High Yield 38 Plus, Astro) or bifenthrin (OnyxPro)* in the second week of April and second week of August.
- Spray bark with dimethoate* before the end of April, but less effective than the sprays above.
- Remove heavily infested trees.

Pine tip moth



Identification

- On pines—mostly ponderosa.
- Dead needles are present at the shoot tip.
- Bud and shoot tip are hollow and may have a small caterpillar inside.

Control

- Spray foliage with acephate (Orthene), spinosad (Conserve), carbaryl (Sevin), permethrin (Hi-Yield 38 Plus, Astro), bifenthrin (Talstar), cyfluthrin (Tempo), tebufenozide (Confirm, Mimic), or dimethoate* in the third week of May and the first week of July.

Ips bark beetles



Identification

- On pines and spruces.
- Many holes about 1/16-inch in diameter are present in the bark.
- Bark may be loose and have tunnels and frass (like sawdust) inside next to the wood.

Control

- Spray bark of trunk with carbaryl (Sevin), permethrin (Hi-Yield 38 Plus, Astro), or bifenthrin (Onyx)* in April, after injury by fire or when freshly cut logs are brought near trees.

*** Trade names are examples of available products. No endorsement is implied. Always follow pesticide label instructions.**

Mark Harrell, Rachel Allison, and Laurie Stepanek
Nebraska Forest Service, University of Nebraska
More information: www.nfs.unl.edu

Photo credits

Spruce needle miner #2: Linnea Gillman, USDA Forest Service, www.forestryimages.org

Pine tussock moth, William M. Ciesla, Forest Health Management International, www.forestryimages.org

Spider mites #1: David Keith, University of Nebraska-Lincoln

Aphids: David Shetlar, Ohio State University



The University of Nebraska-Lincoln does not discriminate based on gender, age, disability, race, color, religion, marital status, veteran's status, national or ethnic origin, or sexual orientation.



Insect Pests of Evergreen Trees

Sawflies



Identification

- On pines—mostly Scotch and ponderosa.
- Needles are missing or chewed.
- Insects are present, often in groups.
- Insects typically rear back when disturbed.
- Insects have 9 or more pairs of legs (other insects have fewer than 9 pairs of legs).

Control

- Spray foliage when the insects appear (mid to late April) with spinosad (Conserve), permethrin (Hi-Yield 38 Plus, Astro), carbaryl (Sevin), imidacloprid (Merit), acephate (Orthene), cyfluthrin (Tempo) or deltamethrin (DeltaGard)*.
- For small trees, knock the insects off the tree with a hard blast of water or gently with a stick.

Bagworm



Identification

- On redcedar, juniper, spruce and pine.
- Foliage is brown or missing.
- Bags covered with dead foliage and up to 2 inches in length hang from branches.

Control

- Spray foliage with Bacillus thuringiensis (Dipel, Thuricide), spinosad (Conserve), permethrin (High Yield 38 Plus, Astro), tebufenozide (Confirm, Mimic), carbaryl (Sevin), acephate (Orthene), malathion, deltamethrin (DeltaGard) or cyfluthrin (Tempo)* when small bags appear (late May to June).
- For small trees, remove and destroy the bags.

Spruce needleminer



Identification

- On Colorado blue spruce, Norway spruce and white spruce (including Black Hills spruce).
- Clusters of dead needles are webbed together.
- Dead needles have a hole at the base.

Control

- Spray foliage with carbaryl (Sevin)* when the new damage is first noticed (June).

Tussock moths

Identification

- On pine—often widespread.
- Needles are missing or chewed.
- Caterpillars are present.

Control

- If needed, spray with Bacillus thuringiensis* or other product labeled for caterpillars on pine.



Spruce spider mite



Identification

- On spruce, juniper, pine and other conifers.
- Yellow speckling appears on needles—more common around the base of the needle and on older needles.
- Damage usually appears in early summer.
- Heavily damaged needles turn reddish brown and fall off.
- Tiny mites look like moving dots on white paper. Hold paper under branch and strike branch to dislodge mites.
- Mites are present April to May and September to October, not in the summer months.

Control

- Spray with growing season oil (NOT on blue or white spruces), insecticidal soap (Safer, Concern), spinosad (Conserve), bifentazate (Floramite), hexythiazox (Hexygon), spiromesifen (Forbid), etoxazole (TetraSan) or abamectin (Avid)* in April to May and September to October. Repeat with the same or different product in 7 to 10 days.
- Reduce applications of insecticides for other pests, which can trigger outbreaks of mites.

Aphids

Identification

- Mostly on spruce and pine.
- Insects are present, usually in groups.
- Shiny sticky material (honeydew) or dark material (sooty mold) may be present on foliage or objects below.

Control

- Reduce applications of nitrogen fertilizers.
- Reduce applications of insecticides for other pests, which could trigger an aphid outbreak.
- Spray with growing season oil (All Seasons, Volck), insecticidal soap (Safer, Concern), permethrin (High Yield 38 Plus, Astro), imidacloprid (Merit), malathion, acephate (Orthene), cyfluthrin (Tempo), or deltamethrin (DeltaGard)* as aphids appear (May through September) or apply a soil treatment with imidacloprid (Merit)* prior to infestation if a serious infestation is expected. The soil treatment has a 60-day delay.



Pine needle scale

Identification

- Mostly on spruce and pine.
- White scales are present on needles.
- Needles may turn yellow then brown.

Control

- Spray with growing season oil (All Seasons, Volck), insecticidal soap (Safer, Concern), acephate (Orthene), permethrin (Eight), malathion, bifenthrin (Talstar) or deltamethrin (DeltaGard)* when eggs hatch and red crawler stage appears (late May to early June and again in July). Repeat in 7 to 10 days unless using oil.

