

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Donald Umstadter Publications

Research Papers in Physics and Astronomy

11-28-2003

Criss-crossing Laser Beams Zoom Electrons Along

P. Zhang

N. Saleh

Shouyuan Chen
schen6@unl.edu

Z.M. Sheng

Donald P. Umstadter
University of Nebraska-Lincoln, donald.umstadter@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/physicsumstadter>




Part of the [Physics Commons](#)

Zhang, P.; Saleh, N.; Chen, Shouyuan; Sheng, Z.M.; and Umstadter, Donald P., "Criss-crossing Laser Beams Zoom Electrons Along" (2003). *Donald Umstadter Publications*. 54.


<https://digitalcommons.unl.edu/physicsumstadter/54>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Donald Umstadter Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Criss-crossing Laser Beams Zoom Electrons Along



Crossing two high-intensity laser beams in a plasma (a collection of charged particles) can have some interesting effects. In a recent experiment performed by researchers at the University of Michigan and the Institute of Physics in China, energy from a higher-power laser pulse was transferred to a lower-power laser pulse. The lower-power pulse had been accelerating electrons with its "wakefield" (like a wave accelerating a surfer). The extra energy to this lower-power pulse enhanced the electron acceleration and decreased the divergence of the electron beam. These features are desirable for proposed "laser particle accelerators" that would be powered by relatively inexpensive and convenient laser light, which is much more powerful than conventional radio-frequency waves.



How does this effect come about? When the laser beams cross at right angles, an interference pattern forms. In the criss-cross zone (indicated by the bright light coming from the intersection region of the two beams in the photo), the interference pattern reflects light in much the same way as does a mirror, redirecting it perpendicular to its original travel. The interference also heats electrons.

In addition to enhancing laser accelerators, this optical beamsplitting effect is expected to play a role in experiments that use lasers to produce fusion energy. In those experiments, high-power laser beams cross each other when heating and compressing a capsule of fusion fuel (for example, solid deuterium). The effects of criss-crossing beams will unavoidably divert some of the laser energy from its original direction, and must be taken into account so as to maximize the effectiveness of using lasers to produce fusion energy.

Thanks to Donald Umstadter, University of Michigan, for the images and help with the caption.

Reported by: [P. Zhang, N. Saleh, S. Chen, Z. M. Sheng, and D. Umstadter](#), *Physical Review Letters*, 28 November 2003.

Published online in *Physics News Graphics*, American Institute of Physics, <http://www.aip.org/mgr/png/2003/209.htm>, December 22, 2003. Copyright ©2003 American Institute of Physics. Used by permission