

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

9-25-2003

Functional Characterization and Expression of a Cytosolic Iron-Superoxide Dismutase from Cowpea Root Nodules

Jose F. Moran

University of Nebraska-Lincoln

Euan K. James

Mario C. Rubio

Gautam Sarath

University of Nebraska-Lincoln, Gautam.sarath@ars.usda.gov

Robert V. Klucas

University of Nebraska-Lincoln

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>



Part of the [Agricultural Science Commons](#)

Moran, Jose F.; James, Euan K.; Rubio, Mario C.; Sarath, Gautam; Klucas, Robert V.; and Becana, Manuel, "Functional Characterization and Expression of a Cytosolic Iron-Superoxide Dismutase from Cowpea Root Nodules" (2003). *Publications from USDA-ARS / UNL Faculty*. 53.
<https://digitalcommons.unl.edu/usdaarsfacpub/53>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Jose F. Moran, Euan K. James, Mario C. Rubio, Gautam Sarath, Robert V. Klucas, and Manuel Becana

First published online September 25, 2003; 10.1104/pp.103.023010

Plant Physiology 133:773-782 (2003)
© 2003 [American Society of Plant Biologists](#)

PLANTS INTERACTING WITH OTHER ORGANISMS

Functional Characterization and Expression of a Cytosolic Iron-Superoxide Dismutase from Cowpea Root Nodules^{1,2}

Jose F. Moran³, Euan K. James, Maria C. Rubio, Gautam Sarath⁴, Robert V. Klucas and Manuel Becana*

Departamento de Nutrición Vegetal, Estación Experimental de Aula Dei, Consejo Superior de Investigaciones Científicas, Apdo 202, 50080 Zaragoza, Spain (J.F.M., M.C.R., M.B.); Centre for High Resolution Imaging and Processing, Medical Sciences Institute/Wellcome Trust Biocentre Complex, School of Life Sciences, University of Dundee, Dundee DD1 5EH, United Kingdom (E.K.J.); and Department of Biochemistry, University of Nebraska, The Beadle Center, P.O. Box 880664, Lincoln, Nebraska 68588 (G.S., R.V.K.)

Abstract: An iron-superoxide dismutase (FeSOD) with an unusual subcellular localization, VuFeSOD, has been purified from cowpea (*Vigna unguiculata*) nodules and leaves. The enzyme has two identical subunits of 27 kD that are not covalently bound. Comparison of its N-terminal sequence (NVAGINLL) with the cDNA-derived amino acid sequence showed that VuFeSOD is synthesized as a precursor with seven additional amino acids. The mature protein was overexpressed in *Escherichia coli*, and the recombinant enzyme was used to generate a polyclonal monospecific antibody. Phylogenetic and immunological data demonstrate that there are at least two types of FeSODs in plants. An enzyme homologous to VuFeSOD is present in soybean (*Glycine max*) and common bean (*Phaseolus vulgaris*) nodules but not in alfalfa (*Medicago sativa*) and pea (*Pisum sativum*) nodules. The latter two species also contain FeSODs in the leaves and nodules, but the enzymes are presumably localized to the chloroplasts and plastids. In contrast, immunoblots of the soluble nodule fraction and immunoelectron microscopy of cryo-processed nodule sections demonstrate that VuFeSOD is localized to the cytosol. Immunoblot analysis showed that the content of VuFeSOD protein increases in senescent nodules with active leghemoglobin degradation, suggesting a direct or indirect (free radical-mediated) role of the released Fe in enzyme induction. Therefore, contrary to the widely held view, FeSODs in plants are not restricted to the chloroplasts and may become an important defensive mechanism against the oxidative stress associated with senescence.

Article, publication date, and citation information can be found at
www.plantphysiol.org/cgi/doi/10.1104/pp.103.023010.