

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Daugherty Water for Food Global Institute:
Faculty Publications

Daugherty Water for Food Global Institute

2016

Anthropogenic Nitrogen and Phosphorus Emissions and Related Grey Water Footprints Caused by EU-271s Crop Production and Consumption

Mesfin Mekonnen

University of Nebraska - Lincoln, mmekonnen2@unl.edu

Stephan Lutter

Vienna University of Economics and Business Administration, stephan.lutter@wu.ac.at

Aldo Martinez

National University of Asuncion, amartinez@pol.una.py

Follow this and additional works at: <https://digitalcommons.unl.edu/wffdocs>



Part of the [Environmental Health and Protection Commons](#), [Environmental Monitoring Commons](#), [Hydraulic Engineering Commons](#), [Hydrology Commons](#), [Natural Resource Economics Commons](#), [Natural Resources and Conservation Commons](#), [Natural Resources Management and Policy Commons](#), [Sustainability Commons](#), and the [Water Resource Management Commons](#)

Mekonnen, Mesfin; Lutter, Stephan; and Martinez, Aldo, "Anthropogenic Nitrogen and Phosphorus Emissions and Related Grey Water Footprints Caused by EU-271s Crop Production and Consumption" (2016). *Daugherty Water for Food Global Institute: Faculty Publications*. 54.
<https://digitalcommons.unl.edu/wffdocs/54>

This Article is brought to you for free and open access by the Daugherty Water for Food Global Institute at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Daugherty Water for Food Global Institute: Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Article

Anthropogenic Nitrogen and Phosphorus Emissions and Related Grey Water Footprints Caused by EU-27's Crop Production and Consumption

Mesfin M. Mekonnen ^{1,*}, Stephan Lutter ² and Aldo Martinez ³

Received: 19 October 2015; Accepted: 14 January 2016; Published: 20 January 2016

Academic Editor: Rolf David Vogt

¹ Twente Water Center, University of Twente, P.O. Box 217, 7500 AE Enschede, The Netherlands² Institute for Ecological Economics, Vienna University of Economics and Business, Welthandelsplatz 1/D4, 1020 Vienna, Austria; stephan.lutter@wu.ac.at³ Energy Systems Research Group (GISE), Polytechnic Faculty, National University of Asunción, Campus San Lorenzo C.C. 2111, Paraguay; amartinez@pol.una.py

* Correspondence: m.m.mekonnen@utwente.nl; Tel.: +31-53-489-6879; Fax: +31-53-489-5377

Abstract: Water is a prerequisite for life on our planet. Due to climate change and pollution, water availability for agricultural production, industry and households is increasingly put at risk. With agriculture being the largest water user as well as polluter worldwide, we estimate anthropogenic nitrogen and phosphorus emissions to fresh water related to global crop production at a spatial resolution level of 5 by 5 arc min and calculate the grey water footprints (GWF) related to EU-27's crop production. A multiregional input-output model is used to trace the the GWF embodied in the final consumption of crop products by the EU-27. The total GWF related to crop production in the EU-27 in 2007 was 1×10^{12} m³/year. Spain contributed about 40% to this total. Production of cereals (wheat, rice and other cereals) take the largest share, accounting for 30% of the GWF, followed by fruits (17%), vegetables (14%), and oil crops (13%). The total agricultural GWF of the EU-27 related to crop consumption was 1830 billion m³/year, which is 3700 m³/year per capita on average. Overall, the EU-27 was able to externalize about 41% of the GWF to the rest of the world through imports of crop products.

Keywords: nitrogen and phosphorus emission; diffuse pollution; grey water footprint; consumption; multi-regional input-output analysis

1. Introduction

The growing global population, crop production, sewage emissions and fossil fuel combustion have significantly altered the global nitrogen (N) and phosphorus (P) cycles. These changes of global nutrient cycles have both positive and negative effects [1]. On the one hand, the past few decades have seen an increase in crop yield that can be attributed to the increased use of N and P fertilizer in agriculture [2]. On the other hand, there is an imbalance between the applied nutrients in agriculture in the form of artificial fertilizer or manure and the nutrients taken up by the crops. Such imbalances—with crops not taking up the complete N and P provided—result in a large fraction of the mobilized N and P entering the freshwater system, causing degradation of the water quality and eutrophication of coastal marine ecosystems [2]. As a consequence, human-induced eutrophication of lakes, estuaries and coastal seas have been reported since the early twentieth century [3–6].

International trade of agricultural products has increased significantly over the last few decades. Taking into account the above described effects of agriculture, this increase results in a globalization of agricultural pollution [7–9]. Importing countries are shifting the environmental damage of agricultural

production to regions that produce and export the products [7–12]. Hence, consumers do not have to bear the environmental cost and, as they may be far away from the regions that produce and supply the products, they are less aware of the environmental damages caused [7–11]. In order to increase awareness as well as to correctly assign responsibilities for environmental impacts to the responsible final consumer, it is necessary to quantify and account for the N and P emissions produced along the supply chain of the traded agricultural products and to relate them to final consumption. This need to measure the environmental pressure generated by intensive agricultural activities has led to the development of indicators that include the Ecological Footprint [13], the grey water footprint (GWF) [10] and the nitrogen footprint [14]. The water footprint (WF) is an indicator of both consumptive water use (water evaporated or incorporated into the product) and water required to assimilate pollutants. We can distinguish the WF into three colours: the green, blue and grey WF. The green WF refers to the consumption of rainwater, the blue WF refers to consumption of surface and ground water. The GWF is an indicator of water pollution. It is defined as the volume of freshwater that is required to assimilate the load of pollutants based on natural background concentration and existing ambient water quality standards [15]. The WF of a nation can be distinguished into: “the WF of national production” and “the WF national consumption”. The WF of national production refers to the total freshwater volume consumed or polluted within the territory of the nation as a result of activities within the agricultural sector, the industrial sector, and the domestic water supply sector. The WF of national consumption, on the other hand, is the total volume of freshwater that is consumed or polluted to produce the goods consumed by the nation’s population. We can further divide the WF of national consumption into two components: the internal and external WF of national consumption. The internal WF refers to the use of domestic water to produce goods consumed by the nation’s population. The external WF, on the other hand, is the volume of water used in other countries to produce goods consumed by the population in the nation under consideration.

Hoekstra and Mekonnen [12] have shown that 430 billion m³ per year of grey water have been traded globally over the period 1996–2005. During the same period, the global total external GWF (*i.e.*, the part of the WF of national consumption falling outside the nation considered) was 218 billion m³ per year. In a recent study, O’Bannon *et al.* [7] showed that the global external GWF increased 136% during the 1986–2010 period.

The aim of this study was to estimate the quantity of nutrients (N and P) emitted and building on these estimates to calculate the GWF of the EU-27 related to both production and consumption of crop products. Further, the sustainability of the EU-27’s GWF of consumption is assessed by identifying priority basins and products. The analysis was carried out using the EXIOBASE v2.2 database (TNO, The Hague, The Netherlands) (base year 2007). EXIOBASE is a global, detailed multi-regional environmentally extended Supply and Use/Input Output (MR EE SUT/IOT) database [16]. The use of EXIOBASE allows tracking the environmental pressure caused by production processed through the supply chains to the final consumer.

The paper is structured as follows: In the following section, the method and data applied in the study is described. This is followed by a section presenting the results of our analysis. The final section provides a discussion of our main findings and the limitation of the study.

2. Materials and Methods

Nitrogen and phosphorus emissions to fresh water. Nitrogen loss through leaching (OUT_{lea} , kg/ha) was calculated using the regression model developed by De Willigen [17]:

$$OUT_{lea} = \left(0.0463 + 0.0037 \times \left(\frac{P}{C \times L} \right) \right) \times (N_f + \alpha \times N_{om} - N_u) \quad (1)$$

where P is the annual precipitation (mm/year), C the clay content (%), L the layer thickness or rooting depth (m), N_f the mineral and manure N (kg N/ha), α the decomposition rate (=1.6% per year), N_{om} the amount of N in soil organic matter (kg N/ha), and N_u the N uptake by harvested crop and crop

residues removed from the field (kg of N/ha per year). N_{om} is calculated by multiplying bulk density (BD , kg/m³), total N in soil ($TOTN$, in kg/kg), and layer thickness L . The data on C , BD , and $TOTN$ were obtained from the derived soil properties on a 5×5 arc-min global grid (version 1.2) from ISRICWISE [18]. The precipitation data for the period 2000–2010 were obtained from the Climate Research Unit of the University of East Anglia (version CRU TS 2.0) [19]. The 30-arc-min precipitation data were converted into 5-arc-min resolution data by assigning the 30-arc-min values to each of the 36 5-arc-min grid cells contained in the 30 by 30 arc-min grid cell. The rooting depth (L , m) for individual crops were obtained from Allen *et al.* [20].

The first part of the regression equation ($0.0463 + 0.0037 \times (P/C \times L)$) determines the fraction of the total mobile N that will leach, and the second part ($N_f + \alpha \times N_{om} - N_u$) determines the available mobile N [21]. To keep the fractional losses less than one, following De Willigen [17], an upper limit of 250 mm/(% m) was set for the combined parameter ($P/C \times L$). Since perennials crops can take up N throughout the year, for these crops, the amount of N in soil organic matter was multiplied by 0.5 in order to further limit overestimation of N leaching [22]. We believe this assumption is reasonable given the fact that other authors have set an upper limit on the total leaching arguing the model might overestimate the leaching volume. For example, Liu *et al.* [23] have set an upper limit on the total leached amount not to exceed 20% of the total mineral fertilizer application and 50% of the N from manure arguing that the model may give unreasonably high leaching values.

We estimated the human-induced (*i.e.*, due to artificial fertilizer and manure application) leaching quantity ($Anthrop_{lea}$) by multiplying the total leached volume (OUT_{lea}) by the fraction of N input in the form of fertilizer and manure (N_f) to the total N input including the amount of N in the soil organic content:

$$Anthrop_{lea} = OUT_{lea} \times \frac{N_f}{(N_f + \alpha \times N_{om})} \quad (2)$$

Modeling the emission of P from diffuse sources at a global scale is more complicated than estimating N. Therefore, we have adopted the approach of Bouwman *et al.* [24], where the amount of P emitted to the water system is assumed to be 12.5% of the P input from the fertilizer and manure application. This value was obtained from the increase of total P river export between 1970 and 2000 as modeled by the global NEWS model [25] and corrected for the global average P retention of 20% in river systems [26].

IFA *et al.* [27] were the primary source of artificial fertilizer application rate per crop per country. This dataset was further complemented with FAO [28] and Heffer [29]. These application rates were adjusted to fit FAO [30] country average artificial fertilizer consumption per year for the period 2007. The manure input was calculated at grid cell level by multiplying livestock density (taken from FAO [31]) by the animal-specific excretion rates (taken from Sheldrick *et al.* [32]). The N and P removal with harvested crops was estimated by multiplying the crop yield by the crop-specific N and P content. Similarly, the N and P removal with crop residues was calculated by multiplying the yield of crop residue by the N and P content of the crop residue and a residual removal factor. The crop yield at 5 arc min per crop was taken from Mekonnen and Hoekstra [33]. The crop-specific N and P contents of major crop and crop residues were taken from IPNI [34]. For other crops and crop groupings, values from FAO [22] and Roy *et al.* [35] were used. For nuts and spices, we have adopted the values of fruits and vegetables respectively from FAO [22]. The crop residues removal factors were obtained from different sources [22,36–40].

Calculating the grey water footprint. The GWFs of N and P are first calculated separately, and then the overall GWF of N and P is determined by the nutrient that is most critical, *i.e.*, the nutrient with the larger GWF. Following Hoekstra *et al.* [15], the GWF (WF_{grey} , m³/year) is calculated by dividing the N and P nutrients load (L , kg/year) by the difference between the ambient water quality

standard for N and P nutrients (the maximum acceptable concentration (c_{\max} , mg/L) and their natural concentration in the receiving water body (c_{nat} , in mg/L)):

$$WF_{\text{grey}} = \frac{L}{(C_{\max} - C_{\text{nat}})} \quad (3)$$

The natural concentration in a receiving water body is the concentration of the water systems that are in pristine condition before human disturbances occur in the catchment. We have adopted the maximum acceptable and natural concentration values for total N and total P from Liu *et al.* [41]. The maximum acceptable concentration values are 3.1 mg N/L and 0.95 mg P/L for total N and total P, respectively, and the natural concentration value of 1.5 mg N/L and 0.52 mg P/L for total N and total P, respectively. Although different ecosystems respond differently to nutrient loads, requiring different maximum allowable concentration values, we have taken a single value for the whole world. Besides, natural concentrations vary from basin to basin. However, obtaining or estimating basin-specific values for both maximum allowable and natural concentration is an elaborate task and impossible for a global study like this one.

To calculate the GWF related to the EU-27 consumption of goods and services, we extend the monetary MRIO model EXIOBASE 2.2 by the physical extensions consisting of GWF values related to the production of 146 different crops in the 48 different countries and regions of the EXIOBASE model. By allocating the GWF to the sectors producing the specific crops, the GWF is “linked” to the trade flows of the crops and ends with them in the countries of final consumption.

The GWF in a river basin is environmentally unsustainable, creating an environmental hotspot when the water pollution level exceeds the waste assimilation capacity. Those basins where the water pollution level creates environmental hotspot are identified as “priority river basins.” The priority river basins and products related to EU-27’s WF of consumption were identified on the basis of the size of the GWF of EU-27’s consumption and the water pollution level (WPL) obtained from Liu *et al.* [41]. Water pollution levels (WPL) can be calculated as the ratio of the total GWF in a catchment to the actual runoff from that catchment. A water pollution level of one means that the waste assimilation capacity has been fully consumed, as the complete runoff is “needed” to assimilate the emissions. When the water pollution level exceeds one, ambient water quality standards are violated. If the EU-27’s consumption causes a GWF in a basin with WPL higher than one, the products contributing a large share to this GWF are identified as priority products as per the definition the basin is defined as priority river basin (see above).

3. Results

In the first part of the results (Section 3.1), we will present the nutrient emission values that are used as input to the EXIOBASE database. Section 3.2 presents the GWF related to the production of crops within the EU-27. The GWF from the production perspective is the total GWF within the territory of the 27 EU countries related to the production of crops. Part of the GWF within the EU-27 can be used to produce goods consumed within the region, and the rest used for producing goods for export to the rest of the world. In Section 3.3, we present the GWF as the result of EU-27 consumption of crop products. For this section, EXIOBASE is used to calculate the GWF embodied in the final consumption of crop products by the EU-27. Part of the GWF related to EU-27 consumption is located within the EU-27 (internal WF) and part of it is located outside the boarder of EU-27 (external WF).

3.1. Anthropogenic Nitrogen and Phosphorus Emissions to Fresh Water Related to EU-27’s Crop Production

Nitrogen and phosphorus emissions to fresh water at high spatial resolution are presented in Figure 1. The results show that the global total N and P emissions from agricultural production in 2007 were 31 and 2.9 million tonnes per year, respectively. However, as Figure 1 illustrates, there are marked spatial variations in the anthropogenic N and P emissions to fresh water across the globe. In Southeastern China and Northeastern India, N loads can reach above 100 kg N/ha. Other places

with large N loads include Western Europe, Central US, and Central America. In Europe, mainly in Germany, Netherlands, Belgium, the UK, Denmark, and Poland, N loads of up to 75 kg N/ha were observed. On the other hand, anthropogenic P load is largest in Northeastern Spain in the Ebro River basin where it reaches as much 50 kg P/ha. In Northeastern China and Northern India, the anthropogenic P load of a little over 5 kg/ha was observed.

The anthropogenic N and P emissions in 2007 within the EU-27 were 0.9 and 0.4 million tonnes per year, respectively. The contribution of major countries to the total EU-27 N and P loads are presented in Table 1. Ireland contributed the largest share (23%) of anthropogenic N load, followed by Germany (19%), UK (14%), Netherlands and Poland, each with 10%. About 50% of the anthropogenic P load in the EU-27 was contributed by Spain, followed by France (11%), and Poland (6%).

In terms of per unit of harvested area, the largest anthropogenic emission occurred in Ireland with 690 kg N/ha and 26 kg P/ha, followed by Netherlands (157 kg N/ha and 17 kg P/ha), Belgium (41 kg N/ha and 9 kg P/ha), and UK (33 kg N/ha and 5 kg P/ha). An excessive nutrient input from manure is mainly responsible for the larger emission in these countries.




Figure 1. Map of global anthropogenic loads of (a) nitrogen and (b) phosphorus due to global production of crops shown in kg/ha on a 5 by 5 arc-min grid. Period: 2007.

Table 1. Anthropogenic nitrogen and phosphorus emissions to water for selected European countries and EU-27 total.

Product	Anthropogenic Nitrogen Emissions		Anthropogenic Phosphorus Emissions	
	k Tonne/Year	kg/ha	k Tonne/Year	kg/ha
Belgium	25	41	6	9
Bulgaria	11	4	6	2
France	70	5	45	3
Germany	176	19	22	2
Greece	13	4	6	2
Ireland	214	690	8	26
Italy	19	3	21	3
Netherlands	90	157	10	17
Poland	91	8	26	2
Portugal	6	5	5	4
Spain	58	5	212	17
UK	139	33	22	5
Other countries	21	1	38	2
Total EU-27	933	11	425	5

3.2. The Grey Water Footprint of EU-27 Related to Crop Production

Figure 2 illustrates the GWFs due to N and P emissions related to the crop production in the EU-27 in 2007. The overall GWF is the larger of the two values; the total overall GWF related to agricultural production in the EU-27 was 1065 billion m³ per year (Table 2). About 78% of this GWF is due to the anthropogenic P emissions, while the remaining 22% is due to anthropogenic N load from crop fields mainly cereals, fruits, and vegetables. Cereals (wheat, rice and other cereals) take the largest share, accounting for 30% of the GWF. The other crops with major contribution to the total GWF are fruits (17%), vegetables (14%) and oil crops (13%). The large GWF by these crops can be explained by their large extent of the harvested area. Cereals (wheat, rice and other cereals) account for 65% of the harvested area, followed by oil crops with 18%, and vegetables, fruits and nuts together with 11%. The other reasons for difference in the contribution of different crops to the overall GWF of EU-27 are the difference in the artificial fertilizer and manure application rate, and the crops nutrient removal rate.

The contribution of different countries to the overall GWF of the EU-27 related to crop production is presented in Figure 2. About 40% of the overall GWF was contributed by Spain, mainly due to anthropogenic P loads from cropland. The other major countries contributing a large share to the EU-27's GWF of production are Ireland (11%), Germany (9%), France (9%), and the UK (7%). In Ireland, Germany and the UK, the larger GWF is due to N loads, while in France anthropogenic load was the dominant one. The large GWF in Ireland is mainly due to large manure application rates, while in Germany, France, and the UK it is mainly due to the large agricultural harvested area combined with high nutrient (fertilizer and manure) application rate.

Table 2. Grey water footprint of production of the EU-27 (million m³ per year). Period: 2007.

Product	Nitrogen	Phosphorus	Overall
Other cereals	105,093	190,130	190,130
Fruits	22,631	175,891	175,891
Vegetables	30,962	144,285	144,285
Oil crops	139,674	95,105	139,674
Wheat	106,408	115,376	115,376
Pulses	11,775	85,151	85,151
fodder crops	62,581	63,121	63,121
Roots and tubers	51,324	32,720	51,324
Sugar crops	41,003	27,681	41,003
Nuts	9580	31,330	31,330
Paddy rice	15	12,869	12,869
Other crops	1849	11,775	11,775
Fibres	43	2985	2985
Total	582,938	988,419	1,064,914




Figure 2. Contribution of different countries to the overall GWF of EU-27 related to production. Period: 2007.

3.3. The Grey Water Footprint Related to EU-27 Consumption

The total agricultural GWF of the EU-27 related to consumption in 2007 was 1834 billion m³/year, which is 3700 m³/year per capita on average (Table 3). Looking at some selected European countries, the difference in the per-capita WF is quite large, ranging from 8900 m³/cap/year for Spain to 1600 m³/cap/year for Estonia.

The extent to which countries externalize their WF varies considerably. For the EU-27 as a whole, the share of the external WF was 41%. The share of the external GWF for the selected countries varies from 26% for Spain to 84% for Estonia. The extent to which single EU-27 countries externalize their GWF to other EU-27 countries or to the rest of the world also varies. For Spain, 42% of its GWF was external but only 4% was externalized to the EU-27, while 22% was to the rest of the world. Estonia, on the other hand, shifted 26% of its grey WF to the EU-27 and 57% to the rest of the world. The large share of the external WF shows that the EU-27 and the other countries displaced a large portion of the water pollution to the rest of the world.

Table 3. Grey water footprint of the EU-27's and some selected countries related to consumption of agricultural products. Period: 2007.

Country	Total GWF (Gm ³ /Year)			Per capita GWF (m ³ /Cap/Year)	
	Internal	External GWF			Total
		Within EU-27	Rest of the World		
EU-27	1080		754	1834	3694
Estonia	0.3	0.6	1.2	2.1	1608
Germany	76	74	178	328	3924
Netherlands	17	17	46	80	4839
Poland	46	11	20	77	2023
Spain	294	15	90	399	8931
UK	72	64	129	265	4322

The origin of the global GWF of the EU-27 related to agricultural products consumption is shown in Figure 3. The figure shows that, although the external GWF of the EU-27 is spread across the rest of the world, its extent is concentrated in only a few places. Large external GWFs of the EU-27 can be observed in the Mississippi river basin (USA), Parana River basin (shared by Argentina, Bolivia, Brazil and Paraguay), Indus River basin, Southeastern China in the Huai He River basin, Yellow River basin and Yangtze River basin, and the Nile Delta in Egypt. The large GWF (both internal and external) is the function of the volume of the different crops supplied to the EU-27 and the GWF per unit of product (m³/tonne) of the consumed crops. Figure 3 shows a number of darker areas within the EU-27, which can be explained by the fact that about 59% of the GWF of consumption of the EU-27 is internal (within EU-27). The other darker area is in Eastern China, which is due to both the large volume of oil crops and cereals imported from China, and an excessive fertilizer application rate in China.

The contribution of different crops to the average GWF of consumption of the EU is presented in Figure 4. One quarter of the GWF relates to the consumption of oil crops (including cotton, soybean, oil palm, sunflower, rapeseed and others). The difference in the GWF among the different crops is mainly due to the rate of application of artificial fertilizers and manure, plus the crops' nutrient removal rate. For example, cereal crops have larger N removal rates compared to oil seeds (e.g., 55 kg N/tonne of wheat compared to 10 kg N/tonne of oil crops). In addition, the fraction of N leaching from the agricultural field depends on the precipitation and the clay content of the soil as shown in Equation 1. The leaching fraction will be larger for areas with low clay soil and high precipitation. The large GWF of oil crops comes mainly from China, which has relatively larger fertilizer application rate and anthropogenic N and P emissions. The other products contributing a large share to the EU's GWF of consumption are other cereals (14%), fruits (12%), wheat (11%), and vegetables (10%). Eastern China

and developed Western countries (EU and USA) are known to have an excessive nutrient application rate, while the fertilizer application rate in developing countries such as Africa is not sufficient to supply the crop nutrient requirement [2]. This is reflected in the high N and P loads in Figure 1 and large GWF in Figure 3, in Eastern China and Western Europe.




Figure 3. The global GWF of the EU-27 related to consumption of agricultural products shown in mm/year on a 5 by 5 arc-min grid. Data per grid cell have been calculated as the WF within a grid cell (in cubic meters per year) divided by the area of the grid cell (in 10^3 m^2). Period: 2007.




Figure 4. The contribution of different crops to the overall GWF of the EU consumption of agricultural products. Period: 2007.

3.4. Priority Basins and Products

A detailed analysis of the GWF of the EU-27 related to consumption of agricultural products shows that about 87% of the total GWF occurs in basins where the WPL is beyond 1.0, *i.e.*, the ambient water quality standard are violated (Table 4). The hotspots with respect to the GWF of the EU-27's consumption of agricultural products are identified based on the WPL in the respective river basins where these GWFs are located. Table 4 presents the top 20 river basins ranked by size of the GWF of the EU related to the consumption of agricultural products. The table also shows the crop products with significant contribution to the GWF within the basin. In all 20 river basins, the WPL exceeds 1.0, indicating that in all basins the water quality standards are violated.

Table 4. The water pollution level and contribution of major crops in the priority basins supporting the largest share of the total GWF related to EU's consumption.

Basin	Percentage the Total GWF EU-27's Consumption Located in the Basin	Water Pollution Level for the Nutrient with Larger GWF	Products with Significant Contribution to the GWF in the Basin (% Contribution)
Yangtze (Chang Jiang)	8.5%	2.50	oil crops-39%, paddy rice-14%, roots and tubers-12%, vegetables-10%, other cereals-8%, wheat-7%
Ebro	6.9%	1.08	fruits-28%, vegetables-18%, other cereals-16%, pulses-14%, oil crops-6%, nuts-5%
Douro	4.2%	1.53	fruits-24%, other cereals-20%, vegetables-17%, pulses-12%, nuts-5%, oil crops-5%
Huai He	3.0%	7.08	oil crops-38%, wheat-19%, other cereals-16%, vegetables-7%, roots and tubers-7%, paddy rice-6%
Guadalquivir	2.9%	8.31	oil crops-24%, fruits-21%, vegetables-15%, pulses-11%, other cereals-9%
Danube	2.8%	1.09	other cereals-29%, Wheat-17%, oil crops-12%, fruits-10%, vegetables-9%, fodder crops-8%
Guadiana	2.6%	1.70	fruits-24%, vegetables-16%, other cereals-15%, pulses-12%, oil crops-11%, nuts-6%
Tejo	2.4%	1.00	fruits-26%, vegetables-17%, other cereals-16%, pulses-13%, oil crops-9%, nuts-6%
Mississippi	2.1%	1.33	oil crops-97%
Xi Jiang	2.0%	2.49	oil crops-29%, sugar crops-21%, vegetables-14%, paddy rice-13%, other cereals-9%, roots and tubers-7%
Rhine	1.5%	2.59	oil crops-39%, other cereals-23%, sugar crops-11%, wheat-9%, roots and tubers-6%
Wisla	1.5%	3.33	fruits-21%, oil crops-18%, wheat-15%, roots and tubers-14%, sugar crops-14%, other cereals-10%, vegetables-8%
Elbe	1.4%	2.72	oil crops-58%, other cereals-16%, sugar crops-9%, wheat-6%, vegetables-5%
Oder	1.4%	2.14	oil crops-37%, sugar crops-16%, fruits-13%, wheat-13%, roots and tubers-7%, other cereals-7%, vegetables-5%
Huang He (Yellow)	1.2%	2.87	other cereals-25%, wheat-24%, oil crops-23%, vegetables-6%, fruits-7%, roots and tubers-7%
Amur	1.1%	1.18	oil crops-34%, other cereals-33%, wheat-12%
Loire	1.1%	1.58	other cereals-44%, oil crops-22%, pulses-19%, Fruits-7%
Indus	1.1%	2.99	wheat-50%, oil crops-35%, sugar crops-6%, pulses-6%
Uruguay	1.0%	1.02	oil crops-98%
Yongding He	0.9%	8.98	other cereals-29%, wheat-26%, oil crops-21%, fruits-9%, vegetables-7%

The Yangtze River basin (China) and Ebro River basin (Spain) are two of the most important priority basins, contributing about 8.5% and 6.9% to the overall GWF of the EU-27's consumption of agricultural products, respectively. In the Yangtze River basin, oil crops production accounts for a significant share of the GWF (39%), followed by rice (14%) and root & tubers (12%). In the Ebro River

basin, fruits are the dominant crops that account for the largest share of the GWF in the basin (28%), followed by vegetables (18%) and other cereals (16%).

The Douro (shared by Spain and Portugal) and Huai He (China) are the other priority basins, where 4.2% and 3% of EU's grey WF is located, respectively. The major crops accounting for a large share of the GWF in the Douro basin are fruits production (24%), other cereals (20%), and vegetables (17%). In the Huai He basins, oil crops (mainly cotton production) contribute about 38% to the GWF, followed by wheat (19%) and other cereals (16%). Next in the list of the priority basins are the Guadalquivir and the Danube, where 2.9% and 2.8% of the GWF of EU's consumption are located, respectively. The GWF in the Guadalquivir river basin is mainly due to the production of oil crops, fruits, and vegetables. In the Danube basin, the major crops responsible for the large GWF are mainly other cereals, wheat, and oil crops production.

The results of this basin prioritization provide the EU-27 with a clear focus on two priority regions for crop production and sourcing: within the EU-27 itself and Eastern Asia (mainly China). Among the 20 priority basins, 11 of them are within the EU-27 and 6 mainly in China.

There is diversity in the priority crops among the basins. However, five crops and crop categories stand out as the major priority crops: oil crops, other cereals, fruits, wheat and vegetables. These five products also account for about 72% of the overall GWF of the EU-27's consumption (Figure 4). Oil crops (including cotton) are the most important crops contributing the largest share of the EU-27 consumers GWF. They are listed as the priority crops in all the priority basins.

4. Discussion and Conclusions

In the current study, we estimate the global N and P emission related to crops production. The study also provides important information on the level of nutrient emission to water and the GWF of the EU-27, both related to the production and consumption of crops.

Our estimate of global total N emission of 31 million tonnes of N per year is 20% lower than the estimate by Bouwman *et al.* [24] (39 million tonne N/year) and 35% above the estimate by Liu *et al.* [23] (23 million tonne N/year). Our estimate of global P emission of 2.9 million tonne P per year is 28% lower than the estimate by Bouwman *et al.* [24] (4 million tonne P/year). It should be noted that the estimate by Bouwman *et al.* [24] includes nutrients flows from grassland, which is not included in the current study. Besides, the two studies refer to two different time periods—while the values in the current study refer to the year 2007, the Bouwman *et al.* [24] study refers to the year 2000.

We further compared our estimate of total N and P emission in from the EU-27's croplands to freshwater with results from earlier studies. Our estimate of total N emission from EU-27's croplands is much lower than the other studies (Table 5): It varies from as low as 30% of the estimate by Leip *et al.* [42] to almost half of the estimate by Westhoek *et al.* [43]. Further, differences in the modeling approach, the large difference between our estimate and the earlier studies, can be explained by the following facts: (a) our estimate refers to cropland, while the estimate from the other studies include nutrients flows from grassland; (b) the other studies have considered inputs through N deposition and fixation, while we consider only fertilizer and manure inputs in croplands; and (c) the study by Leip *et al.* [44] and Westhoek *et al.* [43] have considered N flows from other countries to EU through crop imports. On the other hand, our estimate of the total P emission is 87% larger than the estimate by Grizzetti *et al.* [45]. The estimate by Grizzetti *et al.* [45] refers to nutrient flows to European seas after accounting for nutrient retention in the river networks, while our estimate accounts for all the emission that goes to the freshwater systems. Although the volume of the estimated nutrients emission vary greatly among the different studies, the findings of all of the studies, including ours, underline the need to control nutrient emission in EU-27 and around the world to manage the undesired consequences on the aquatic ecosystems.

Table 5. Comparison of the Estimated Overall Diffuse N Load from the EU-27/S Croplands to Fresh Water with the Results from Previous Studies.

Study	N Leaching and Runoff to Fresh Water from Diffuse Sources (Million Tonne N/Year)	P Leaching and Runoff to Fresh Water from Diffuse Sources (Million Tonne P/Year)	Study Period
Leip <i>et al.</i> [42]	5.98	–	2002
Liep <i>et al.</i> [44]	5.0	–	2004
Westhoek <i>et al.</i> [43]	3.3	–	2004
Grizzetti <i>et al.</i> [45]	4.77	0.23	2000
Current study	1.68 *	0.43	2007

* Note: This refers to the total N leaching and runoff as estimated by Equation (1), which includes both natural and anthropogenic emissions.

The nutrient emission estimates are derived based on a large number of inputs that may include uncertainties and possible errors. Possible sources of uncertainties include: the fertilizer and manure application rate, the crops nutrient uptake rate, the maximum allowable nutrient concentration, and the natural nutrient concentration. For the GWF in particular, the maximum allowable concentration and the natural nutrient concentrations values taken will have large impacts on the final result. For example, Liu *et al.* [41] have shown that a slight increase/decrease results in a large change in the final values of the WPL. In a recent study, Mekonnen and Hoekstra [46] showed that the uncertainty of the N related global GWF due to uncertainty in input data and parameters ranges from –33% to +60%. Such large uncertainty ranges highlights the need for further detailed assessment of the nutrient emission and GWF based on local data and better modeling tools. In general, depending on the elements included and the data on the maximum allowable concentration and natural concentration used, the final volume of the GWF differs substantially. To address this and other criticism directed at the GWF concept, Vanham and Bidoglio [47] suggest further standardization of the GWF method.

The other potential uncertainties are related to the models used in estimating both the N and P emission per grid cell. The regression model of De Willigen [17] used here in estimating N emission is based on experimental data, and could therefore give unreasonably high leaching values when used for input parameters (precipitation, clay content, and layer thickness) beyond those used in deriving the regression equation. Liu *et al.* [23], for example, have set an upper limit on the total leached amount not to exceed 20% of the total mineral fertilizer application and 50% of the N from manure, arguing that the model may give unreasonably high leaching values for extreme precipitation. On the other hand, the model accounts for the effect of precipitation neglecting the effect of irrigation, potentially underestimating the N leaching from irrigated fields. The P emission is also estimated using a single factor derived from extensive modeling exercise applied at a global scale. Since it does not account for the local conditions (e.g., precipitation and soil P content) that might influence P runoff from agricultural field, the estimates at grid level are subject to large uncertainties and errors. Given the fact that 78% of the GWF is related to anthropogenic P emission, further refinement of the approach in modeling P emission will improve the final result. Given the uncertainties in the input variables and the modeling, the results in this paper should be taken as indicative of the nutrient loads and the grey WF magnitudes rather than as an absolute value.

The study also does not assess the final effect of eutrophication of lakes, estuaries and coastal seas associated with the emitted nutrients, as this is beyond the scope of the present paper. In future studies, the assessment of such effects on fish, the threat on biodiversity, on the tourism and recreation activities would add value to the analysis results.

The increases in international trade of agricultural products in the last few decades have also increased the globalization of virtual water flows as well as agricultural pollution. Consumers in one part of the world are shifting the environmental damage brought about by fertilizer application to regions that produce and export the products often without being aware of the environmental damages. The methodology of tracing global virtual water flows through an environmentally extended input-output framework such as the EXIOBASE has proven to be a valuable tool for providing insight

into the environmental pressure put by consumers in one part of the world on regions that produce and export the product. By combining the data on direct and indirect water consumption per country with data on water pollution, it is possible to trace water pollution induced by final consumption.

This study is a major contribution to understanding the sustainability of the EU's consumption with respect to water pollution both within the EU and the rest of the world. By tracing the global GWF of the EU, the study made the link between the crops consumed in the EU to the water pollution problem in other places. It further assessed the geographically specific extent to which the EU has externalized its environmental damage by identifying priority basins and products. Prioritizing or identifying the basins with serious water pollution and the associated products responsible for the large GWF can help the EU policy makers and governments to formulate policies and strategies on how to share responsibility for reducing GWFs where it is most needed.

As a consequence, analyses like the one presented in this paper may further encourage EU policy makers to develop GWF reduction targets that account for the assimilation capacity of river basins. These targets can be translated into maximum acceptable loads per river basin to be downscaled to the specific crops production. In many regions of the world, crops receive excessive amounts of nutrient. In these regions, the excessive application of nutrients can be reduced without affecting agricultural productivities [2]. Reducing the application of N and P in agriculture provides both environmental and economic benefits. The EU could implement a number of regulations including the certification of imported crop products in order to promote environmentally sustainable crop production in other countries.

Acknowledgments: The EXIOBASE database used in this research was developed through EU's Sixth and Seventh Framework Programmes. This research was partly supported by the European Community FP7 project CREEA (Compiling and Refining Environmental and Economic Accounts), grant agreement No. 265134.

Author Contributions: Mesfin Mekonnen designed and performed research. Stephan Lutter and Aldo Martinez carried out the input-output modeling. Mesfin Mekonnen wrote the paper, and Stephan Lutter provided comments. All authors approved the final manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Galloway, J.N.; Cowling, E.B. Reactive nitrogen and the world: 200 Years of Change. *AMBIO J. Hum. Environ.* **2002**, *31*, 64–71. [[CrossRef](#)]
2. Vitousek, P.M.; Naylor, R.; Crews, T.; David, M.B.; Drinkwater, L.E.; Holland, E.; Johnes, P.J.; Katzenberger, J.; Martinelli, L.A.; Matson, P.A.; *et al.* Nutrient imbalances in agricultural development. *Science* **2009**, *324*, 1519–1520. [[CrossRef](#)] [[PubMed](#)]
3. Smith, V. Eutrophication of freshwater and coastal marine ecosystems a global problem. *Environ. Sci. Pollut. Res.* **2003**, *10*, 126–139. [[CrossRef](#)]
4. Landsberg, J.H. The effects of harmful algal blooms on aquatic organisms. *Rev. Fish. Sci.* **2002**, *10*, 113–390. [[CrossRef](#)]
5. Lewitus, A.J.; Horner, R.A.; Caron, D.A.; Garcia-Mendoza, E.; Hickey, B.M.; Hunter, M.; Huppert, D.D.; Kudela, R.M.; Langlois, G.W.; Largier, J.L.; *et al.* Harmful algal blooms along the north american west coast region: History, Trends, Causes, and Impacts. *Harmful Algae* **2012**, *19*, 133–159. [[CrossRef](#)]
6. Anderson, D.; Glibert, P.; Burkholder, J. Harmful algal blooms and eutrophication: Nutrient Sources, Composition, and Consequences. *Estuaries* **2002**, *25*, 704–726. [[CrossRef](#)]
7. O'Bannon, C.; Carr, J.; Seekell, D.A.; D'Odorico, P. Globalization of agricultural pollution due to international trade. *Hydrol. Earth Syst. Sci.* **2014**, *18*, 503–510. [[CrossRef](#)]
8. Galloway, J.N.; Townsend, A.R.; Erisman, J.W.; Bekunda, M.; Cai, Z.; Freney, J.R.; Martinelli, L.A.; Seitzinger, S.P.; Sutton, M.A. Transformation of the nitrogen cycle: Recent Trends, Questions, and Potential Solutions. *Science* **2008**, *320*, 889–892. [[CrossRef](#)] [[PubMed](#)]
9. Naylor, R.; Steinfeld, H.; Falcon, W.; Galloway, J.; Smil, V.; Bradford, E.; Alder, J.; Mooney, H. Losing the links between livestock and land. *Science* **2005**, *310*, 1621–1622. [[CrossRef](#)] [[PubMed](#)]

10. Hoekstra, A.Y.; Chapagain, A.K. *Globalization of Water: Sharing the Planet's Freshwater Resources*; Blackwell: Oxford, UK, 2008.
11. Steen-Olsen, K.; Weinzettel, J.; Cranston, G.; Ercin, A.E.; Hertwich, E.G. Carbon, land, and water footprint accounts for the European Union: Consumption, Production, and Displacements through International Trade. *Environ. Sci. Technol.* **2012**, *46*, 10883–10891. [[CrossRef](#)] [[PubMed](#)]
12. Hoekstra, A.Y.; Mekonnen, M.M. The water footprint of humanity. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 3232–3237. [[CrossRef](#)] [[PubMed](#)]
13. Wackernagel, M.; Onisto, L.; Bello, P.; Callejas Linares, A.; Susana López Falfán, I.; Méndez García, J.; Isabel Suárez Guerrero, A.; Guadalupe Suárez Guerrero, M. National natural capital accounting with the ecological footprint concept. *Ecol. Econ.* **1999**, *29*, 375–390. [[CrossRef](#)]
14. Leach, A.M.; Galloway, J.N.; Bleeker, A.; Erisman, J.W.; Kohn, R.; Kitzes, J. A nitrogen footprint model to help consumers understand their role in nitrogen losses to the environment. *Environ. Dev.* **2012**, *1*, 40–66. [[CrossRef](#)]
15. Hoekstra, A.Y.; Chapagain, A.K.; Aldaya, M.M.; Mekonnen, M.M. *The Water Footprint Assessment Manual: Setting the Global Standard*; Earthscan: London, UK, 2011.
16. Tukker, A.; Bulavskaya, T.; Giljum, S.; de Koning, A.; Lutter, S.; Simas, M.; Stadler, K.; Wood, R. *The Global Resource Footprint of Nations. Carbon, Water, Land and Materials Embodied in Trade and Final Consumption Calculated with Exiobase 2.1*; The Netherlands Organisation for Applied Scientific Research/Leiden University/Vienna University of Economics and Business/Norwegian University of Science and Technology: Leiden/Delft, The Netherlands; Vienna, Austria; Trondheim, Norway, 2014.
17. De Willigen, P. *An Analysis of the Calculation of Leaching and Denitrification Losses as Practised in the Nutmon Approach*; Plant Research International: Wageningen, The Netherlands, 2000; Volume 18.
18. Batjes, N.H. *Isric-Wise Derived Soil Properties on a 5 by 5 Arc-Min Global Grid (ver 1.2b)*; ISRIC—World Soil Information: Wageningen, The Netherlands, 2012.
19. Mitchell, T.D.; Jones, P.D. An improved method of constructing a database of monthly climate observations and associated high-resolution grids. *Int. J. Climatol.* **2005**, *25*, 693–712. [[CrossRef](#)]
20. Allen, R.G.; Pereira, L.S.; Raes, D.; Smith, M. *Crop. Evapotranspiration: Guideline for Computing Crop Water Requirement*; Food and Agriculture Organization: Rome, Italy, 1998.
21. Smaling, E.M.A.; Roscoe, R.; Lesschen, J.P.; Bouwman, A.F.; Comunello, E. From forest to waste: Assessment of the Brazilian Soybean Chain, Using Nitrogen as a Marker. *Agric. Ecosyst. Environ.* **2008**, *128*, 185–197. [[CrossRef](#)]
22. FAO. *Scaling Soil Nutrient Balances*; Food and Agriculture Organization: Rome, Italy, 2004; Volume 15.
23. Liu, J.; You, L.; Amini, M.; Obersteiner, M.; Herrero, M.; Zehnder, A.J.B.; Yang, H. A high-resolution assessment on global nitrogen flows in cropland. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 8035–8040. [[CrossRef](#)] [[PubMed](#)]
24. Bouwman, L.; Goldewijk, K.K.; van der Hoek, K.W.; Beusen, A.H.W.; van Vuuren, D.P.; Willems, J.; Rufino, M.C.; Stehfest, E. Exploring global changes in nitrogen and phosphorus cycles in agriculture induced by livestock production over the 1900–2050 period. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 20882–20887. [[CrossRef](#)] [[PubMed](#)]
25. Seitzinger, S.P.; Mayorga, E.; Bouwman, A.F.; Kroeze, C.; Beusen, A.H.W.; Billen, G.; van Drecht, G.; Dumont, E.; Fekete, B.M.; Garnier, J.; et al. Global river nutrient export: A Scenario Analysis of Past and Future Trends. *Glob. Biogeochem. Cycle* **2010**, *24*, GB0A08. [[CrossRef](#)]
26. Harrison, J.A.; Seitzinger, S.P.; Bouwman, A.F.; Caraco, N.F.; Beusen, A.H.W.; Vörösmarty, C.J. Dissolved inorganic phosphorus export to the coastal zone: Results from a Spatially Explicit, Global Model. *Glob. Biogeochem. Cycle* **2005**, *19*, GB4S03. [[CrossRef](#)]
27. IFA; IFDC; IPI; PPI; FAO. *Fertilizer Use by Crops*; Food and Agriculture Organization: Rome, Italy, 2002.
28. FAO. *Fertistat on-Line Database*; Food and Agriculture Organization: Rome, Italy, 2012.
29. Heffer, P. *Assessment of Fertilizer Use by Crop at the Global Level 2006/07–2007/08*; International Fertilizer Industry Association: Paris, France, 2009.
30. FAO. *Faostat on-Line Database*; Food and Agriculture Organization: Rome, Italy, 2012.
31. FAO. Global Livestock Densities. Available online: http://www.fao.org/ag/againfo/resources/en/glw/GLW_dens.html (accessed on 20 January 2016).

32. Sheldrick, W.; Keith Syers, J.; Lingard, J. Contribution of livestock excreta to nutrient balances. *Nutr. Cycl. Agroecosyst.* **2003**, *66*, 119–131. [[CrossRef](#)]
33. Mekonnen, M.M.; Hoekstra, A.Y. The green, blue and grey water footprint of crops and derived crop products. *Hydrol. Earth Syst. Sci.* **2011**, *15*, 1577–1600. [[CrossRef](#)]
34. IPNI. Ipn Estimates of Nutrient Uptake and Removal. International Plant Nutrition Institute. Available online: <http://www.ipni.net/article/IPNI-3296> (accessed on 12 July 2012).
35. Roy, R.N.; Finck, A.; Blair, G.J.; Tandon, H.L.S. *Plant. Nutrition for Food Security a Guide for Integrated Nutrient Management*; Food and Agriculture Organization: Rome, Italy, 2006; Volume 16.
36. Ravindranath, N.H.; Somashekar, H.I.; Nagaraja, M.S.; Sudha, P.; Sangeetha, G.; Bhattacharya, S.C.; Abdul Salam, P. Assessment of sustainable non-plantation biomass resources potential for energy in india. *Biomass Bioenergy* **2005**, *29*, 178–190. [[CrossRef](#)]
37. Graham, R.L.; Nelson, R.; Sheehan, J.; Perlack, R.D.; Wright, L.L. Current and potential U.S. Corn stover supplies. *Agron. J.* **2007**, *99*, 1–11. [[CrossRef](#)]
38. Nelson, R.G. Resource assessment and removal analysis for corn stover and wheat straw in the eastern and midwestern united states—Rainfall and wind-induced soil erosion methodology. *Biomass Bioenergy* **2002**, *22*, 349–363. [[CrossRef](#)]
39. Perlack, R.D.; Wright, L.L.; Turhollow, A.F.; Graham, R.L.; Stokes, B.J.; Erbach, D.C. *Biomass as Feedstock for a Bioenergy and Bioproducts Industry: The Technical Feasibility of a Billion-Ton Annual Supply*; DOE/GO-102005-2135ORNL/TM-2005/66; Oak Ridge National Laboratory, US Department of Energy: Oak Ridge, TN, USA, 2005.
40. Krausmann, F.; Erb, K.H.; Gingrich, S.; Lauk, C.; Haberl, H. Global patterns of socioeconomic biomass flows in the year 2000: A Comprehensive Assessment of Supply, Consumption And Constraints. *Ecol. Econ.* **2008**, *65*, 471–487. [[CrossRef](#)]
41. Liu, C.; Kroeze, C.; Hoekstra, A.Y.; Gerbens-Leenes, W. Past and future trends in grey water footprints of anthropogenic nitrogen and phosphorus inputs to major world rivers. *Ecol. Indic.* **2012**, *18*, 42–49. [[CrossRef](#)]
42. Leip, A.; Achermann, B.; Billen, G.; Bleeker, A.; Bouwman, A.F.; de Vries, W.; Dragosits, U.; Döring, U.; Fernall, D.; Geupel, M.; *et al.* Integrating nitrogen fluxes at the european scale. In *The European Nitrogen Assessment*; Sutton, M.A., Howard, C.M., Erisman, J.W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., Grizzetti, B., Eds.; Cambridge University Press: Cambridge, UK, 2011; pp. 345–376.
43. Westhoek, H.; Lesschen, J.P.; Rood, T.; Wagner, S.; de Marco, A.; Murphy-Bokern, D.; Leip, A.; van Grinsven, H.; Sutton, M.A.; Oenema, O. Food choices, health and environment: Effects of Cutting Europe's Meat and Dairy Intake. *Glob. Environ. Chang.* **2014**, *26*, 196–205. [[CrossRef](#)]
44. Leip, A.; Weiss, F.; Lesschen, J.P.; Westhoek, H. The nitrogen footprint of food products in the european union. *J. Agric. Sci.* **2014**, *152*, 20–33. [[CrossRef](#)]
45. Grizzetti, B.; Bouraoui, F.; Aloe, A. Changes of nitrogen and phosphorus loads to european seas. *Glob. Change Biol.* **2012**, *18*, 769–782. [[CrossRef](#)]
46. Mekonnen, M.M.; Hoekstra, A.Y. Global gray water footprint and water pollution levels related to anthropogenic nitrogen loads to fresh water. *Environ. Sci. Technol.* **2015**, *49*, 12860–12868. [[CrossRef](#)] [[PubMed](#)]
47. Vanham, D.; Bidoglio, G. A review on the indicator water footprint for the eu28. *Ecol. Indic.* **2013**, *26*, 61–75. [[CrossRef](#)]

