

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Kenneth Bloom Publications

Research Papers in Physics and Astronomy

---

12-17-2001

## Search for Gluinos and Squarks Using Like-Sign Dileptons in $p\bar{p}$ Collisions at $\sqrt{s}= 1.8$ TeV

T. Affolder

*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California*

Kenneth A. Bloom

*University of Nebraska-Lincoln, kbloom2@unl.edu*

Collider Detector at Fermilab Collaboration

Follow this and additional works at: <https://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

---

Affolder, T.; Bloom, Kenneth A.; and Fermilab Collaboration, Collider Detector at, "Search for Gluinos and Squarks Using Like-Sign Dileptons in  $p\bar{p}$  Collisions at  $\sqrt{s}= 1.8$  TeV" (2001). *Kenneth Bloom Publications*. 78.

<https://digitalcommons.unl.edu/physicsbloom/78>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Search for Gluinos and Squarks Using Like-Sign Dileptons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV

T. Affolder,<sup>23</sup> H. Akimoto,<sup>45</sup> A. Akopian,<sup>37</sup> M. G. Albrow,<sup>11</sup> P. Amaral,<sup>8</sup> D. Amidei,<sup>25</sup> K. Anikeev,<sup>24</sup> J. Antos,<sup>1</sup> G. Apollinari,<sup>11</sup> T. Arisawa,<sup>45</sup> A. Artikov,<sup>9</sup> T. Asakawa,<sup>43</sup> W. Ashmanskas,<sup>8</sup> F. Afzar,<sup>30</sup> P. Azzi-Bacchetta,<sup>31</sup> N. Bacchetta,<sup>31</sup> H. Bachacou,<sup>23</sup> S. Bailey,<sup>16</sup> P. de Barbaro,<sup>36</sup> A. Barbaro-Galtieri,<sup>23</sup> V. E. Barnes,<sup>35</sup> B. A. Barnett,<sup>19</sup> S. Baroiant,<sup>5</sup> M. Barone,<sup>13</sup> G. Bauer,<sup>24</sup> F. Bedeschi,<sup>33</sup> S. Belforte,<sup>42</sup> W. H. Bell,<sup>15</sup> G. Bellettini,<sup>33</sup> J. Bellinger,<sup>46</sup> D. Benjamin,<sup>10</sup> J. Bensinger,<sup>4</sup> A. Beretvas,<sup>11</sup> J. P. Berge,<sup>11</sup> J. Berryhill,<sup>8</sup> A. Bhatti,<sup>37</sup> M. Binkley,<sup>11</sup> D. Bisello,<sup>31</sup> M. Bishai,<sup>11</sup> R. E. Blair,<sup>2</sup> C. Blocker,<sup>4</sup> K. Bloom,<sup>25</sup> B. Blumenfeld,<sup>19</sup> S. R. Blusk,<sup>36</sup> A. Bocci,<sup>37</sup> A. Bodek,<sup>36</sup> W. Bokhari,<sup>32</sup> G. Bolla,<sup>35</sup> Y. Bonushkin,<sup>6</sup> D. Bortoletto,<sup>35</sup> J. Boudreau,<sup>34</sup> A. Brandl,<sup>27</sup> S. van den Brink,<sup>19</sup> C. Bromberg,<sup>26</sup> M. Brozovic,<sup>10</sup> E. Brubaker,<sup>23</sup> N. Bruner,<sup>27</sup> E. Buckley-Geer,<sup>11</sup> J. Budagov,<sup>9</sup> H. S. Budd,<sup>36</sup> K. Burkett,<sup>16</sup> G. Busetto,<sup>31</sup> A. Byon-Wagner,<sup>11</sup> K. L. Byrum,<sup>2</sup> S. Cabrera,<sup>10</sup> P. Calafiura,<sup>23</sup> M. Campbell,<sup>25</sup> W. Carithers,<sup>23</sup> J. Carlson,<sup>25</sup> D. Carlsmith,<sup>46</sup> W. Caskey,<sup>5</sup> A. Castro,<sup>3</sup> D. Cauz,<sup>42</sup> A. Cerri,<sup>33</sup> A. W. Chan,<sup>1</sup> P. S. Chang,<sup>1</sup> P. T. Chang,<sup>1</sup> J. Chapman,<sup>25</sup> C. Chen,<sup>32</sup> Y. C. Chen,<sup>1</sup> M.-T. Cheng,<sup>1</sup> M. Chertok,<sup>5</sup> G. Chiarelli,<sup>33</sup> I. Chirikov-Zorin,<sup>9</sup> G. Chlachidze,<sup>9</sup> F. Chlebana,<sup>11</sup> L. Christofek,<sup>18</sup> M. L. Chu,<sup>1</sup> Y. S. Chung,<sup>36</sup> C. I. Ciobanu,<sup>28</sup> A. G. Clark,<sup>14</sup> A. Connolly,<sup>23</sup> J. Conway,<sup>38</sup> M. Cordelli,<sup>13</sup> J. Cranshaw,<sup>40</sup> R. Cropp,<sup>41</sup> R. Culbertson,<sup>11</sup> D. Dagenhart,<sup>44</sup> S. D'Auria,<sup>15</sup> F. DeJongh,<sup>11</sup> S. Dell'Agnello,<sup>13</sup> M. Dell'Orso,<sup>33</sup> L. Demortier,<sup>37</sup> M. Deninno,<sup>3</sup> P. F. Derwent,<sup>11</sup> T. Devlin,<sup>38</sup> J. R. Dittmann,<sup>11</sup> A. Dominguez,<sup>23</sup> S. Donati,<sup>33</sup> J. Done,<sup>39</sup> M. D'Onofrio,<sup>33</sup> T. Dorigo,<sup>16</sup> N. Eddy,<sup>18</sup> K. Einsweiler,<sup>23</sup> J. E. Elias,<sup>11</sup> E. Engels, Jr.,<sup>34</sup> R. Erbacher,<sup>11</sup> D. Errede,<sup>18</sup> S. Errede,<sup>18</sup> Q. Fan,<sup>36</sup> R. G. Feild,<sup>47</sup> J. P. Fernandez,<sup>11</sup> C. Ferretti,<sup>33</sup> R. D. Field,<sup>12</sup> I. Fiori,<sup>3</sup> B. Flaughner,<sup>11</sup> G. W. Foster,<sup>11</sup> M. Franklin,<sup>16</sup> J. Freeman,<sup>11</sup> J. Friedman,<sup>24</sup> Y. Fukui,<sup>22</sup> I. Furic,<sup>24</sup> S. Galeotti,<sup>33</sup> A. Gallas,<sup>16,\*</sup> M. Gallinaro,<sup>37</sup> T. Gao,<sup>32</sup> M. Garcia-Sciveres,<sup>23</sup> A. F. Garfinkel,<sup>35</sup> P. Gatti,<sup>31</sup> C. Gay,<sup>47</sup> D. W. Gerdes,<sup>25</sup> P. Giannetti,<sup>33</sup> P. Giromini,<sup>13</sup> V. Glagolev,<sup>9</sup> D. Glenzinski,<sup>11</sup> M. Gold,<sup>27</sup> J. Goldstein,<sup>11</sup> I. Gorelov,<sup>27</sup> A. T. Goshaw,<sup>10</sup> Y. Gotra,<sup>34</sup> K. Goulianos,<sup>37</sup> C. Green,<sup>35</sup> G. Grim,<sup>5</sup> P. Gris,<sup>11</sup> L. Groer,<sup>38</sup> C. Grosso-Pilcher,<sup>8</sup> M. Guenther,<sup>35</sup> G. Guillian,<sup>25</sup> J. Guimaraes da Costa,<sup>16</sup> R. M. Haas,<sup>12</sup> C. Haber,<sup>23</sup> S. R. Hahn,<sup>11</sup> C. Hall,<sup>16</sup> T. Handa,<sup>17</sup> R. Handler,<sup>46</sup> W. Hao,<sup>40</sup> F. Happacher,<sup>13</sup> K. Hara,<sup>43</sup> A. D. Hardman,<sup>35</sup> R. M. Harris,<sup>11</sup> F. Hartmann,<sup>20</sup> K. Hatakeyama,<sup>37</sup> J. Hauser,<sup>6</sup> J. Heinrich,<sup>32</sup> A. Heiss,<sup>20</sup> M. Herndon,<sup>19</sup> C. Hill,<sup>5</sup> K. D. Hoffman,<sup>35</sup> C. Holck,<sup>32</sup> R. Hollebeck,<sup>32</sup> L. Holloway,<sup>18</sup> R. Hughes,<sup>28</sup> J. Huston,<sup>26</sup> J. Huth,<sup>16</sup> H. Ikeda,<sup>43</sup> J. Incandela,<sup>11</sup> G. Introzzi,<sup>33</sup> J. Iwai,<sup>45</sup> Y. Iwata,<sup>17</sup> E. James,<sup>25</sup> M. Jones,<sup>32</sup> U. Joshi,<sup>11</sup> H. Kambara,<sup>14</sup> T. Kamon,<sup>39</sup> T. Kaneko,<sup>43</sup> K. Karr,<sup>44</sup> H. Kasha,<sup>47</sup> Y. Kato,<sup>29</sup> T. A. Keaffaber,<sup>35</sup> K. Kelley,<sup>24</sup> M. Kelly,<sup>25</sup> R. D. Kennedy,<sup>11</sup> R. Kephart,<sup>11</sup> D. Khazins,<sup>10</sup> T. Kikuchi,<sup>43</sup> B. Kilminster,<sup>36</sup> B. J. Kim,<sup>21</sup> D. H. Kim,<sup>21</sup> H. S. Kim,<sup>18</sup> M. J. Kim,<sup>21</sup> S. B. Kim,<sup>21</sup> S. H. Kim,<sup>43</sup> Y. K. Kim,<sup>23</sup> M. Kirby,<sup>10</sup> M. Kirk,<sup>4</sup> L. Kirsch,<sup>4</sup> S. Klimentenko,<sup>12</sup> P. Koehn,<sup>28</sup> K. Kondo,<sup>45</sup> J. Konigsberg,<sup>12</sup> A. Korn,<sup>24</sup> A. Korytov,<sup>12</sup> E. Kovacs,<sup>2</sup> J. Kroll,<sup>32</sup> M. Kruse,<sup>10</sup> S. E. Kuhlmann,<sup>2</sup> K. Kurino,<sup>17</sup> T. Kuwabara,<sup>43</sup> A. T. Laasanen,<sup>35</sup> N. Lai,<sup>8</sup> S. Lami,<sup>37</sup> S. Lammel,<sup>11</sup> J. Lancaster,<sup>10</sup> M. Lancaster,<sup>23</sup> R. Lander,<sup>5</sup> A. Lath,<sup>38</sup> G. Latino,<sup>33</sup> T. LeCompte,<sup>2</sup> A. M. Lee IV,<sup>10</sup> K. Lee,<sup>40</sup> S. W. Lee,<sup>39</sup> S. Leone,<sup>33</sup> J. D. Lewis,<sup>11</sup> M. Lindgren,<sup>6</sup> T. M. Liss,<sup>18</sup> J. B. Liu,<sup>36</sup> Y. C. Liu,<sup>1</sup> D. O. Litvintsev,<sup>11</sup> O. Lobban,<sup>40</sup> N. Lockyer,<sup>32</sup> J. Loken,<sup>30</sup> M. Loreti,<sup>31</sup> D. Lucchesi,<sup>31</sup> P. Lukens,<sup>11</sup> S. Lusin,<sup>46</sup> L. Lyons,<sup>30</sup> J. Lys,<sup>23</sup> R. Madrak,<sup>16</sup> K. Maeshima,<sup>11</sup> P. Maksimovic,<sup>16</sup> L. Malferrari,<sup>3</sup> M. Mangano,<sup>33</sup> M. Mariotti,<sup>31</sup> G. Martignon,<sup>31</sup> A. Martin,<sup>47</sup> J. A. J. Matthews,<sup>27</sup> J. Mayer,<sup>41</sup> P. Mazzanti,<sup>3</sup> K. S. McFarland,<sup>36</sup> P. McIntyre,<sup>39</sup> E. McKigney,<sup>32</sup> M. Menguzzato,<sup>31</sup> A. Menzione,<sup>33</sup> C. Mesropian,<sup>37</sup> A. Meyer,<sup>11</sup> T. Miao,<sup>11</sup> R. Miller,<sup>26</sup> J. S. Miller,<sup>25</sup> H. Minato,<sup>43</sup> S. Miscetti,<sup>13</sup> M. Mishina,<sup>22</sup> G. Mitselmakher,<sup>12</sup> N. Moggi,<sup>3</sup> E. Moore,<sup>27</sup> R. Moore,<sup>25</sup> Y. Morita,<sup>22</sup> T. Moulik,<sup>35</sup> M. Mulhearn,<sup>24</sup> A. Mukherjee,<sup>11</sup> T. Muller,<sup>20</sup> A. Munar,<sup>33</sup> P. Murat,<sup>11</sup> S. Murgia,<sup>26</sup> J. Nachtman,<sup>6</sup> V. Nagaslaev,<sup>40</sup> S. Nahn,<sup>47</sup> H. Nakada,<sup>43</sup> I. Nakano,<sup>17</sup> C. Nelson,<sup>11</sup> T. Nelson,<sup>11</sup> C. Neu,<sup>28</sup> D. Neuberger,<sup>20</sup> C. Newman-Holmes,<sup>11</sup> C.-Y. P. Ngan,<sup>24</sup> H. Niu,<sup>4</sup> L. Nodulman,<sup>2</sup> A. Nomerotski,<sup>12</sup> S. H. Oh,<sup>10</sup> Y. D. Oh,<sup>21</sup> T. Ohmoto,<sup>17</sup> T. Ohsugi,<sup>17</sup> R. Oishi,<sup>43</sup> T. Okusawa,<sup>29</sup> J. Olsen,<sup>46</sup> W. Orejudos,<sup>23</sup> C. Pagliarone,<sup>33</sup> F. Palmonari,<sup>33</sup> R. Paoletti,<sup>33</sup> V. Papadimitriou,<sup>40</sup> D. Partos,<sup>4</sup> J. Patrick,<sup>11</sup> G. Pauletta,<sup>42</sup> M. Paulini,<sup>23,†</sup> C. Paus,<sup>24</sup> D. Pellett,<sup>5</sup> L. Pescara,<sup>31</sup> T. J. Phillips,<sup>10</sup> G. Piacentino,<sup>33</sup> K. T. Pitts,<sup>18</sup> A. Pompos,<sup>35</sup> L. Pondrom,<sup>46</sup> G. Pope,<sup>34</sup> M. Popovic,<sup>41</sup> F. Prokoshin,<sup>9</sup> J. Proudfoot,<sup>2</sup> F. Ptohos,<sup>13</sup> O. Pukhov,<sup>9</sup> G. Punzi,<sup>33</sup> A. Rakitine,<sup>24</sup> F. Ratnikov,<sup>38</sup> D. Reher,<sup>23</sup> A. Reichold,<sup>30</sup> A. Ribon,<sup>31</sup> W. Riegler,<sup>16</sup> F. Rimondi,<sup>3</sup> L. Ristori,<sup>33</sup> M. Riveline,<sup>41</sup> W. J. Robertson,<sup>10</sup> A. Robinson,<sup>41</sup> T. Rodrigo,<sup>7</sup> S. Rolli,<sup>44</sup> L. Rosenson,<sup>24</sup> R. Roser,<sup>11</sup> R. Rossin,<sup>31</sup> A. Roy,<sup>35</sup> A. Ruiz,<sup>7</sup> A. Safonov,<sup>5</sup> R. St. Denis,<sup>15</sup> W. K. Sakumoto,<sup>36</sup> D. Saltzberg,<sup>6</sup> C. Sanchez,<sup>28</sup> A. Sansoni,<sup>13</sup> L. Santi,<sup>42</sup> H. Sato,<sup>43</sup> P. Savard,<sup>41</sup> P. Schlabach,<sup>11</sup> E. E. Schmidt,<sup>11</sup> M. P. Schmidt,<sup>47</sup> M. Schmitt,<sup>16,\*</sup> L. Scodellaro,<sup>31</sup> A. Scott,<sup>6</sup> A. Scribano,<sup>33</sup> S. Segler,<sup>11</sup> S. Seidel,<sup>27</sup> Y. Seiya,<sup>43</sup> A. Semenov,<sup>9</sup> F. Semeria,<sup>3</sup> T. Shah,<sup>24</sup> M. D. Shapiro,<sup>23</sup> P. F. Shepard,<sup>34</sup> T. Shibayama,<sup>43</sup> M. Shimojima,<sup>43</sup> M. Shochet,<sup>8</sup> A. Sidoti,<sup>31</sup> J. Siegrist,<sup>23</sup> A. Sill,<sup>40</sup>

P. Sinervo,<sup>41</sup> P. Singh,<sup>18</sup> A. J. Slaughter,<sup>47</sup> K. Sliwa,<sup>44</sup> C. Smith,<sup>19</sup> F. D. Snider,<sup>11</sup> A. Solodsky,<sup>37</sup> J. Spalding,<sup>11</sup> T. Speer,<sup>14</sup> P. Sphicas,<sup>24</sup> F. Spinella,<sup>33</sup> M. Spiropulu,<sup>16</sup> L. Spiegel,<sup>11</sup> J. Steele,<sup>46</sup> A. Stefanini,<sup>33</sup> J. Strologas,<sup>18</sup> F. Strumia,<sup>14</sup> D. Stuart,<sup>11</sup> K. Sumorok,<sup>24</sup> T. Suzuki,<sup>43</sup> T. Takano,<sup>29</sup> R. Takashima,<sup>17</sup> K. Takikawa,<sup>43</sup> P. Tamburello,<sup>10</sup> M. Tanaka,<sup>43</sup> B. Tannenbaum,<sup>6</sup> M. Tecchio,<sup>25</sup> R. Tesarek,<sup>11</sup> P. K. Teng,<sup>1</sup> K. Terashi,<sup>37</sup> S. Tether,<sup>24</sup> A. S. Thompson,<sup>15</sup> R. Thurman-Keup,<sup>2</sup> P. Tipton,<sup>36</sup> S. Tkaczyk,<sup>11</sup> D. Toback,<sup>39</sup> K. Tollefson,<sup>36</sup> A. Tollestrup,<sup>11</sup> D. Tonelli,<sup>33</sup> H. Toyoda,<sup>29</sup> W. Trischuk,<sup>41</sup> J. F. de Troconiz,<sup>16</sup> J. Tseng,<sup>24</sup> N. Turini,<sup>33</sup> F. Ukegawa,<sup>43</sup> T. Vaiciulis,<sup>36</sup> J. Valls,<sup>38</sup> S. Vejcik III,<sup>11</sup> G. Velez,<sup>11</sup> G. Veramendi,<sup>23</sup> R. Vidal,<sup>11</sup> I. Vila,<sup>7</sup> R. Vilar,<sup>7</sup> I. Volobouev,<sup>23</sup> M. von der Mey,<sup>6</sup> D. Vucinic,<sup>24</sup> R. G. Wagner,<sup>2</sup> R. L. Wagner,<sup>11</sup> N. B. Wallace,<sup>38</sup> Z. Wan,<sup>38</sup> C. Wang,<sup>10</sup> M. J. Wang,<sup>1</sup> B. Ward,<sup>15</sup> S. Waschke,<sup>15</sup> T. Watanabe,<sup>43</sup> D. Waters,<sup>30</sup> T. Watts,<sup>38</sup> R. Webb,<sup>39</sup> H. Wenzel,<sup>20</sup> W. C. Wester III,<sup>11</sup> A. B. Wicklund,<sup>2</sup> E. Wicklund,<sup>11</sup> T. Wilkes,<sup>5</sup> H. H. Williams,<sup>32</sup> P. Wilson,<sup>11</sup> B. L. Winer,<sup>28</sup> D. Winn,<sup>25</sup> S. Wolbers,<sup>11</sup> D. Wolinski,<sup>25</sup> J. Wolinski,<sup>26</sup> S. Wolinski,<sup>25</sup> S. Worm,<sup>27</sup> X. Wu,<sup>14</sup> J. Wyss,<sup>33</sup> W. Yao,<sup>23</sup> G. P. Yeh,<sup>11</sup> P. Yeh,<sup>1</sup> J. Yoh,<sup>11</sup> C. Yosef,<sup>26</sup> T. Yoshida,<sup>29</sup> I. Yu,<sup>21</sup> S. Yu,<sup>32</sup> Z. Yu,<sup>47</sup> A. Zanetti,<sup>42</sup> F. Zetti,<sup>23</sup> and S. Zucchelli<sup>3</sup>  
(CDF Collaboration)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439*

<sup>3</sup>*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*

<sup>4</sup>*Brandeis University, Waltham, Massachusetts 02254*

<sup>5</sup>*University of California at Davis, Davis, California 95616*

<sup>6</sup>*University of California at Los Angeles, Los Angeles, California 90024*

<sup>7</sup>*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

<sup>8</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

<sup>9</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

<sup>10</sup>*Duke University, Durham, North Carolina 27708*

<sup>11</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

<sup>12</sup>*University of Florida, Gainesville, Florida 32611*

<sup>13</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

<sup>14</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*

<sup>15</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*

<sup>16</sup>*Harvard University, Cambridge, Massachusetts 02138*

<sup>17</sup>*Hiroshima University, Higashi-Hiroshima 724, Japan*

<sup>18</sup>*University of Illinois, Urbana, Illinois 61801*

<sup>19</sup>*The Johns Hopkins University, Baltimore, Maryland 21218*

<sup>20</sup>*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*

<sup>21</sup>*Center for High Energy Physics, Kyungpook National University, Taegu 702-701, Korea*

*and SungKyunKwan University, Suwon 440-746, Korea*

<sup>22</sup>*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan*

<sup>23</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720*

<sup>24</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

<sup>25</sup>*University of Michigan, Ann Arbor, Michigan 48109*

<sup>26</sup>*Michigan State University, East Lansing, Michigan 48824*

<sup>27</sup>*University of New Mexico, Albuquerque, New Mexico 87131*

<sup>28</sup>*The Ohio State University, Columbus, Ohio 43210*

<sup>29</sup>*Osaka City University, Osaka 588, Japan*

<sup>30</sup>*University of Oxford, Oxford OX1 3RH, United Kingdom*

<sup>31</sup>*Universita di Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova, I-35131 Padova, Italy*

<sup>32</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104*

<sup>33</sup>*Istituto Nazionale di Fisica Nucleare, University and Scuola Normale Superiore of Pisa, I-56100 Pisa, Italy*

<sup>34</sup>*University of Pittsburgh, Pittsburgh, Pennsylvania 15260*

<sup>35</sup>*Purdue University, West Lafayette, Indiana 47907*

<sup>36</sup>*University of Rochester, Rochester, New York 14627*

<sup>37</sup>*Rockefeller University, New York, New York 10021*

<sup>38</sup>*Rutgers University, Piscataway, New Jersey 08855*

<sup>39</sup>*Texas A&M University, College Station, Texas 77843*

<sup>40</sup>*Texas Tech University, Lubbock, Texas 79409*

<sup>41</sup>*Institute of Particle Physics, University of Toronto, Toronto, Canada M5S 1A7*

<sup>42</sup>*Istituto Nazionale di Fisica Nucleare, University of Trieste, Udine, Italy*

<sup>43</sup>*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*

<sup>44</sup>*Tufts University, Medford, Massachusetts 02155*<sup>45</sup>*Waseda University, Tokyo 169, Japan*<sup>46</sup>*University of Wisconsin, Madison, Wisconsin 53706*<sup>47</sup>*Yale University, New Haven, Connecticut 06520*

(Received 18 June 2001; published 30 November 2001)

We present results of the first search for like-sign dilepton ( $e^+e^+$ ,  $e^+\mu^\pm$ ,  $\mu^\pm\mu^\pm$ ) events associated with multijets and large missing energy using  $106 \text{ pb}^{-1}$  of data in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8 \text{ TeV}$  collected during 1992–1995 by the CDF experiment. Finding no events that pass our selection, we examine pair production of gluinos ( $\tilde{g}$ ) and squarks ( $\tilde{q}$ ) in a constrained framework of the minimal supersymmetric standard model. At  $\tan\beta = 2$  and  $\mu = -800 \text{ GeV}/c^2$ , we set 95% confidence level limits of  $M_{\tilde{g}} > 221 \text{ GeV}/c^2$  for  $M_{\tilde{g}} = M_{\tilde{q}}$ , and  $M_{\tilde{g}} > 168 \text{ GeV}/c^2$  for  $M_{\tilde{q}} \gg M_{\tilde{g}}$ , both with small variation as a function of  $\mu$ .

DOI: 10.1103/PhysRevLett.87.251803

PACS numbers: 14.80.Ly, 11.30.Pb, 12.60.Jv, 13.85.Rm

The standard model (SM) of particle physics is enormously successful in explaining a wide variety of phenomena. In spite of this, there are a number of structural defects in the model, such as the quadratic mass divergence of the Higgs boson. Supersymmetry (SUSY) provides a promising solution and in the minimal supersymmetric standard model (MSSM) [1] each SM particle has a SUSY partner which is required to be lighter than or of the order of  $1 \text{ TeV}/c^2$  [1]. Conservation of  $R$  parity [2] requires SUSY particles to be produced in pairs and the lightest SUSY particle (LSP) to be stable.

At the Fermilab Tevatron, pair production and sequential decays of supersymmetric quarks (squarks,  $\tilde{q}$ ) and supersymmetric gluons (gluinos,  $\tilde{g}$ ) can result in events with final state leptons. The  $\tilde{q}$  can decay to the lightest chargino ( $\tilde{\chi}_1^\pm$ ) or the next-to-lightest neutralino ( $\tilde{\chi}_2^0$ ) via  $\tilde{q} \rightarrow q'\tilde{\chi}_1^\pm$  or  $\tilde{q} \rightarrow q\tilde{\chi}_2^0$ , and the  $\tilde{q} \rightarrow q\tilde{g}$  decay occurs when kinematically allowed. The decays of the  $\tilde{g}$  are  $\tilde{g} \rightarrow q\tilde{q}'\tilde{\chi}_1^\pm$  or  $\tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_2^0$ . Each  $\tilde{q}$  and  $\tilde{g}$  decay can eventually produce isolated leptons and missing transverse energy ( $\cancel{E}_T$ ) [3] via the decays  $\tilde{\chi}_1^\pm \rightarrow \ell^\pm \nu_\ell \tilde{\chi}_1^0$  or  $\tilde{\chi}_2^0 \rightarrow \ell^+ \ell^- \tilde{\chi}_1^0$  where  $\tilde{\chi}_1^0$  is the LSP [4] which exits the detector without interacting. Thus,  $\tilde{g}\tilde{g}$ ,  $\tilde{g}\tilde{q}$ , and  $\tilde{q}\tilde{q}$  production can lead to the like-sign (LS) dilepton signatures of  $e^\pm e^\pm$ ,  $e^\pm \mu^\pm$ , and  $\mu^\pm \mu^\pm$  [5] with two or more jets and appreciable  $\cancel{E}_T$ . The fraction of dilepton events which are LS can be as large as 30% in some regions of MSSM parameter space.

The  $\ell^\pm \ell^\pm + \geq 2 \text{ jets} + \cancel{E}_T$  channel is a clean signature to search for SUSY. It has an advantage over the opposite-sign (OS) dilepton channel as there are only small SM backgrounds. Even without the  $\cancel{E}_T$  requirement the LS analysis is also useful for testing other theories beyond the SM, including  $R$  parity violating SUSY [6]. The dilepton decay channels are a natural complement to other direct searches for squarks and gluinos in the  $\cancel{E}_T$  plus multijet channel [7–12].

In this Letter, we present the results of the first search for  $\ell^\pm \ell^\pm + \geq 2 \text{ jets} + \cancel{E}_T$  events using  $106 \text{ pb}^{-1}$  of data from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8 \text{ TeV}$ . The data were collected by the Collider Detector at Fermilab (CDF) [13] during the 1992–1995 run of the Tevatron. We briefly describe the detector subsystems relevant to this analysis.

The location of the  $p\bar{p}$  collision event vertex ( $z_{\text{vertex}}$ ) is measured along the beam direction with a time projection chamber. The  $p_T$  of charged particles are measured in the region  $|\eta| < 1.1$  by a central tracking chamber (CTC) which is located in a 1.4 T solenoidal magnetic field. The momentum resolution is  $\delta p_T/p_T^2 \approx 0.001$  where  $p_T$  is measured in  $\text{GeV}/c$ . Electromagnetic and hadronic calorimeters are segmented in a projective tower geometry surrounding the solenoid and cover the region  $|\eta| < 4.2$ . A muon detector is located outside the hadron calorimeter and covers the region  $|\eta| < 1.0$ .

The analysis begins with a sample of 515 699 loosely selected dilepton events [14,15] from which we select an initial dilepton plus dijet sample. To ensure that the trigger is fully efficient, we require each event to have a lepton with  $p_T \geq 11 \text{ GeV}/c$  and  $|\eta| < 1.0$  for electrons or  $|\eta| < 0.6$  for muons. A second electron or muon is required with  $p_T \geq 5 \text{ GeV}/c$  and  $|\eta| < 1.0$ . If there are more than two isolated leptons, we take the two leading- $p_T$  leptons. Each lepton is required to be isolated such that there is no more than 4 GeV of transverse energy (measured by the calorimeter or CTC) in a cone of  $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.4$  around the direction of the lepton. To ensure that both leptons originated from the same collision event and are well measured, we require  $|z_{\text{vertex}}| \leq 60 \text{ cm}$  and  $|z_{\text{lepton}} - z_{\text{vertex}}| \leq 5 \text{ cm}$  for each lepton, where  $z_{\text{lepton}}$  is measured along the beam line. In addition to the leptons, we require two or more jets with  $E_T \geq 15 \text{ GeV}$  and  $|\eta| < 2.4$ .

Since the OS sample is used as a check of our understanding of the LS backgrounds, we place the same cuts on both samples in parallel, but with additional cuts on the OS events so as to remove events which might give a kinematic bias. To reduce the large  $J/\psi$  and  $Y$  component of the background we remove the events with  $M_{\ell\ell} < 12 \text{ GeV}/c^2$ . A total of 239 OS and 16 LS dilepton events pass the requirement.

The dominant SM backgrounds are from Drell-Yan ( $\gamma^*/Z^0$ ),  $t\bar{t}$ ,  $b\bar{b}$ ,  $c\bar{c}$ , and diboson ( $W^+W^-$ ,  $W^\pm Z^0$ ,  $Z^0 Z^0$ ) production. Each is estimated using the ISAJET Monte Carlo (MC) event generator [16] and a simulation of the CDF detector. The cross sections for  $\gamma^*/Z^0$  and  $t\bar{t}$




FIG. 1. Distributions for the dilepton + dijet data after the  $M_{\ell\ell} > 12 \text{ GeV}/c^2$  and  $Z^0$  veto requirements. (a) and (b) show the  $\cancel{E}_T$  distributions for OS and LS samples, respectively. The data (points) are compared to the standard model background (shaded line) with a SUSY contribution (solid line) for  $\tan\beta = 2$ ,  $\mu = -800 \text{ GeV}/c^2$ ,  $M_{\tilde{g}} = 210 \text{ GeV}/c^2$ , and  $M_{\tilde{q}} = 211 \text{ GeV}/c^2$ . (c) and (d) show the  $M_{\ell\ell}$  distributions in the OS and LS samples for the same requirements.

production and contributions due to  $B^0\bar{B}^0$  mixing events are normalized to CDF measurements [17–19]. We use next-to-leading order (NLO) cross sections for diboson production [20]. The contribution from  $W(\rightarrow \ell\nu\ell) + \geq 3$  jets events where one of the jets is misidentified as a lepton is found to be negligible.

Given the large  $\cancel{E}_T$  signature from SUSY, we require at least 25 GeV of  $\cancel{E}_T$  for all dilepton events. In the OS sample, we also remove all same-flavor OS dilepton events with  $76 < M_{\ell^+\ell^-} < 106 \text{ GeV}/c^2$ . Figure 1 compares the  $\cancel{E}_T$  and  $M_{\ell\ell}$  distributions for the data and the SM backgrounds for the OS and LS samples after the  $Z^0$  veto but

before the  $\cancel{E}_T$  requirement. After all cuts, we observe 19 OS (4  $ee$ , 10  $e\mu$ , 5  $\mu\mu$ ) events and no LS events in agreement with the SM expectation of  $14.1 \pm 1.3$  (stat)  $\pm 2.8$  (syst) OS events and  $0.55 \pm 0.25 \pm 0.08$  LS events. Tables I and II show a comparison of the data reduction and the SM backgrounds. We note that  $t\bar{t}$  and  $Z^0 \rightarrow \tau^+\tau^-$  are two major SM sources of the  $e\mu$  events. The 19 event sample also contains six dilepton (1  $ee$ , 5  $e\mu$ ) events out of nine  $t\bar{t} \rightarrow W^+bW^-\bar{b} \rightarrow (\ell^+\nu b)(\ell^-\bar{\nu}\bar{b})$  event candidates (1  $ee$ , 7  $e\mu$ , 1  $\mu\mu$ ) from the CDF  $t\bar{t}$  analysis in the dilepton channel [21]. The remaining three top dilepton event candidates are not in our final sample because our lepton isolation requirement for the second lepton is stricter than the top analysis. There is no evidence for new particle production.

We examine the exclusion region of  $M_{\tilde{q}}$  and  $M_{\tilde{g}}$  in a constrained framework of the MSSM. We assume five squarks ( $\tilde{u}, \tilde{d}, \tilde{s}, \tilde{c}, \tilde{b}$ ) with nearly mass-degenerate left and right helicity states. Production of top squarks is not considered even though the lighter of the two top-squark mass eigenstates can be lighter than the other squarks [22]. We impose common scalar and common gaugino masses at a grand unified theory scale as in the minimal supergravity model [23], and use the renormalization group equations [24] that relate the mass parameters, leading to a general prediction:  $M_{\tilde{q}} \geq 0.9M_{\tilde{g}}$ . To avoid a region in MSSM parameter space where there are significant branching ratios of chargino and neutralino decays into Higgs particles, the pseudoscalar Higgs mass is set to  $500 \text{ GeV}/c^2$  which is above the  $\tilde{\chi}_1^\pm$  and  $\tilde{\chi}_2^0$  masses. With these assumptions, the sensitivity of our search can be studied as a function of four parameters: the gluino mass ( $M_{\tilde{g}}$ ), the squark mass ( $M_{\tilde{q}}$ ), the ratio of the vacuum expectation values of the two Higgs fields ( $\tan\beta$ ), and the Higgs mass parameter ( $\mu$ ). Since we choose to decouple our search from the Higgs sector we scan a range of  $\mu$  that is both consistent with LEP results [9,25] and less than the SUSY mass scale:  $100 \leq |\mu| \leq 1000 \text{ GeV}/c^2$ .

The acceptance for SUSY processes is estimated by performing the final data selection on events simulated with ISAJET [16] using CTEQ3L [26] parton distribution functions (PDFs). These events are then passed through the CDF detector simulation. We define the acceptance as the ratio of the number of LS dilepton events that pass our cuts

TABLE I. A comparison of the event reduction for the data, standard model (SM) backgrounds and a model of SUSY production with  $\tan\beta = 2$ ,  $\mu = -800 \text{ GeV}/c^2$ ,  $M_{\tilde{g}} = 210 \text{ GeV}/c^2$ , and  $M_{\tilde{q}} = 211 \text{ GeV}/c^2$ .

Selection	Data	SM backgrounds	SUSY
Dilepton dataset	515,699		
Dilepton dijet	350		
$M_{\ell\ell} \geq 12 \text{ GeV}/c^2$	255	$279 \pm 9 \pm 79$	$27 \pm 1 \pm 5$
$Z^0(\rightarrow \ell^+\ell^-)$ veto	128	$158 \pm 7 \pm 45$	$27 \pm 1 \pm 5$
$\cancel{E}_T \geq 25 \text{ GeV}$	19	$14.1 \pm 1.3 \pm 2.8$	$24 \pm 1 \pm 5$
Like-sign dilepton	0	$0.55 \pm 0.25 \pm 0.08$	$5.9 \pm 0.6 \pm 1.4$

TABLE II. The expected backgrounds from standard model contributions to the final data selection after all but the LS requirement in Table I. Opposite-sign and like-sign dilepton events are listed.

Source	Opposite-sign	Like-sign
Drell-Yan	$8.7 \pm 0.9 \pm 2.5$	$0.00^{+0.01}_{-0.00} \quad ^{+0.01}_{-0.00}$
$t\bar{t}$	$4.0 \pm 0.3 \pm 1.2$	$0.08 \pm 0.04 \pm 0.02$
$b\bar{b}/c\bar{c}$	$0.9 \pm 0.9 \pm 0.3$	$0.23 \pm 0.23 \pm 0.07$
Diboson	$0.5 \pm 0.1 \pm 0.1$	$0.24 \pm 0.10 \pm 0.04$
Total	$14.1 \pm 1.3 \pm 2.8$	$0.55 \pm 0.25 \pm 0.08$
Data	19	0

to the total number of generated SUSY events which contain at least two leptons. For a nominal SUSY scenario of  $M_{\tilde{g}} = 200 \text{ GeV}/c^2$ , infinite squark mass (and hence infinite slepton mass),  $\tan\beta = 2$  and  $\mu = -800 \text{ GeV}/c^2$ , the acceptance is 1%, due mostly to the lower  $p_T$  values of the leptons. For the case where  $M_{\tilde{q}} \approx M_{\tilde{g}} \approx 200 \text{ GeV}/c^2$ , the slepton ( $\tilde{\ell}_R$ ) mass is lighter. This enhances the leptonic branching ratio due to  $\tilde{\chi}_2^0 \rightarrow \ell\tilde{\ell}_R$ , resulting in an increase of LS dilepton events in  $\tilde{g}\tilde{g}, \tilde{g}\tilde{q}, \tilde{q}\tilde{q}$  production and a rise of the overall acceptance to 3%. Table I and Fig. 1 compare the data reduction, the expectations from SM processes, and a SUSY scenario.

The total systematic uncertainty on the expected number of LS signal events comes from uncertainties on the theoretical calculation of the production cross section of gluinos and squarks, the event acceptance, and the integrated luminosity. The NLO cross section depends mainly on the choices of the QCD renormalization scale ( $Q^2$ ) and PDFs [27]. The nominal choice of  $Q^2$  is  $m^2$ , where  $m$  is  $M_{\tilde{q}}, M_{\tilde{g}}$ , or  $\frac{1}{2}\sqrt{M_{\tilde{q}}^2 + M_{\tilde{g}}^2}$  for  $\tilde{q}\tilde{q}/\tilde{q}\tilde{q}, \tilde{g}\tilde{g}$ , or  $\tilde{q}\tilde{g}$  production, respectively. The uncertainty due to the choice of  $Q^2$  is determined to be 21% by taking the larger of the variation of the cross section at  $Q^2 = (m/2)^2$  and at  $Q^2 = (2m)^2$  from the nominal cross section value. Similarly, the variation of the cross section due to the choice of PDFs yields an 8% uncertainty, estimated as the maximum deviation between the nominal choice of CTEQ3M [26] and MRS(G) [28] or GRV94HO [29]. Uncertainty on the signal acceptance is due to uncertainties on the efficiencies of the lepton trigger, identification and isolation efficiencies, as well as on the jet energy scale and the amount of gluon radiation. By varying the measured lepton trigger and identification efficiencies by 1 standard deviation, the acceptance uncertainties are estimated to be 5% and 3%, respectively. Since the lepton isolation efficiency depends on jet multiplicity, the uncertainty is estimated using  $Z^0(\rightarrow \ell^+\ell^-) + \geq 2$  jet events and is found to be 11%. By varying the jet energy scale by 1 standard deviation, we find a 5% effect on the acceptance. The uncertainty due to the initial and final state gluon radiation (ISR and FSR) is estimated by turning the ISR and/or FSR radiation off, which gives at most 7% variation in the acceptance. Enough

MC events are generated so as to keep the statistical uncertainty below 3%. The uncertainty on the luminosity is 4%. The combined uncertainty is calculated by adding all uncertainties in quadrature, and is found to be 28%.

Since no LS events pass our cuts, we calculate the upper limit on the number of SUSY events at the 95% confidence level (C.L.) using a frequentist algorithm [30] with a systematic uncertainty of 28% and no background subtraction. This corresponds to 3.46 events which we use to exclude regions in the  $M_{\tilde{q}}-M_{\tilde{g}}$  plane. Figure 2 shows the exclusion region for  $\tan\beta = 2$  and  $\mu = -800 \text{ GeV}/c^2$ . We set 95% C.L. limits at  $M_{\tilde{g}} > 168 \text{ GeV}/c^2$  for  $M_{\tilde{q}} \gg M_{\tilde{g}}$  and  $M_{\tilde{g}} > 221 \text{ GeV}/c^2$  for  $M_{\tilde{g}} \approx M_{\tilde{q}}$ . These results are better than the previous limits from complementary searches by about  $5 \text{ GeV}/c^2$  [10,11].

We examine the dependence of the mass limit as  $\tan\beta$  and  $\mu$  are varied in the region  $M_{\tilde{g}} \approx M_{\tilde{q}}$ . For  $\mu = -800 \text{ GeV}/c^2$ , the variation in the mass limit is smaller than 2% in the range of  $\tan\beta$  between 1.7 and 10 if the mixings of the third generation SUSY particles (especially  $\tilde{\tau}$ ) are minimal. In the case of maximal  $\tilde{\tau}$  mixing, the mass limit remains the same for  $\tan\beta$  up to about 3. For  $\tan\beta = 2$ , the limit deviates by at most 3.6% from the  $221 \text{ GeV}/c^2$  limit in the range  $\mu \leq -150 \text{ GeV}/c^2$ , while the limits in  $\mu \geq 150 \text{ GeV}/c^2$  are systematically 8%–12% lower.

In conclusion, we have searched for new physics using LS dilepton events in association with two or more jets and  $\cancel{E}_T$  in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.8 \text{ TeV}$ . Production of both OS and LS dilepton events is consistent with the SM expectations. Within a framework of constrained MSSM




FIG. 2. Limit in the  $M_{\tilde{q}}-M_{\tilde{g}}$  plane at the 95% confidence level for a constrained MSSM scenario ( $M_{\tilde{q}} \geq 0.9M_{\tilde{g}}$ ) for  $\tan\beta = 2$  and  $\mu = -800 \text{ GeV}/c^2$ . The results of other direct, but complementary, searches are also presented [8–11].

(five degenerate squarks,  $M_{\tilde{q}} \geq 0.9M_{\tilde{g}}$ ), for small  $\tan\beta$  we set mass limits of  $M_{\tilde{g}} > 168 \text{ GeV}/c^2$  for  $M_{\tilde{q}} \gg M_{\tilde{g}}$ , and  $M_{\tilde{g}} > 221 \text{ GeV}/c^2$  for  $M_{\tilde{q}} \approx M_{\tilde{g}}$ , both with small variation as a function of  $\mu$ .

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and the National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Science, Sports and Culture of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A. P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korea Science and Engineering Foundation (KoSEF); the Korea Research Foundation; and the Comision Interministerial de Ciencia y Tecnologia, Spain.

\*Now at Northwestern University, Evanston, IL 60208.

†Now at Carnegie Mellon University, Pittsburgh, PA 15213.

- [1] H. P. Nilles, Phys. Rep. **110**, 1 (1984); H. E. Haber and G. L. Kane, *ibid.* **117**, 75 (1985).
- [2] A. Salam and J. Strathdee, Nucl. Phys. **B76**, 477 (1974); P. Fayet, *ibid.* **B90**, 104 (1975); G. Farrar and P. Fayet, Phys. Lett. **76B**, 575 (1978).
- [3] We use a coordinate system where  $\theta$  and  $\phi$  are the polar and azimuthal angles, respectively, with respect to the proton beam direction ( $z$  axis). The pseudorapidity  $\eta$  is defined as  $-\ln[\tan(\theta/2)]$ . The transverse momentum of a particle is denoted as  $p_T = p \sin\theta$ . The missing transverse energy,  $\cancel{E}_T$ , is the magnitude of  $\vec{\cancel{E}}_T \equiv -\sum E_T^i \hat{n}_i$ , where  $\hat{n}_i$  is the unit vector in the transverse plane pointing from the interaction point to the energy deposition in calorimeter cell  $i$ .
- [4] H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983); J. Ellis *et al.*, Nucl. Phys. **238B**, 453 (1984).
- [5] H. Baer *et al.*, Phys. Lett. **161B**, 175 (1985).
- [6] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **83**, 2133 (1999).
- [7] UA1 Collaboration, C. Albajar *et al.*, Phys. Lett. B **198**, 261 (1987); UA2 Collaboration, J. Alitti *et al.*, Phys. Lett. B **235**, 363 (1990).
- [8] L3 Collaboration, M. Acciarri *et al.*, Phys. Lett. B **471**, 308 (1999).
- [9] ALEPH Collaboration, R. Barate *et al.*, Phys. Lett. B **499**, 67 (2001).
- [10] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **69**, 3439 (1992); Phys. Rev. D **56**, 1357 (1997).
- [11] D0 Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **75**, 618 (1995).
- [12] M. Spiropulu, Ph.D. thesis, Harvard University, 2000.
- [13] CDF Collaboration, F. Abe *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **271**, 387 (1988); Phys. Rev. D **50**, 2966 (1994).
- [14] For details on the lepton, jet, and  $\cancel{E}_T$  identification, see J. Done, Ph.D. thesis, Texas A&M University, 1999.
- [15] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **76**, 4307 (1996); **80**, 5275 (1998).
- [16] H. Baer, F. E. Paige, S. D. Protopopescu, and X. Tata, hep-ph/9810440. We use ISAJET version 7.20.
- [17] CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **49**, 1 (1994); **59**, 052002 (1999).
- [18] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **80**, 2773 (1998).
- [19] CDF Collaboration, F. Abe *et al.*, Phys. Rev. D **55**, 2546 (1997).
- [20] J. Ohnemus and J. Owens, Phys. Rev. D **43**, 3626 (1991); J. Ohnemus, *ibid.* **44**, 1403 (1991); **44**, 3477 (1991).
- [21] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **80**, 2779 (1998).
- [22] See, for example, K. Hikasa and M. Kobayashi, Phys. Rev. D **36**, 724 (1987); H. Baer *et al.*, Phys. Rev. D **44**, 725 (1991); H. Baer, J. Sender, and X. Tata, Phys. Rev. D **50**, 4517 (1994).
- [23] A. H. Chamseddine, R. Arnowitt, and P. Nath, Phys. Rev. Lett. **49**, 970 (1982); **50**, 232 (1983); R. Barbieri, S. Ferrara, and C. A. Savoy, Phys. Lett. **119B**, 343 (1982); L. Hall, J. Lykken, and S. Weinberg, Phys. Rev. D **27**, 2359 (1983); P. Nath, R. Arnowitt, and A. H. Chamseddine, Nucl. Phys. **B227**, 121 (1983).
- [24] H. Baer *et al.*, Phys. Rev. D **47**, 2739 (1992); M. Drees and M. Nojiri, Nucl. Phys. **B369**, 54 (1992).
- [25] L3 Collaboration, M. Acciarri *et al.*, Phys. Lett. B **350**, 109 (1995); OPAL Collaboration, G. Abbiendi *et al.*, Eur. Phys. J. C **8**, 255 (1999); DELPHI Collaboration, P. Abreu *et al.*, Report No. CERN-EP-2000-133 [Eur. Phys. J. C (to be published)].
- [26] CTEQ Collaboration, H. L. Lai *et al.*, Phys. Rev. D **51**, 4763 (1995).
- [27] W. Beenakker and R. Höpker, M. Spira, and P. M. Zerwas, Nucl. Phys. **B492**, 51 (1997).
- [28] A. D. Martin, R. G. Roberts, and W. J. Stirling, Phys. Lett. B **387**, 419 (1996).
- [29] M. Glück, E. Reya, and A. Vogt, Z. Phys. C **67**, 433 (1995).
- [30] G. Zech, Nucl. Instrum. Methods Phys. Res., Sect. A **277**, 608 (1989); T. Huber *et al.*, Phys. Rev. D **41**, 2709 (1990).