

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Agricultural Research Magazine

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

3-2013

Enhancing Yogurt With Healthful Fiber From Oats

Mukti Singh

USDA-ARS, mukti.singh@ars.usda.gov

Sanghoon Kim

USDA-ARS, sanghoon.kim@ars.usda.gov

Marcia Wood

USDA-ARS

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaagresmag>



Part of the [Agriculture Commons](#), [Animal Sciences Commons](#), [Food Science Commons](#), and the [Plant Sciences Commons](#)

Singh, Mukti; Kim, Sanghoon; and Wood, Marcia, "Enhancing Yogurt With Healthful Fiber From Oats" (2013). *Agricultural Research Magazine*. 79.

<https://digitalcommons.unl.edu/usdaagresmag/79>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agricultural Research Magazine by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Enhancing Yogurt With Healthful Fiber From Oats

PEGGY GREB (D1073-2)

Adding about one-quarter teaspoon of a fiber-rich component of oats boosts the nutritional value of low-fat yogurt without noticeably affecting the taste or texture of this increasingly popular dairy food.

Oat fiber is of interest to foodmakers and nutritionists alike. Studies with volunteers have shown that it can lower serum cholesterol, which may help improve heart health.

Agricultural Research Service food technologist Mukti Singh, chemist Sanghoon Kim, and their colleagues experimented with adding fiber-containing oat beta-glucan to what's known in the dairy industry as "low-fat yogurt mix." It is made up of low-fat milk and a selection of common, safe-to-eat bacteria that ferment the milk. Best known bacteria include *Lactobacillus acidophilus* or various *Bifidobacterium* species.

Singh's intent was to see how much fiber she could add without altering key qualities of yogurt, including texture, viscosity, color, pH, and fermentation time.

In experiments at the ARS National Center for Agricultural Utilization Research in Peoria, Illinois, Singh's team added 0.1, 0.2, 0.3, 0.4, or 0.5 percent purified oat beta-glucan—a light, fine-textured, whitish powder—to low-fat yogurt mix and compared the results to yogurt mix with no oat fiber.

The idea of adding edible fiber to yogurt isn't new. But in studies such as this, Singh is filling in some of the gaps in our knowledge of how fiber may affect yogurt-mix qualities that are important to foodmakers and to yogurt fans.

In the oat beta-glucan work, the team determined that up to 0.3 percent highly purified (95 percent pure) oat beta-glucan, which translates to 0.3 grams of beta-

glucan per 100 grams of yogurt mix, could be added without significantly altering key yogurt qualities. But adding 0.4 percent or higher changed the yogurt's color, contributed to unwanted hardening, and slowed fermentation.

The 0.3 percent level of fortification totals out at 0.75 grams of fiber per 8-ounce serving of yogurt. Most Americans don't get enough fiber, so even this small addition to a familiar dairy product helps.


A peer-reviewed article published in 2012 in the *Journal of Food Science* by Singh, Kim, and Sean X. Liu, research leader of the Functional Foods Research Unit at Peoria, documents the research.—
By **Marcia Wood**, ARS.

This research is part of Quality and Utilization of Agricultural Products, an ARS national program (#306) described at www.nps.ars.usda.gov.

*Mukti Singh is in the Functional Foods Research Unit and Sanghoon Kim is in the Plant Polymer Research Unit, USDA-ARS National Center for Agricultural Utilization Research, 1815 N. University St., Peoria, IL 61604; (309) 681-6357 [Singh], (309) 681-6260 [Kim], mukti.singh@ars.usda.gov, sanghoon.kim@ars.usda.gov.**



ARS scientists have added different levels of oat beta-glucan to low-fat yogurt mix to increase its fiber content.



STEPHEN AUSMUS (D2763-1)

At the ARS National Center for Agricultural Utilization Research in Peoria, Illinois, chemist Sanghoon Kim (left) and food technologist Mukti Singh review a microscope image showing the effects of oat beta-glucan during yogurt fermentation.