

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications: Department of  
Entomology

Entomology, Department of

---

1-16-2004

## Partial purification and characterization of a methyl-parathion resistance-associated general esterase in *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae)

Xuguo Zhou

University of Nebraska-Lincoln

Michael E. Scharf

Purdue University-West Lafayette, IN

Gautam Sarath

University of Nebraska-Lincoln, Gautam.sarath@ars.usda.gov

Lance J. Meinke

University of Nebraska-Lincoln, lmeinke1@unl.edu

Laurence D. Chandler

USDA-ARS

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

---

Zhou, Xuguo; Scharf, Michael E.; Sarath, Gautam; Meinke, Lance J.; Chandler, Laurence D.; and Siegfried, Blair, "Partial purification and characterization of a methyl-parathion resistance-associated general esterase in *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae)" (2004). *Faculty Publications: Department of Entomology*. 76.

<https://digitalcommons.unl.edu/entomologyfacpub/76>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Xuguo Zhou, Michael E. Scharf, Gautam Sarath, Lance J. Meinke, Laurence D. Chandler, and Blair Siegfried



# Partial purification and characterization of a methyl-parathion resistance-associated general esterase in *Diabrotica virgifera virgifera* (Coleoptera: Chrysomelidae)

Xuguo Zhou,<sup>a</sup> Michael E. Scharf,<sup>b</sup> Gautam Sarath,<sup>c</sup> Lance J. Meinke,<sup>a</sup> Laurence D. Chandler,<sup>d</sup> and Blair D. Siegfried<sup>a,\*</sup>

<sup>a</sup> Department of Entomology, University of Nebraska-Lincoln, Lincoln, NE 68583, USA

<sup>b</sup> Department of Entomology, Purdue University-West Lafayette, IN 47907, USA

<sup>c</sup> Department of Biochemistry, University of Nebraska-Lincoln, Lincoln, NE 68583, USA

<sup>d</sup> USDA-ARS Red River Valley Experiment Station, Fargo, ND 58105, USA

Received 4 August 2003; accepted 27 October 2003

## Abstract

Increased hydrolytic metabolism of organophosphate insecticides has been associated with resistance among Nebraska western corn rootworm populations. In this study, resistance-associated esterases were partially purified by differential centrifugation, ion exchange, and hydroxyapatite column chromatography, with a final purification factor of 100-fold and recovery of approximately 10%. Kinetic analysis of the partially purified enzyme indicated that the  $K_m$  of the group II esterases was identical for the two populations, although  $V_{max}$  was consistently threefold higher in the resistant population. A putative esterase, DvvII, was further purified to homogeneity by preparative polyacrylamide gel electrophoresis. DvvII is a monomer with a molecular weight of approximately 66 kDa, although three distinct isoforms with similar pIs were evident based on isoelectric focusing gel electrophoresis. Immunoassays with the *Myzus persicae* E4 antiserum indicated that group II esterases from *D. v. virgifera* were cross-reactive and expressed at much higher titers in the resistant population relative to the susceptible counterpart. These results suggest that the resistance is likely associated with overproduction of an esterase isozyme in resistant *D. v. virgifera* populations.

© 2003 Elsevier Inc. All rights reserved.

**Keywords:** Esterase; Insecticide resistance; Purification; Characterization; *Diabrotica virgifera virgifera*

## 1. Introduction

The western corn rootworm, *Diabrotica virgifera virgifera* LeConte, is a serious pest of field

corn, *Zea mays* (L.) in the United States [1,2]. Crop rotation and chemical control have been the primary management strategies [2], although *D. v. virgifera* has become increasingly difficult to control because of its remarkable ability to adapt to these management practices. Chlorinated hydrocarbons were the first synthetic

\* Corresponding author. Fax: 1-402-472-4687.

E-mail address: [bsiegfried1@unl.edu](mailto:bsiegfried1@unl.edu) (B.D. Siegfried).

insecticide class to be introduced in Nebraska for larval control in the late 1940s [3]. However, as a consequence of their broadcast application and extreme persistence, resistance to cyclodienes was detected in less than a decade [3,4].

In certain areas of the Platte Valley of south central Nebraska, adult rootworm control was adopted after the development of cyclodiene resistance [5]. Microencapsulated methyl-parathion (PennCap-M, Elf Atochem North America, Philadelphia, PA) has been the most commonly used organophosphate insecticide to control adult *D. v. virgifera* in Nebraska because of its low cost and relatively long persistence. PennCap failures were first reported in the 1990s [5], and preliminary studies implied the involvement of hydrolytic metabolism in resistant rootworm populations [6]. In vivo metabolism experiments with <sup>14</sup>C-labeled ethyl-parathion, synergism bioassays with the esterase inhibitor DEF<sup>1</sup> (*S,S,S*-tributyl phosphorotriothioate), and esterase activity assays with model substrates suggested a common esterase-mediated resistance in both larvae and adult *D. v. virgifera* [6,8–11].

Recent investigations have identified three distinct esterase isozyme groups (I, II, and III) from *D. v. virgifera* based on their mobility in native polyacrylamide gel electrophoresis (PAGE) [11]. Elevated activity of group II esterase isozymes is highly correlated with resistance and the activity of these enzymes is strongly inhibited by a variety of organophosphate and carbamate insecticides [7, unpublished]. These enzymes have provided a reliable biochemical marker for detection of methyl-parathion resistance in individual *D. v. virgifera* adults [10]. In this investigation, we report the purification and characterization of the methyl-parathion resistance-associated group II esterases and provide a basis for defining a molecular mechanism of resistance.

## 2. Materials and methods

### 2.1. *D. v. virgifera* populations

Two populations were collected in Nebraska during the summer of 1998 and 1999. The Saunders County population was collected from areas where adult management is not practiced and has previously been shown to be highly susceptible to methyl-parathion. The resistant Phelps County population originated from an area where continuous corn production and adult control practice have been used for more than two decades and where field control failures have been reported. Topical bioassays with methyl-parathion indicated a 16.4-fold level of resistance in the Phelps County population relative to the Saunders County population [5]. In addition, the Saunders and Phelps County populations exhibited 97.0 and 7.0% mortality, respectively, after exposure to a diagnostic concentration of methyl-parathion [10]. Field populations were collected before seasonal application of adulticides had been initiated. The field populations were maintained in the laboratory at ambient temperature on a 14:10 h (L:D) photoperiod and were fed on a standard diet of head lettuce and sweet corn before being harvested and stored at –80 °C.

### 2.2. Chemicals

DEAE-Sepharose, hydroxyapatite, and all other reagents and solvents were purchased from Sigma Chemical (St. Louis, MO) unless stated otherwise. Electrophoresis reagents and ampholytes were purchased from Bio-Rad Laboratories (Hercules, CA). Protein assay reagents and bovine serum albumin were purchased from Pierce (Rockford, IL).

### 2.3. Protein and esterase assays

Protein concentration was determined with a commercially available bicinchoninic acid protein assay kit (Pierce, Rockford, IL) using bovine serum albumin as the standard. During purification, esterase activity was estimated by colorimetric measurement of  $\alpha$ -naphthol formation from

<sup>1</sup> Abbreviations used: DEF, *S,S,S*-tributyl phosphorotriothioate; PAGE, polyacrylamide gel electrophoresis; SDS, sodium dodecyl sulfate; IEF, isoelectric focusing; PNPA, *p*-nitrophenyl acetate; Mops, 3-(*N*-morpholino) propanesulfonic acid; PVDF, polyvinylidene difluoride.

$\alpha$ -naphthyl acetate. The approach of Van Asperen [12] was adapted to a 96-well microplate format. The assay mixture contained 188  $\mu$ l of 0.02 M sodium phosphate buffer (pH 7.0), 2  $\mu$ l 25 mM  $\alpha$ -naphthyl acetate (in acetone), and 10  $\mu$ l enzyme preparation. The reaction was stopped by the addition of 33.2  $\mu$ l of fast blue BN (tetrazotized *O*-dianisidine) (0.3% in 3.5% sodium lauryl sulfate). The plate was incubated at 30 °C for 15 min, the absorbance was recorded at 600 nm using a Bio Kinetics Reader (BIO-TEK, Winooski, VT). Formation of product was determined from a standard curve of  $\alpha$ -naphthol.

#### 2.4. Electrophoresis

Continuous non-denaturing PAGE was performed in a vertical Mini Protean II electrophoresis apparatus (Bio-Rad, Richmond, CA) using a 7.5% separating gel and 4% stacking gel with a continuous Tris/glycine running buffer system (50 mM, pH 8.3). Samples were diluted 1:1 with 2 $\times$  sample buffer [20% sucrose (w/v), 0.1% bromophenol blue in 50 mM Tris/glycine running buffer (pH 8.3)]. Gels were run at 120 V constant voltage for 75 min at 4 °C. Esterase bands were then visualized by incubating the gels in 100 ml of 0.02 M sodium phosphate buffer (pH 7.0), with 2% (v/v) 30 mM  $\alpha$ -naphthyl acetate dissolved in acetone at 30 °C for 15 min before adding 0.04% (w/v) fast blue BN salt (tetrazotized *O*-dianisidine).

Denaturing SDS (sodium dodecyl sulfate)–PAGE was used to examine protein composition of crude homogenates and column fractions during the purification process. SDS–PAGE was performed along with molecular size standard proteins (Kaleidoscope Prestained Standards;  $M_r$ , 6700–205,000; Bio-Rad, Hercules, CA) using a Bio-Rad Mini-Protean II electrophoresis unit and a 10% acrylamide concentration as described by Laemmli [13]. Coomassie brilliant blue G was used for staining protein on SDS–PAGE gels. The molecular weight of purified esterase was determined using a calibration curve of the protein standards.

Isoelectric focusing electrophoresis (IEF) was performed on a vertical unit (Model MGV-100, Scientific, Del Mar, CA) using a 7.5% polyacryl-

amide gel with narrow pH range ampholytes (5–7). Gels were run at ambient temperature and at 200 V for the first 2 h followed by 400 V for an additional 2 h. Esterase bands were visualized by the method previously described for non-denaturing PAGE. In order to determine the pH values at various points across the gel, an empty lane (50 mm in width) adjacent to the sample was excised and used as a pH ladder. Ten evenly spaced segments (2 mm  $\times$  50 mm) from the pH ladder were excised and placed in 0.5 ml double distilled water for at least 30 min. The pH of the ampholytes eluted from each segment was measured by an AR15 pH meter (Fisher Scientific, Bohemia, NY), which was standardized with a buffer of known pH (Fisher Certified Buffer). Sample lanes were stained with esterase activity as described previously, and the *pI* of each isozyme band was determined based on a calibration curve of pH versus distance from the cathode [14].

#### 2.5. Enzyme purification

Unless stated otherwise, all purification procedures were performed at 4 °C. An initial purification involving differential centrifugation was employed since the group II esterase isozymes are primarily cytosolic in nature [11]. Abdomens of 20 rootworm beetles (stored at –80 °C) were homogenized in 1.5 ml ice-cold 0.1 M sodium phosphate homogenization buffer (pH 7.8) with a Potter–Elvehjem homogenizer. The homogenate was centrifuged in an Eppendorf microcentrifuge (Brinkman Instruments, Westbury, NY) at 10,000g for 20 min. The 10,000g supernatant was filtered through glass wool to remove lipids and centrifuged in a Beckman Optima TLX-1 M-1 ultracentrifuge (Allendale, NJ) at 100,000g for 1 h. The resulting supernatant was applied to a DEAE-Sepharose ion exchange column (Pharmacia 1.6 cm ID  $\times$  40 cm length) equilibrated with 30 mM MOPS [3-(*N*-morpholino) propanesulfonic acid] buffer (pH 8.0). Esterase activity was eluted with a salt gradient (0–0.5 M NaCl in MOPS buffer) at 1 ml/min for 100 min. Fractions of 2 ml were collected and elution profiles determined for esterase activity and protein. Peak fractions were assayed by non-denaturing PAGE, and the

fractions containing group II esterase isozymes were combined and concentrated in an Amicon 8050 stirred cell (Beverly, MA) with a 10,000 kDa ultrafiltration membrane. The concentrated fraction was then desalted with 30 mM Mops buffer (pH 8.0) in the same stirred cell.

The desalted and concentrated ion exchange fraction was applied to a hydroxyapatite column (Pharmacia 1 cm ID  $\times$  12 cm length) equilibrated with 10 mM sodium phosphate buffer (pH 6.8). Group II esterase isozymes were eluted with a phosphate gradient (10–500 mM, pH 6.8) at 1 ml/min for 70 min and fractions of 1 ml were collected. Elution profiles were determined for esterase activity and protein. Hydroxyapatite chromatography was performed at ambient temperature. Esterase activity was eluted in one major peak, and peak fractions were examined by non-denaturing PAGE. The fractions containing group II esterase activity were combined, concentrated, and exchanged with 30 mM MOPS buffer (pH 8.0) as previously described.

The concentrated hydroxyapatite fraction was further concentrated to less than 0.1 ml with NANOSEP microconcentrators (10,000 kDa cutoff, Pall, Ann Arbor, MI), and subjected to preparative electrophoresis. The concentrated sample was examined by SDS-PAGE and protein bands were visualized by zinc staining (Bio-Rad, Hercules, CA). The bands of interests were excised, destained, and then eluted overnight with SDS running buffer in a Bio-Rad Model 422 Electro-Eluter (Bio-Rad, Hercules, CA). Eluted samples were dialyzed for 24 h against three changes of 5 mM Tris buffer (2 L) in individual Slide-A-Lyzer Dialysis Cassettes (10,000 kDa cutoff, Pierce, Rockford, IL). The dialyzed samples were concentrated in a Centrivap Concentrator (Labconco, Kansas City, MO) and the purity examined by SDS-PAGE.

## 2.6. Kinetic analysis

Michaelis–Menten constants ( $K_m$ ) and maximal velocities ( $V_{max}$ ) for the partially purified group II esterase (post-ion exchange fraction) from resistant and susceptible populations were determined by measuring activities toward a series of *p*-nitrophenyl acetate (PNPA) concentrations

(2.6–168.8 mM). The 210  $\mu$ l assay mixture consisted of 5  $\mu$ l PNPA in acetonitrile, 195  $\mu$ l of 0.02 M sodium phosphate buffer (pH 7.0), and 10  $\mu$ l enzyme preparation. The assay was initiated by the addition of PNPA, and the change in absorbance was recorded at 405 nm. An extinction coefficient of 6.53 mM<sup>-1</sup> cm<sup>-1</sup> was used to convert the absorbance in mOD/min to nanomoles of *p*-nitrophenol produced per minute.

## 2.7. Gel filtration

After ion exchange chromatography, the native molecular weight of the group II esterase was determined using a Superose 12 HR 10/30 column (LKB Pharmacia, Bromma, Sweden) equilibrated with 20 mM Mops buffer (pH 8.0). The column was calibrated at 4 °C with standard proteins ( $M_r$ , 12,400–160,000) (LKB Pharmacia, Bromma, Sweden). A 0.2 ml sample volume was applied with a flow rate of 0.4 ml/min and fractions of 0.4 ml were collected and elution profiles were determined for esterase activity and protein. The native molecular weight corresponding to the group II esterases was determined from a standard curve of the elution volume of standard proteins plotted against the logarithmic value of their molecular weight.

## 2.8. Western blotting

The polyclonal antiserum prepared from the E4 esterase of *Myzus persicae* (provided by A. Devonshire, IRAC Rothamsted, UK) was tested for cross-reactivity with the *D. v. virgifera* group II esterase by Western blotting. SDS-PAGE was performed in a vertical Mini Protean II electrophoresis apparatus using a 10% separating gel and 4% stacking gel. Individual wells were loaded with equal amounts of proteins and gels were run at constant 100 V for 90 min at ambient temperature. Each gel was then transferred to a polyvinylidene difluoride (PVDF) membrane (Millipore, Boston, MA) and probed with a polyclonal antiserum to *M. persicae* E4. Transfer was made using a Mini Trans-Blot Electrophoresis Transfer Cell (Bio-Rad, Hercules, CA) with 120 mM Tris, 40 mM glycine, and 20% methanol (v/v) as transfer buffer. Procedures for probing the blot with primary and

secondary antibody were according to instructions for the Western-Light and Western-Star System, a chemiluminescent immunoblot detection system with CDP-Star substrate (Applied Biosystems, Bedford, MA). The chemiluminescence was visualized with a Fluor-S Multiimager (Bio-Rad, Hercules, CA).

### 2.9. Amino acid sequence analysis

Approximately 100 pmol of the post-preparative electrophoresis fraction, partially purified group II esterases, were subjected to 10% SDS-PAGE and electroblotted onto PVDF membrane (Millipore, Bedford, MA). The blotted proteins were submitted to the W.M. Keck Foundation, Biotechnology Resource Laboratory, Yale University and analyzed by automated Edman degradation methods to determine NH<sub>2</sub>-terminal amino acid sequences of the putative group II esterase.

### 2.10. Statistical analyses

Standard curves for protein,  $\alpha$ -naphthol, molecular weight, and isoelectric focusing point were determined by least squares linear regression analysis.

## 3. Results

### 3.1. Purification of a group II esterase

Data from a representative purification of resistance-associated esterases are provided in Table 1. Ion exchange chromatography resulted in approximately 10-fold purification with 56.6% recovery. The ion exchange elution profiles obtained

from the resistant and susceptible populations (Fig. 1) indicated similar chromatographic behavior for protein and esterase activity in both strains although the activity peak corresponding to group II esterases was significantly larger from the resistant population. These results are consistent with denaturing PAGE results, in which peak fractions containing group II esterase activity from the resistant population were elevated relative to the susceptible population (Figs. 2A and B). Analysis of esterase isozyme composition by non-denaturing PAGE revealed that the three peak fractions corresponded to the three esterase groups previously identified. The group I esterases did not appear on the separating gel (Fig. 2A) because the majority of their activity remained in the stacking gel [10,11]. Hydroxyapatite column chromatography gave one major activity peak, which corresponded to the group II esterases based on non-denaturing PAGE (data not shown) and resulted in an additional 11-fold purification with 9.89% recovery.

Post-hydroxyapatite fractions containing group II esterases were subjected to preparative SDS-PAGE to isolate individual proteins. Examination of peak fractions from both ion-exchange and hydroxyapatite chromatography by SDS-PAGE revealed three protein bands of approximately 60, 25, and 20 kDa that exhibited increased staining intensity in the resistant strain (Fig. 3), in which a single protein band of expected size (ca. 60 kDa) was identified as the putative group II esterase based on molecular weight of other resistance associated esterases (Table 2) and cross-reactivity with the E4 esterase antiserum from *M. persicae* (see below). The 60 kDa band was purified to homogeneity following preparative electrophoresis, zinc staining, excision, destaining, and electroelution. The pure protein, referred to as DvII,

Table 1  
Purification of group II esterase from a resistant *D. v. virgifera* population

Purification step	Protein (mg)	Specific activity <sup>a</sup>	Units <sup>b</sup>	Recovery (%)	Purification factor
350,000g supernatant	28.29	397.82	11254.33	100.00	1.00
DEAE-Sepharose column	1.62	3946.13	6373.00	56.63	9.92
Hydroxyapatite column	0.08	42393.65	3338.50	9.89	106.56

<sup>a</sup> Specific activity is expressed as nmol/min/mg protein.

<sup>b</sup> Units are expressed as nmol/min.




Fig. 1. Elution profiles for protein and esterase activity using  $\alpha$ -naphthyl acetate as substrate and a DEAE-Sepharose ion exchange column. (A) Resistant population. (B) Susceptible population. See text for details of chromatography.

was concentrated for subsequent amino terminal sequencing. The purity and molecular weight of DvvII as determined by SDS-PAGE after the final purification procedure is presented in Fig. 3.

### 3.2. Physical properties of group II esterase

Isoelectric focusing of the partially purified group II esterases revealed three different esterase isozymes in the resistant population with isoelec-

tric points (pI) of  $5.32 \pm 0.04$ ,  $5.21 \pm 0.01$ , and  $5.07 \pm 0.02$ , respectively (Fig. 4). The monomeric molecular weight of the putative group II esterase, DvvII, estimated by SDS-PAGE was  $65.6 \pm 3.3$  kDa (mean  $\pm$  SE,  $n = 3$ ). The mass of the native form of this group II esterase estimated by gel filtration chromatography was 65.6 kDa ( $R^2 = 0.99$ , data not shown). These data in combination suggest that the native form of this esterase is monomeric.






Fig. 2. (A) Non-denaturing PAGE stained for esterase activity toward  $\alpha$ -naphthyl acetate of crude homogenates and three ion exchange peak fractions (Group I, II, and III) from both susceptible (S) and resistant (R) populations and (B) SDS-PAGE stained with Coomassie blue of peak ion exchange fractions (Group I, II, and III) from both susceptible (S) and resistant (R) populations and the resistant hydroxyapatite fraction (R<sub>HTP</sub>).




Fig. 3. Molecular weight of the partially purified group II esterase on SDS-PAGE. M<sub>r</sub>, molecular mass marker; E<sub>PURE</sub>, purified group II esterase; S<sub>C</sub> and R<sub>C</sub>, crude homogenates from the susceptible and resistant populations, respectively.




Fig. 4. Isoelectric point (pI) of the native group II esterase from a resistant *D. v. virgifera* population. Peak fractions from ion exchange chromatography and hydroxyapatite chromatography (HTP) used as enzyme source.

Table 2  
Physical properties of insect esterases from selected resistant species

Insect species/esterases	Order	$pI$	$M_r$ (kDa), native	$M_r$ (kDa), denatured	Molecular form	References
<i>Culex quinquefasciatus</i> A2	Diptera	5.2	67.0	67.0	Monomer	[31]
<i>Culex quinquefasciatus</i> A2	Diptera	NA*	58.0–67.0	58.0–67.0	Monomer	[32]
<i>Culex quinquefasciatus</i> B2	Diptera	5.0	60.0	60.0	Monomer	[33]
<i>Culex quinquefasciatus</i> B2	Diptera	5.0	NA	63.0	Monomer	[34]
<i>Culex quinquefasciatus</i> Est $\beta$ 1	Diptera	4.8	66.0	66.0	Monomer	[35]
<i>Culex tarsalis</i>						
Susceptible	Diptera	6.8	59.0	NA	NA	[36,37]
Resistance		6.8/6.2	59.0	NA	NA	
<i>Culex pipiens pallens</i>	Diptera	4.6	NA	66.0	NA	[38]
<i>Anopheles albimanus</i>	Diptera	4.6	60.0	NA	NA	[39]
<i>Lucilia cuprina</i>	Diptera	5.5	60.5	60.5	Monomer	[40,41]
<i>Myzus persicae</i>						
E4	Homoptera	NA	65.0	57.0/65.0	Monomer	[42,43]
FE4		NA	66.0	58.0/66.0	Monomer	
<i>Myzus persicae</i>						
E4	Homoptera	NA	65.0	58.8	Monomer	[44]
FE4		NA	66.0	60.2	Monomer	
<i>Myzus nicotianae</i>	Homoptera	4.6	67.3	66.0	Monomer	[45]
<i>Nilaparvata lugens</i> E <sub>1</sub> –E <sub>3</sub>	Homoptera	4.7–4.9	NA	62.0–64.0	Monomer	[46]
<i>Nilaparvata lugens</i>	Homoptera	4.7–5.0	NA	66.0–68.0	NA	[30]
<i>Laodelphax striatellus</i>	Homoptera	4.7–5.3	NA	66.0–70.0	NA	[47]
<i>Schizaphis graminum</i> Type II	Homoptera	4.8	102.0	52.0/56.0	Dimer	[26]
<i>Pseudophilus includens</i>	Lepidoptera	4.6–4.8	NA	80.0	NA	[48]
<i>Leptinotarsa decemlineata</i>	Coleoptera	4.8	46.0–48.0	46.0–48.0	Monomer	[49,50]
		4.5	57.0–59.0	30.0–33.0	Dimer	
<i>Leptinotarsa decemlineata</i>	Coleoptera	6.2	NA	NA	NA	[51]
<i>Tribolium castaneum</i>						
Resistant	Coleoptera	7.3	62.0	NA	Monomer	[52]
Susceptible		6.6	61.7	NA	Monomer	
<i>Oryzaephilus surinaemensis</i>	Coleoptera	NA	130.0	65.0	Dimer	[53]
<i>Oryzaephilus surinaemensis</i>	Coleoptera	NA	NA	71.0	Monomer	[54]
<i>Diabrotica virgifera virgifera</i>	Coleoptera	5.1–5.3	65.6	65.6	Monomer	This study
<i>Nephotettix cincticeps</i> E1–E4	Hemiptera	4.8–5.1	NA	E3: 58.6	NA	[55]
<i>Anisopteromalus calandrae</i>	Hymenoptera	5.2	NA	NA	NA	[56]
<i>Blattella germanica</i> E6	Dictyoptera	4.8	55.0	55.0	Monomer	[57]

\*NA, not available.

### 3.3. Kinetic analysis

Results of kinetic analysis of partially purified group II esterases from the resistant and susceptible population are presented in Fig. 5. Similar  $K_m$  values were obtained for the two populations although the  $V_{max}$  was consistently two- to threefold higher in the resistant population.  $K_m$  and  $V_{max}$  values for the resistant and susceptible population were  $8.40 \times 10^{-4}$  M and  $1.09 \mu\text{mol}/\text{min}/\text{ml}$ , and  $7.39 \times 10^{-4}$  M and  $0.304 \mu\text{mol}/\text{min}/\text{ml}$ , respectively. Similar  $K_m$  values indicate that enzymes

from susceptible and resistant populations interact similarly with the model substrate, *p*-nitrophenyl acetate. The higher  $V_{max}$  of the resistant strain suggests that there are greater amounts of the group II esterase in the resistant strain.

### 3.4. Cross-reactivity of *M. persicae* E4 antiserum with *D. v. virgifera* group II esterases

Fig. 6 shows that the *M. persicae* E4 antiserum was cross-reactive with a group II esterase isozyme weighing approximately 66 kDa in size from both




Fig. 5. Lineweaver–Burk plot of partially purified group II esterases from the resistant and susceptible populations.

resistant and susceptible *D. v. virgifera*. However, the enzyme preparation from the resistant population exhibited much stronger signal intensity than preparations from the susceptible counterpart. A crude homogenate prepared from the greenbug, *Schizaphis graminum*, was used as a positive control as it was previously shown that




Fig. 6. Cross-reaction of *D. v. virgifera* group II esterases with *M. persicae* E4 polyclonal antiserum. GB, type I Greenbug; S, *S. graminum* (a positive control for *M. persicae* E4 antiserum); R, resistant Phelps County population, NE; S, susceptible Saunders County population, NE; M, Kaleidoscope Prestained Standards (Bio-Rad, Hercules, CA).

greenbug esterase isozymes exhibit strong cross-reactivity with *M. persicae* E4 antiserum [15].

### 3.5. N-terminal amino acid analysis

The first 16 N-terminal amino acid residues of the putative group II esterase DvvII were identified as: Asp, Ile, Val, Leu, Asn, Lys, Lys, Arg, Phe, Pro, Lys, Asp, Phe, Met/Gly, Phe, and Gly. Comparisons of these 16 residues with N-terminal regions from other resistance-associated insect esterase revealed little or no homology with other known esterase sequences (Fig. 7).

Species	1			5			10			15						
Hva1	D	D	E	-	R	E	V	R	T	A	Q	G	P	L	R	G
CuB1	S	L	E	S	L	T	V	Q	T	K	Y	G	P	V	R	G
DmE6	T	D	D	P	L	L	V	Q	L	P	Q	G	K	L	R	G
MpE4/FE4	-	S	N	T	P	K	V	Q	V	H	S	G	E	I	A	G
BesB	-	-	E	S	P	R	V	T	V	K	H	G	T	L	-	G
N1-Est1	N	S	V	P	V	V	H	D	T	A	S	G	D	L	S	G
GbT2Est (56)	-	-	N	P	-	V	V	R	I	T	N	G	A	I	R	G
HH012f/MH130f	E	N	D	F	P	F	V	S	T	K	L	G	D	I	K	-
7012/malRR	-	N	D	F	P	F	V	S	T	K	L	-	G	I	GA	-
DvvII	D	I	V	L	N	K	K	R	F	P	K	D	F	M/G	F	G

Fig. 7. Comparison of the first 16 N-terminal sequences of the *D. v. virgifera* group II esterase DvvII with other amino terminal ends from: *Heliothis virescens* (Hva1), *Culex* spp. esterase B1 (CuB1), *Drosophila melanogaster* esterase-6 (DmE6), *Myzus persicae* E4 and FE4 (MpE4/FE4), *Bombyx mori* blood esterase B (BesB), *Nilaparvata lugens* esterase (N1-Est1), 56kDa subunit of Type II *Schizaphis graminum* esterase [GbT2Est (56)], and esterases from two different organophosphate-resistant *Oryzaephilus surinamensis* populations (HH012f/MH130f, and 7012/malRR). The sequences were aligned by hand using results of PILEUP analysis (Wisconsin Package Version 10.2, Genetics Computer Group, Madison, WI). Letters in bold denote amino acid residues conserved among the 10 sequences.

#### 4. Discussion

A putative *D. v. virgifera* group II esterase, DvvII, was partially purified by the combined techniques of differential centrifugation, ion exchange chromatography, hydroxyapatite chromatography, and preparative SDS–PAGE. The molecular form of DvvII is consistent with most resistance-associated esterases from other insect species (Table 2). The isoelectric point as well as the molecular weight estimated by both SDS–PAGE and gel filtration lie within the range of pH 4–7 and 55–70 kDa in size, respectively, which are typical of other insect esterases (Table 2) [16–18]. Although DvvII has similar physical properties with many other resistance-associated esterases (Table 2), it exhibits little homology at the N-terminal sequence. However, it should be noted that an esterase isolated from another organophosphate-resistant Coleopteran, *Oryzaephilus surinamensis*, also exhibits little homology with N-terminal sequences from other insect esterases [19].

The combined results of this investigation indicate that the differences in group II esterase activity between resistant and susceptible *D. v. virgifera* populations are the result of quantitative differences in the amount of enzymes [20–25]. Following purification, a 66 kDa band was clearly visible after Coomassie staining in the resistant population, but barely visible in the susceptible population. The difference in intensity between populations indicates a quantitative mechanism of resistance involving over-production of the same enzyme that is present in susceptible insects. This conclusion is supported by results of western blotting using the E4 antiserum of *M. persicae* where increased signal intensity of the 66 kDa band was observed in the resistant strain. The cross-reactivity exhibited between the 66 kDa band from both resistant and susceptible *D. v. virgifera* with *M. persicae* E4 antiserum suggests that the resistance associated esterase is not immunologically distinct from the susceptible esterase and the intensity difference is likely the result of the quantitative rather than qualitative modification of group II esterases [26–29].

If resistance were the result of a mutant ali-esterase with increased catalytic efficiency, qualita-

tive differences in physical and chemical properties of the enzyme between the resistant and susceptible population should have been evident. Similarities between the resistant and susceptible population in chromatographic behavior, physical and immunological properties of group II esterases suggest the existence of identical esterases in both populations. However, it is still not clear whether the genetic basis of this esterase-mediated resistance is the result of multiple gene copies (i.e., gene amplification), or if the overproduction is caused by a change in the sequences of regulatory DNA outside the protein coding region that leads to increased protein synthesis.

#### Acknowledgments

The authors thank Z.B. Mayo and David Taylor for valuable comments on an earlier draft of the manuscript. Jim Brown and Jenny Stebbing helped collect and maintain *D. v. virgifera* field populations. Support for this project was provided by DuPont, FMC, Zeneca, American Cyanamid, Dow Agrosciences, USDA Specific Cooperative Agreement 58-5447-6-116, USDA Initiative for Future Agricultural and Food Science Grant No. 00-52103-9699) and NIH Grant No. 1P20RR16569 from the BRIN Program of the National Center for Research Resources.

#### References

- [1] H.C. Chiang, Bionomics of the northern and western corn rootworms, *Annu. Rev. Entomol.* 18 (1973) 47–72.
- [2] E. Levine, S.H. Oloumi, Management of *Diabrotica* rootworms in corn, *Annu. Rev. Entomol.* 36 (1991) 229–255.
- [3] H.J. Ball, G.T. Weekman, Insecticide resistance in the western corn rootworm in Nebraska, *J. Econ. Entomol.* 55 (1963) 439–441.
- [4] R.L. Metcalf, Foreword, in: J.L. Krysan, T.A. Smith (Eds.), *Methods for the Study of Pest Diabrotica*, Springer-Verlag, New York, 1986.
- [5] L.J. Meinke, B.D. Siegfried, R.J. Wright, L.D. Chandler, Adult susceptibility of Nebraska western corn rootworm (Coleoptera: Chrysomelidae) populations to selected insecticides, *J. Econ. Entomol.* 91 (1998) 594–600.
- [6] F. Miota, M.E. Scharf, M. Ono, P. Marcon, L.J. Meinke, R.J. Wright, L.D. Chandler, B.D. Siegfried, Mechanisms

- of methyl and ethyl parathion resistance in the western corn rootworm (Coleoptera: Chrysomelidae), *Pestic. Biochem. Physiol.* 61 (1998) 39–52.
- [7] M.E. Scharf, L.J. Meinke, R.J. Wright, L.D. Chandler, B.D. Siegfried, Metabolism of carbaryl by insecticide-resistant and -susceptible western corn rootworm populations (Coleoptera: Chrysomelidae), *Pestic. Biochem. Physiol.* 63 (1999) 85–96.
- [8] M.E. Scharf, L.J. Meinke, B.D. Siegfried, R.J. Wright, L.D. Chandler, Carbaryl susceptibility, diagnostic concentration determination, and synergism for U.S. populations of western corn rootworm (Coleoptera: Chrysomelidae), *J. Econ. Entomol.* 92 (1999) 33–39.
- [9] R.J. Wright, M.E. Scharf, L.J. Meinke, X. Zhou, B.D. Siegfried, L.D. Chandler, X.G. Zhou, Larval susceptibility of an insecticide-resistant western corn rootworm (Coleoptera: Chrysomelidae) population to soil insecticides: laboratory bioassays, assays of detoxification enzymes, and field performance, *J. Econ. Entomol.* 93 (2000) 7–13.
- [10] X.G. Zhou, M.E. Scharf, P. Srinivas, L.J. Meinke, R.J. Wright, L.D. Chandler, B.D. Siegfried, Diagnostic assays based on esterase-mediated resistance mechanisms in western corn rootworms (Coleoptera: Chrysomelidae), *J. Econ. Entomol.* 95 (2002) 1261–1266.
- [11] X.G. Zhou, M.E. Scharf, L.J. Meinke, L.D. Chandler, B.D. Siegfried, Characterization of general esterases from methyl-parathion resistant and susceptible populations of western corn rootworm (Coleoptera: Chrysomelidae), *J. Econ. Entomol.* (2003) (in press).
- [12] Van Asperen, A study of housefly esterases by means of a sensitive colorimetric method, *J. Insect Physiol.* 8 (1962) 401–416.
- [13] U.K. Laemmli, Cleavage of structural proteins during the assembly of the head of bacteriophage T4, *Nature* 227 (1970) 680–685.
- [14] C.M. Conyers, A.D. MacNicoll, N.R. Price, Purification and characterization of an esterase involved in resistance to organophosphorous insecticides in the saw-toothed grain beetle, *Oryzaephilus surinamensis* (Coleoptera: Silvanidae), *Insect Biochem. Mol. Biol.* 28 (1998) 435–448.
- [15] B.D. Siegfried, J.J. Swanson, A.L. Devonshire, Immunological detection of greenbug (*Schizaphis graminum*) esterases associated with resistance to organophosphate insecticides, *Pestic. Biochem. Physiol.* 57 (1997) 165–170.
- [16] G.J. Small, J. Hemingway, Differential glycosylation produces heterogeneity in elevated esterases associated with insecticide resistance in the brown planthopper *Nilaparvata lugens* Stal, *Insect Biochem. Mol. Biol.* 30 (2000) 443–453.
- [17] E. Heymann, Carboxylesterases and amidases, in: W.B. Jakoby (Ed.), *Enzymatic Basis of Detoxication*, Academic Press, New York, 1980, p. 291.
- [18] M. Cygler, S. Wu, A. Zdanov, D.R. Bundle, D.R. Rose, Recognition of a carbohydrate antigenic determinant of *Salmonella* by an antibody, *Biochem. Soc. Trans.* 21 (1993) 441.
- [19] L.C. Rossiter, R.V. Gunning, H.A. Rose, The use of polyacrylamide gel electrophoresis for the investigation and detection of fenitrothion and chlorpyrifos-methyl resistance in *Oryzaephilus surinamensis* (Coleoptera: Silvanidae), *Pestic. Biochem. Physiol.* 69 (2001) 27–34.
- [20] J. Hemingway, S.H.P.P. Karunaratne, Mosquito carboxylesterases: a review of the molecular biology and biochemistry of a major insecticide resistance mechanism, *Med. Vet. Entomol.* 12 (1998) 1–12.
- [21] S.H.P.P. Karunaratne, K.G.I. Jayawardena, J. Hemingway, A.J. Ketterman, Characterization of a B-type esterase involved in insecticide resistance from the mosquito *Culex quinquefasciatus*, *Biochem. J.* 294 (1993) 575–579.
- [22] J. Hemingway, The molecular basis of two contrasting metabolic mechanisms of insecticide resistance, *Insect Biochem. Mol. Biol.* 30 (2000) 1009–1015.
- [23] F.J. Oppenoorth, K. Van Asperen, Allelic genes in the housefly producing modified enzymes that cause organophosphate resistance, *Science* 132 (1960) 298.
- [24] A.L. Devonshire, L.M. Field, Gene amplification and insecticide resistance, *Annu. Rev. Entomol.* 36 (1991).
- [25] L.M. Field, A.L. Devonshire, B.G. Forde, Molecular evidence that insecticide resistance in peach-potato aphids *Myzus persicae* Sulz. results from amplification of an esterase gene, *Biochem. J.* 251 (1988) 309–312.
- [26] B.D. Siegfried, M. Ono, J.J. Swanson, Purification and characterization of a carboxylesterase associated with organophosphate resistance in the greenbug, *Schizaphis graminum* (Homoptera: Aphididae), *Arch. Insect Biochem. Physiol.* 36 (1997) 229–240.
- [27] M. Ono, J.J. Swanson, L.M. Field, A.L. Devonshire, B.D. Siegfried, Amplification and methylation of an esterase gene associated with insecticide-resistance in greenbugs, *Schizaphis graminum* (Rondani) (Homoptera: Aphididae), *Insect Biochem. Mol. Biol.* 29 (1999) 1065–1073.
- [28] C. Mouches, N. Pasteur, J.B. Berge, O. Hyrien, M. Raymond, B.R. Saint-Vincent, M. de Silvestri, G.P. Georghiou, Amplification of an esterase gene is responsible for insecticide resistance in a California *Culex* mosquito, *Science* 233 (1986) 778–780.
- [29] S.H.P.P. Karunaratne, A. Vaughan, M.G. Paton, J. Hemingway, Amplification of a serine esterase gene is involved in insecticide resistance in Sri Lankan *Culex tritaeniorhynchus*, *Insect Mol. Biol.* 7 (1998) 307–315.
- [30] G.J. Small, J. Hemingway, Molecular characterization of the amplified carboxylesterase gene associated with organophosphorus insecticide resistance in the brown planthopper, *Nilaparvata lugens*, *Insect Mol. Biol.* 9 (2000) 647–653.
- [31] A.J. Ketterman, K.G.I. Jayawardena, J. Hemingway, Purification and characterization of a carboxylesterase involved in insecticide resistance from the mosquito *Culex quinquefasciatus*, *Biochem. J.* 287 (1992) 355–360.
- [32] A. Vaughan, J. Hemingway, Mosquito carboxylesterase Est-alpha-2-1 (A-2): cloning and sequence of the full-length cDNA for a major insecticide resistance gene worldwide in the mosquito *Culex quinquefasciatus*, *J. Biol. Chem.* 270 (1995) 17044–17049.

- [33] A.J. Ketterman, S.H.P.P. Karunaratne, K.G.I. Jayawardena, J. Hemingway, Qualitative differences between populations of *Culex quinquefasciatus* in both the esterases A2 and B2 which are involved in insecticide resistance, *Pestic. Biochem. Physiol.* 47 (1993) 142–148.
- [34] K.G.I. Jayawardena, S.H.P.P. Karunaratne, A.J. Ketterman, J. Hemingway, Determination of the role of elevated B2 esterase in insecticide resistance in *Culex quinquefasciatus* (Diptera: Culicidae) from studies on the purified enzyme, *Bull. Entomol. Res.* 84 (1994) 39–44.
- [35] G.J. Small, S.H.P.P. Karunaratne, J. Hemingway, Characterization of amplified esterase Estbeta12 associated with organophosphate resistance in a multi-resistant population of the mosquito *Culex quinquefasciatus* from Cuba, *Med. Vet. Entomol.* 12 (1998) 187–191.
- [36] S. Whyard, A.E.R. Downe, V.K. Walker, Isolation of an esterase conferring insecticide resistance in the mosquito *Culex tarsalis*, *Insect Biochem. Mol. Biol.* 24 (1994) 819–827.
- [37] S. Whyard, A.E.R. Downe, V.K. Walker, Characterization of a novel esterase conferring insecticide resistance in the mosquito *Culex tarsalis*, *Arch. Insect Biochem. Physiol.* 29 (1995) 329–342.
- [38] J. Liang, J. Shen, G. Lin, W. Fu, J. Liang, J.H. Shen, G.F. Lin, W.J. Fu, Purification and biochemical property of resistance related carboxylesterase in mosquito *Culex pipiens pallens*, *Acta Entomol. Sin.* 44 (2001) 161–169.
- [39] W.G. Brogdon, A.M. Barber, Fenitrothion deltamethrin cross-resistance conferred by esterases in Guatemalan *Anopheles albimanus*, *Pestic. Biochem. Physiol.* 37 (1990) 130–139.
- [40] S. Whyard, R.J. Russell, V.K. Walker, Insecticide resistance and malathion carboxylesterase in the sheep blowfly, *Lucilia cuprina*, *Biochem. Genet.* 32 (1994) 9–24.
- [41] S. Whyard, V.K. Walker, Characterization of malathion carboxylesterase in the sheep blowfly *Lucilia cuprina*, *Pestic. Biochem. Physiol.* 50 (1994) 198–206.
- [42] A.L. Devonshire, G.D. Moores, A carboxylesterase with broad substrate specificity causes organophosphorus, carbamate and pyrethroid resistance in peach-potato aphids (*Myzus persicae*), *Pestic. Biochem. Physiol.* 18 (1982) 235–246.
- [43] A.L. Devonshire, L.M. Searle, G.D. Moores, Quantitative and qualitative variation in the mRNA for carboxylesterases in insecticide-susceptible and resistant *Myzus persicae* (Sulz), *Insect Biochem.* 16 (1986) 659–665.
- [44] L.M. Field, M.S. Williamson, G.D. Moores, A.L. Devonshire, Cloning and analysis of the esterase genes conferring insecticide resistance in the potato-peach aphid, *Myzus persicae*, *Biochem. J.* 294 (1993) 569–574.
- [45] M.A. Wolff, Y.A.I. Abdel-Aal, D.K.S. Goh, E.P. Lampert, R.M. Roe, Organophosphate resistance in the tobacco aphid (Homoptera: Aphididae): purification and characterization of a resistance-associated esterase, *J. Econ. Entomol.* 87 (1994) 1157–1164.
- [46] W. Chen, C. Sun, Purification and characterization of carboxylesterases of a rice brown planthopper, *Nilaparvata lugens* Stal, *Insect Biochem. Mol. Biol.* 24 (1994) 347–355.
- [47] K. Sakata, T. Miyata, Biochemical characterization of carboxylesterase in the small brown planthopper *Laodelphax striatellus* (Fallen), *Pestic. Biochem. Physiol.* 50 (1994) 247–256.
- [48] P.F. Dowd, T.C. Sparks, Characterization of a transpermethrin hydrolyzing enzyme from the midgut of *Pseudophusia includens* (Walker), *Pestic. Biochem. Physiol.* 25 (1986) 73–81.
- [49] S.H. Lee, J.M. Clark, Purification and characterization of multiple-charged forms of permethrin carboxylesterase(s) from the hemolymph of resistant Colorado potato beetle, *Pestic. Biochem. Physiol.* 60 (1998) 31–47.
- [50] S.H. Lee, J.M. Clark, Antibody capture immunoassay for the detection of permethrin carboxylesterase in Colorado potato beetle, *Leptinotarsa decemlineata* Say, *Pestic. Biochem. Physiol.* 64 (1999) 66–75.
- [51] D.D. Anspaugh, G.G. Kennedy, R.M. Roe, Purification and characterization of a resistance-associated esterase from the Colorado potato beetle, *Leptinotarsa decemlineata* (Say), *Pestic. Biochem. Physiol.* 53 (1995) 84–96.
- [52] E. Haubruge, M. Amichot, A. Cuany, J.B. Berge, L. Arnaud, Purification and characterization of a carboxylesterase involved in malathion-specific resistance from *Tribolium castaneum* (Coleoptera: Tenebrionidae), *Insect Biochem. Mol. Biol.* 32 (2002) 1181–1190.
- [53] S. Lee, E.M. Lees, B.C. Campbell, S.E. Lee, Purification and characterization of an esterase conferring resistance to fenitrothion in *Oryzaephilus surinamensis* (L.) (Insecta, Coleoptera, Silvanidae), *J. Agric. Food Chem.* 48 (2000) 4991–4996.
- [54] L.C. Rossiter, C.M. Conyers, A.D. MacNicoll, H.A. Rose, Two qualitatively different B-esterases from two organophosphate-resistant strains of *Oryzaephilus surinamensis* (Coleoptera: Silvanidae) and their roles in fenitrothion and chlorpyrifos-methyl resistance, *Pestic. Biochem. Physiol.* 69 (2001) 118–130.
- [55] S.W. Chiang, C.N. Sun, Purification and characterization of carboxylesterases of a rice green Leafhopper *Nephotettix cincticeps* Uhler, *Pestic. Biochem. Physiol.* 54 (1996) 181–189.
- [56] J.E. Baker, J.A. Fabrick, K.Y. Zhu, Characterization of esterases in malathion-resistant and susceptible strains of the pteromalid parasitoid *Anisopteromalus calandrae*, *Insect Biochem. Mol. Biol.* 28 (1998) 1039–1050.
- [57] S.K. Prabhakaran, S.T. Kamble, Purification and characterization of an esterase isozyme from insecticide resistant and susceptible strains of German cockroach, *Blattella germanica* (L.), *Insect Biochem. Mol. Biol.* 25 (1995) 519–524.