

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Bulletin of the University of Nebraska State
Museum

Museum, University of Nebraska State

1969

A New Rhinoceros from the Nebraska Miocene

Lloyd G. Tanner

Follow this and additional works at: <https://digitalcommons.unl.edu/museumbulletin>



Part of the [Entomology Commons](#), [Geology Commons](#), [Geomorphology Commons](#), [Other Ecology and Evolutionary Biology Commons](#), [Paleobiology Commons](#), [Paleontology Commons](#), and the [Sedimentology Commons](#)

Tanner, Lloyd G., "A New Rhinoceros from the Nebraska Miocene" (1969). *Bulletin of the University of Nebraska State Museum*. 52.

<https://digitalcommons.unl.edu/museumbulletin/52>

This Article is brought to you for free and open access by the Museum, University of Nebraska State at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Bulletin of the University of Nebraska State Museum by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

BULLETIN OF
The University of Nebraska State Museum VOLUME 8, NUMBER 6
APRIL 1969

Lloyd G. Tanner

A New Rhinoceros from
the Nebraska Miocene





The University of Nebraska

The Board of Regents

RICHARD E. ADKINS
J. G. ELLIOTT
B. N. GREENBERG, M.D.
RICHARD L. HERMAN
ROBERT L. RAUN
EDWARD SCHWARTZKOPF
G. ROBERT ROSS, *Corporation Secretary*

The Chancellor

MERK HOBSON, *Acting**

*Chancellor Clifford M. Hardin on leave as Secretary of Agriculture in the Cabinet of the President of the United States.

Lloyd G. Tanner

*A New Rbinoceros from
the Nebraska Miocene*

A B S T R A C T

A New Rhinoceros from the Nebraska Miocene

LLOYD G. TANNER

A new species of rhinoceros, *Menoceras marslandensis*, is here proposed. It is based upon a nearly complete skull from the upper part of the Marsland Formation, Hemingford Group (Miocene), of Box Butte County, Nebraska. The new species was probably derived from *Menoceras arikareense* (Barbour), collected from the Agate Springs Quarries (Arikaree Group, Harrison Formation) of Sioux County, Nebraska, with which both *Diceratherium niobrarense* Peterson and *D. cooki* Peterson are likely synonymous. From *Menoceras arikareense*, the presumed ancestral stock, *Menoceras marslandensis* differs in possessing the following characters: (1) very long, fused but slightly cleft, nasals, with flattened, rugose areas at the tips for the support of the horns; and (2) a convex frontal, also roughened, indicating a base for a well-developed frontal horn. This new species provides additional evidence that the generic separation of *Menoceras* Troxell from *Diceratherium* Marsh is warranted, and these two genera evidently lived side-by-side in the Medial to Late Miocene of the Central Great Plains. Both stocks, each greatly modified, may have survived into the Pliocene.

A New Rhinoceros from the Nebraska Miocene

INTRODUCTION

Knowledge of the Tertiary rhinoceroses of North America has proceeded rather slowly during the last thirty-five years, but this has not been due to the lack of new materials. Rather, because of the wealth of fossil rhinoceros specimens found in this period (particularly by expeditions from the University of Nebraska State Museum and the Frick Laboratory, American Museum of Natural History), there has developed a need for a complete revision of the family. This has been accompanied by a realization of the magnitude of the task and the futility of additional descriptions of new forms without a better understanding of the phylogeny.

Although such a revision has not been accomplished, there has been some progress toward it, and a tentative phylogenetic scheme has resulted (Fig. 1). It is the purpose of this paper to present some of the evidence that has accumulated with respect to the distinction of two Miocene rhinoceros genera, *Diceratherium* and *Menoceras*, and to place on record a new species of the latter.

DICERATHERIUM *versus* MENOCERAS

These Miocene rhinoceroses are quite distinct (Troxell, 1921, pp. 206-207), but they have become confused in the literature since

¹ Associate Curator of Vertebrate Paleontology, University of Nebraska State Museum.

1921 and generally have been considered inseparable (Matthew, 1931, p. 8; Simpson, 1945, p. 142). As with any group of fossil mammals, understanding at the generic level must begin with the types of the typical species, fitted into a modern frame of reference that is fundamentally stratigraphic and composed of all available specimens.

Diceratherium has priority, having been established by Marsh at the same time that he described the type species, *D. armatum*, from the "Miocene beds, near the John Day River, in Eastern Oregon" (Marsh, 1875, pp. 242-243). He considered the specimens available for *D. armatum* indicated "an animal about two-thirds the size of the Indian Rhinoceros," but he included a general description of the type skull and feet, together with a table of measurements.

This type skull (Y.P.M. 10003,² Figs. 2, 3, and 4) was kindly made available by the staff members of the Yale Peabody Museum for this study. Additional information regarding its derivation was furnished from the records of the Yale Peabody Museum: "The specimen was collected by L. S. Davis, in early 1874. In a letter dated 'May 12, 1874,' he notified Marsh of the shipment of fossils 'all from the Cove,' among which was the specimen in question. 'The Cove' is *Turtle Cove*, from which many of Marsh's John Day specimens came. (Personal communication from James A. Hopson, June 22, 1964)."

The correlation of the John Day sediments of Oregon with the rocks of the Great Plains has not been clearly established, but it must include an interval about equivalent to the Harrison Formation of Nebraska and eastern Wyoming. After the revision of several groups of oreodonts, Schultz and Falkenbach (1949, p. 92) concluded that their oreodont "examples from the John Day indicated a geologic age approximating that of the Harrison of Nebraska and Wyoming." Other discussions of the geologic placement of the type and referred specimens of *Diceratherium armatum* are to be found in the papers of Osborn (1898), Troxell (1921), Wood (1933), Green (1958, p. 591), and Macdonald (1963, p. 296).

Menoceras was proposed by Troxell (1921, p. 206-207), with *D. cooki* Peterson (1906c, p. 282) as the type species; the latter is here considered to be a synonym of *D. arikareense* Barbour (1906a). Both types have been available for study. These are but a part of a large

²The following abbreviations are utilized to refer to institutions cited: A.M.N.H., American Museum of Natural History, New York City; C.M., Carnegie Museum, Pittsburgh; S.D.S.M., South Dakota School of Mines, Rapid City; U.N.S.M., University of Nebraska State Museum, Lincoln; U.S.N.M., United States National Museum, Washington, D. C.; and Y.P.M., Yale Peabody Museum, New Haven.

A New Rhinoceros from the Nebraska Miocene




FIG. 1—The two lineages of diceratheres in the Miocene (Arikaree and Hemingford) sediments of the Central Great Plains (High Plains) of Nebraska and adjacent states.

sample from the famous locality near Agate, Sioux County, Nebraska (Peterson, 1906a-c; Schultz, 1966; and Roberts, 1966), now authorized as the "Agate Fossil Beds National Monument." The University of Nebraska State Museum expeditions, under the leadership of Erwin H. Barbour, Captain James Cook, and Harold J. Cook, together with the Carnegie Museum party of O. A. Peterson, developed the fossil quarries here during the early 1900's. However, many other institutions and individuals worked at Agate during this same period of time (Schultz, 1966). The principal museums of the world display one or more specimens from these sites. The fossiliferous level of the three quarries (designated "A," "B," and "C" by Holland and Peterson, 1913, Fig. 1) is in the Harrison Formation, in the upper part of the Arikaree Group, and this horizon is usually correlated as medial Miocene (Schultz and Stout, 1961, pp. 6-8, 30, 49, Figs. 2-3, 21-22, also Fig. 10, *this paper*).

There are two composite mounted skeletons of *Menoceras arikarensis* in the collections of the University of Nebraska State Museum³ from the Agate Springs Fossil Quarries (U.N.S.M. 1147 and 1250), one of which is now on display at the University of Nebraska Trailside Museum at Fort Robinson, near Crawford, Nebraska

³ The "panel mount" figured by Barbour (1909, Pl. 1) may have been disassembled after the fire of March 6, 1912.

(Tanner, 1962). In addition, the Museum's study collection now contains five skulls, several mandibular rami, and much other skeletal material of this species. Of particular interest in the present study is the excellent skull of a mature male (U.N.S.M. 1251).

The characters of *Menoceras arikareense* shown by this mature skull (Figs. 5-6) are: (1) a saddle-shape when viewed from the side, with a slight convexity in the anterior part of the frontal; and (2) paired nasal bosses, as round and rugose knobs situated near the tips of the nasals, with the rugosity limited superiorly. The female of this species was probably nearly hornless, since only vestigial knobs are present on skulls with undistorted nasals.

By contrast, the male characters for the very much larger *Diceratherium armatum*, as shown by the type skull (Figs. 2-4), are: (1) only a slight concavity in the frontal region when viewed in profile; and (2) paired nasal bosses are elongate and elevated ridges, instead of near-spherical knobs, and situated far behind rather than over the nasal tips.

TABLE 1
CONTRASTED SKULL CHARACTERS FOR MALES OF *Menoceras* AND *Diceratherium*.

| <i>Menoceras</i> | <i>Diceratherium</i> |
|--|---|
| 1. Approximately one-third smaller than <i>Diceratherium</i> | Much larger than <i>Menoceras</i> |
| 2. In side view, saddle-shaped | In profile, only slightly concave, <i>not</i> saddle-shaped |
| 3. Frontal convex | Frontal flat |
| 4. Frontal rugose | Frontal probably smooth |
| 5. Narial notch retracted, above P ² | Narial notch not retracted, situated above P ¹ |
| 6. Nasal bosses subrounded knobs | Nasal bosses elongate, elliptical ridges |
| 7. Nasal bosses above and near tips of nasals | Nasal bosses posterior to tips of nasals |
| 8. Palate narrow and deep | Palate broad and shallow |
| 9. Zygomata expanded posteriorly and roughened | Zygomata smooth |
| 10. Complicated dental patterns | Relatively simple dental patterns |
| 11. Cristae present on M ³ , with only faint cingulae | Cristae absent on M ³ , with cingulae moderately developed |
| 12. Diameter of M ¹ -M ³ = 223 mm. | Diameter of M ¹ -M ³ = 254 mm. |

A New Rhinoceros from the Nebraska Miocene




FIG. 2—Lateral view of holotype male skull (Y.P.M. 10003) of *Diceratherium armatum* Marsh, from the John Day Miocene of Oregon: X $\frac{1}{4}$




FIG. 3—Dorsal view of the same skull shown as Fig. 2: X $\frac{1}{4}$




FIG. 4—Palatal view of the same skull shown as Figs. 2 and 3: X $\frac{1}{4}$



FIG. 5—Lateral view of male skull (U.N.S.M. 1251), referred to *Menoceras ari-kareense* (Barbour), from the Agate Springs Fossil Quarries (Harrison Formation), Sioux County, Nebraska: X $\frac{1}{3}$




FIG. 6—Dorsal view of the same skull shown as Fig. 5: X $\frac{1}{3}$

A New Rhinoceros from the Nebraska Miocene

The distinctions between the two male skulls, when considered together with the great size difference, are believed to be of generic rank, indicative of two distinct lineages (Fig. 1; Table 1).

THE *MENOCERAS* LINEAGE

Assuming that there are two lineages of North American diceratheres, the male skull of *Menoceras arikarensense* serves as a convenient point of reference for reconstructing the *Menoceras* lineage.

Moving upwardly in the Central Great Plains Miocene succession, the next higher horizon that has yielded remains of this genus is the base of the Marsland Formation, regarded also as the base of the Hemingford Group, and variously referred to the Medial or Late Miocene (Schultz and Stout, 1961, pp. 7-8, 51; Figs. 2-3, Fig. 10, *this paper*). This is the level of the Bridgeport Quarries of the University of Nebraska State Museum (U.N.S.M. Collecting Localities Mo-113, -114, -115, -116 and -118), in Morrill County, Nebraska, worked extensively by the expeditions of 1932-1935 and 1940. Two composite diceratheres skeletons from the Bridgeport Quarries, one with a male skull and the other a female skull, are on display in the University of Nebraska State Museum (U.N.S.M. 1238 and 1241), and there are additionally several thousand specimens in the study collections. The composite skeletons have been figured by Schultz (1943, p. 275) and later considered as a "geologic variety" of *Diceratherium niobrarense* Peterson (Stecher, Schultz, and Tanner, 1962). Some teeth, probably from these same sites, were suggested, by Wood (1964, pp. 378-380), to be very similar to *Diceratherium (Menoceras) barboursi* Wood, the holotype of which is from the well known "Thomas Farm Locality" in Florida.

In order to clarify the status of "*Diceratherium niobrarense*" [sic] it should be noted that the type skull of this species came from near Agate, Sioux County, Nebraska, at a site Peterson designated "Quarry A" (Peterson, 1920, p. 425). He states that "Quarry A": ". . . is only a short distance (300 yards) to the north of the main quarries and may possibly represent a somewhat earlier time; or, more probably, the sediments accumulated at this spot represent a different stream which had its origin in, and flowed through a locality more favorable to this species." (Peterson, 1920, p. 431).

This same site (Quarry A) is shown with other quarries on an unscaled "Rough outline topographical map" drawn by W. J. Holland and published in Holland and Peterson (1913, p. 191, Fig. 1). These authors, in their history of excavations at the Agate Springs Fossil Quarries, relate that "Quarry A" was the first to be excavated: "A few days after the arrival of Mr. Peterson he was conducted by

Mr. Harold Cook, the oldest son of his host, to the locality just beyond the eastern limits of Mr. Cook's property, designated as 'Quarry A' in the map (Fig. 1) accompanying this memoir." (Holland and Peterson, 1913, p. 189).

After Peterson worked a few days at this locality, he examined "two small buttes" situated about 300 yards south of "Quarry A." These buttes were subsequently named "University Hill" and "Carnegie Hill" by E. H. Barbour (1908) and are now considered the "main quarries" at the Agate Springs Fossil Quarries. Three pits were opened on Carnegie Hill (Quarry B with Pits 1, 2, and 3). At University Hill ("Quarry C") the excavation was assigned number 4. (Holland and Peterson, 1913, p. 191, Fig. 1).

Quarry A seems to be stratigraphically close in time to Quarries B and C.

The exact allocation of the specimens from the Bridgeport Quarries is still uncertain, despite the availability of the holotype of Peterson's *Diceratherium niobrarense* for the present study. The type skull, described by Peterson (1906c, pp. 466, 468, Fig. 2) from the Harrison Formation was examined in 1965, through the courtesy of the staff members of the Carnegie Museum, Pittsburgh. The illustrations of this young male skull (C.M. 1271) given by Peterson do not show the slight convexity on the frontal just behind the fronto-nasal suture, but a weak frontal convexity is present. This character is less pronounced than for specimens from the Marsland, which is to be expected, as the holotype came from the Harrison Formation.

A second partial skull in the Carnegie Museum collection from this same "Quarry A" (C.M. 1273) is of an older male, and it displays a more definite convexity on the frontal, together with slight depressions for the nutritive arteries radiating from the apex of this convexity. The nasal bosses are also markedly rugose anteriorly. The "free portion" of the nasals, from the narial notch to the tips of the nasals, measures 94 mm., and the narial notch is situated above the front margin of the second upper premolar.

It now seems clear that Peterson's species should be considered a synonym of *Menoceras arikareense* (Barbour), and that most of the dicerathere specimens from the Bridgeport Quarries should be separated from it, tentatively as *Menoceras* sp. From these quarry sites, all situated at the same geologic and topographic level in the lower part of the Marsland Formation, one skull (U.N.S.M. 62006) has been selected to demonstrate the male characters at this horizon. These are: (1) larger size than for *M. arikareense*, with the occipito-nasal length, from the rear margins of the occipital condyles to the

A New Rhinoceros from the Nebraska Miocene

tips of the nasals, being 447 mm. compared with 350 mm. for the latter species; (2) a similar, saddle-shaped skull, but with a noticeable frontal convexity, and with flaked and crescentic ridges at the anterior margin of the frontal; and (3) nasal bosses located behind the tips of the nasals and more elevated, being swollen anteriorly as well as quite rugose. In each of these characters, and most noticeably in size, the skulls from the basal Marsland are intermediate between specimens (*M. arikarensis*) from the Harrison Formation and specimens from the upper part of the Marsland Formation. The latter constitute the basis for a new species.

A NEW SPECIES OF *MENOCERAS*

Menoceras marslandensis, new species

Figs. 7-9

Holotype.—A nearly complete skull, U.N.S.M. 62003 (Figs. 7-9). The left premaxilla and both of the upper premolars are lacking, but the alveoli for these missing premolars are preserved. The right paroccipital process and right occipital condyle are absent.

Type Locality.—From about four miles east and six miles north of Hemingford, Nebraska, at U.N.S.M. Coll. Loc. Bx-28 (in the SW. $\frac{1}{4}$, NE. $\frac{1}{4}$, SW. $\frac{1}{4}$, sec. 12, T. 28 N., R. 49 W.), Box Butte Northwest Quadrangle (7.5 minute series, U.S. Geol. Survey map), Box Butte County.

Type Level.—From the upper part of the Marsland Formation (Fig. 10), in a light gray ashy sand (quarry level) situated 45-60 feet below the top of the Marsland and 5-15 feet above the upper of two volcanic ash beds. Variously considered as Medial or Late Miocene.

Diagnosis.—Male skull, larger in nearly all measurements than male skulls of either *Menoceras arikarensis* (Barbour) or *M. niobrarensis* (Peterson); see Table 2. Unlike *M. arikarensis*, the frontal probably supported a horn, evidenced by the presence of a rugose convexity on the anterior part of the frontal from which flaked and crescentic ridges radiate (Fig. 8). Also in the new species, there are flattened, rugose areas at the tips of the very long, fused but slightly cleft nasals. The length of the "free portion" of the nasals, from the narial notch to the tips of the nasals, measures 194 mm., compared with 87 mm. for *M. arikarensis* (U.N.S.M. 1147) and 94 mm. for the referred skull of *M. niobrarensis* (C.M. 1273). In the holotype of the proposed new species, the nasals and maxillar region

behind the narial notch are noticeably heavier, and there are both lateral expansion and heavier rugosity of the posterior portion of the zygomata. Associated with these strengthened characters, there was a marked weakening of the premaxillae and apparent loss of

TABLE 2
MEASUREMENTS (IN MILLIMETERS) OF THE TWO MALE SKULLS OF *Menoceras marslandensis*, NEW SPECIES, COMPARED WITH MALE SKULLS OF *M. arikareense* (BARBOUR), AND *Diceratherium armatum* MARSH.

| | <i>M. marslandensis</i> ^a | | <i>M. arikareense</i> ^b | <i>M. arikareense</i> ^c | <i>D. armatum</i> ^d |
|---|--------------------------------------|-----|------------------------------------|------------------------------------|--------------------------------|
| Occipital condyles to tips of nasals | 438 | 474 | 351 | 450 | (512) |
| Midpoint occipital crest to tips of nasals ... | 408 | 416 | 348 | ... | (503) |
| Anterior margin of P ¹ to occipital condyles | 388 | 402 | 320 | 392 | 452 |
| Narial notch to occipital crest | 323 | 324 | 295 | ... | 453 |
| Palatal notch to foramen magnum | 238 | 234 | 205 | 228 | 272 |
| Palatal notch to palatal foramina | 153 | 154 | 107 | 117 | 170 |
| Narial notch to tips of nasals | 93 | 164 | 87 | 94 | ... |
| Zygomatic breadth (maximum) | 245 | 250 | 236 | 258 | 278 |
| Width across palate to buccal sides M ² | 158 | 132 | 121 | ... | 174 |
| Orbital breadth (between notches) | 167 | 143 | 133 | ... | 210 |
| Occipital height, base condyles to crest | 135 | 140 | 122 | 160 | 159 |
| Occipital width (maximum) | 148 | 145 | 135 | ... | 150 |
| Condylar width (outer margins occ. cond.) | 80 | ... | 72 | 99 | 107 |
| Tooth row, P ¹ -M ³ (midline, to rear of M ³) | 223 | 212 | 165 | ... | 254 |
| Tooth row, P ² -M ³ (midline, to rear of M ³) | 203 | 200 | 150 | ... | ... |
| Premolars (midline, alveolus of P ¹) | 107 | 100 | 80 | ... | 123 |
| Premolars along buccal ridges | 109 | ... | 82 | ... | 127 |
| Length P ² -P ⁴ (midline, to rear of P ⁴) | 88 | ... | 63 | ... | 95 |
| Length P ² -P ⁴ (along buccal ridges) | 94 | 88 | 69 | ... | 98 |
| Molars (midline, to rear of M ³) | 118 | 112 | 89 | 113 | 133 |
| Length P ¹ (maximum) | 20 | ... | 13.5 | ... | 27 |
| Width P ¹ (maximum) | 18 | ... | 13.5 | ... | 23 |
| Length P ² (maximum) | 24 | 28 | 20 | ... | 27 |
| Width P ² (maximum) | 31 | 35 | 27 | ... | 37 |
| Length P ³ (maximum) | 30 | 28 | 20 | ... | 33 |
| Width P ³ (maximum) | 28 | 41 | 34.5 | ... | 43 |
| Length P ⁴ (maximum) | 30 | 30 | 23 | ... | 31 |
| Width P ⁴ (maximum) | 25 | 46 | 38 | ... | 47 |
| Length M ¹ (maximum) | 35 | 32 | 27.5 | ... | 47 |
| Width M ¹ (maximum) | 25 | 52 | 40 | ... | 52 |
| Length M ² (maximum) | 35 | 40 | 32 | ... | 50 |
| Width M ² (maximum) | 25 | 52 | 40 | ... | 56 |
| Length M ³ (maximum) | ... | 38 | 34.5 | ... | 37 |
| Width M ³ (maximum) | ... | 47 | 36 | ... | 47 |

^a Young male at left, U.N.S.M. 62004; holotype at right, U.N.S.M. 62003.

^b Referred, U.N.S.M. 1147.

^c Young male, holotype of *Diceratherium niobrarense* Peterson, C.M. 1271.

^d Holotype, Y.P.M. 10003.

() Estimated dimension.



FIG. 7—Lateral view of holotype male skull (U.N.S.M. 62003) of *Menoceras marslandensis*, new species, from a fossil quarry northeast of Hemingford (U.N.S.M. Coll. Loc. Bx-28), Box Butte County, Nebraska (upper part of Marsland Formation): X 11/32



FIG. 8—Dorsal view of the same skull shown as Fig. 7: X 11/32

A New Rhinoceros from the Nebraska Miocene



FIG. 9—Palatal view of the same skull shown as Figs. 7 and 8: X 11/32

| NEBRASKA MIOCENE CORRELATION CHART | | | |
|--|--|---------------------|---|
| By C. Bertrand Schultz and Thompson M. Stout | | | |
| Provincial Ages | PLIOCENE - Valentinian above | | European Standard |
| HEMINGFORDIAN | <p>• SHEEPCREEKIAN</p> <p>HEMINGFORD GROUP</p> <p>SHEEP CREEK FORMATION</p> <p>UPPER MEMBER (=LOWER SNAKE CREEK, =SAND CANYON, in pt.) Lower Snake Creek L.F.</p> <p>LOWER MEMBER (=SHEEP CREEK, s.s., =SPOTTEDTAIL, in part) Sheep Creek L.F.</p> | <p>BOX BUTTE(?)</p> | SAR-MAT-IAN |
| | <p>• MARSLANDIAN</p> <p>HEMINGFORD GROUP</p> <p>MARSLAND FORMATION</p> <p>MARSLAND AND HEMINGFORD QUARRIES</p> <p>BRIDGEPORT QUARRIES</p> | | <p>HELVETIAN - TORTONIAN</p> <p>VINDOBONIAN</p> |
| ARIKAREAN | <p>ARIKAREE GROUP</p> <p>HARRISON FORMATION</p> <p>AGATE SPRING QUARRIES</p> | | BURDIGALIAN |
| | <p>MONROE CREEK FORMATION</p> | | |
| | <p>GERING FORMATION</p> | | |
| | OLIGOCENE - Whitneyan below | | |

FIG. 10—Correlation chart of the Miocene of the Central Great Plains (High Plains) of Nebraska and adjacent states, showing the geologic placement of the principal fossil quarries that yield diceratheres. (From Schultz and Stout, 1961, Fig. 3). The black dots indicate suggested "Provincial Ages."

A New Rhinoceros from the Nebraska Miocene

the upper incisors, compared with the normal condition of these characters in *M. arikareense*. Further, the bosses for the prominent nasal horns have become more rugose anteriorly, more spatulate, and situated farther forward with respect to the nasal protuberance; these accentuate the saddle shape when the cranium is viewed from the side (Fig. 7).

SUMMARY

Two lineages of diceratheres seem to be present in the Miocene sediments of Nebraska and adjacent states, as proposed by Troxell (1921) but were confused by later workers. These are considered to represent two distinct genera, *Diceratherium* Marsh and *Menoceras* Troxell, each with successional species.

The new species here proposed, *Menoceras marslandensis*, from fossil quarries near Marsland and Hemingford in western Nebraska, is believed to be a direct descendant of the best-known dicerathere, *Menoceras arikareense* (Barbour), from the geologically older Agate Springs Fossil Quarries. The latter name preoccupies Peterson's "*Diceratherium niobrarenensis*" and "*D. cooki*." An undescribed, possibly new, species of *Menoceras* occurs in the geologically intermediate Bridgeport Quarries, and it is taken to be the intermediate species in this *Menoceras* lineage. *Diceratherium*, originally established on *D. armatum* Marsh, from the John Day Miocene of Oregon, is not only larger than *Menoceras* but anatomically distinct. *Diceratherium armatum* has also been reported from the Miocene deposits of the Central Great Plains. The *Menoceras* and *Diceratherium* lineages may have both survived into the Late Miocene and Pliocene, but more study regarding this is necessary. Similarly, the Oligocene ancestors need clarification, although it now seems virtually certain that *Subhyracodon occidentalis* (Leidy), from the Medial Oligocene (Orella Member of the Brule Formation), gave rise in the High Plains Miocene to *Diceratherium*.

ACKNOWLEDGMENTS

I am indebted to Professors C. Bertrand Schultz, Thompson M. Stout, and Harvey L. Gunderson of the University of Nebraska State Museum Staff for their aid in the preparation of this publication. Also to Dr. Craig C. Black and Dr. Mary C. Dawson of the Carnegie Museum and Dr. Elwyn L. Simons of Yale Peabody Museum for the permission to use type specimens necessary for the study, and to Dr. Malcolm C. McKenna, Messrs. Morris F. Skinner

and Beryl E. Taylor of the Frick Laboratory, American Museum of Natural History, New York, for use of comparative material from the collections. Mesdames Norma Wagner, Mary Cutler and Miss Charlene Call assisted in the final typing of the manuscript.

REFERENCES

- Barbour, Erwin H. 1906a. Notice of a new fossil rhinoceros from Sioux County, Nebraska: *Diceratherium arikarense*. Publ. Nebr. Geol. Survey, ser. 1, 2 (4): ii + 313-318, Figs. 1-5. [Published June 15, 1906, according to affidavit of R. D. Woodruff of Woodruff Printing Company, 1968.]
- . 1906b. Notice of a new Miocene rhinoceros, *Diceratherium arikarense*. Science, new ser. (ser. 2), 24 (624): 780-781, Figs. 1-2. [Published December 14, 1906; submitted July 4, 1906.]
- . 1908. The skull of *Moropus*. Publ. Nebr. Geol. Survey, ser. 1, 3 (2): 12 pages, Figs. 1-5, Pls. 1-2.
- . 1909a. A slab from the bone beds of Sioux County. Publ. Nebr. Geol. Survey, ser. 1, 3 (7): 3 pages, Pl. 1.
- . 1909b. Restoration of *Diceratherium arikarense*, a new form of panel mount. *Ibid.*, ser. 1, 3 (8): 3 pages, Pl. 1.
- Colbert, Edwin H. 1935. Nebraska—fifteen million years ago. Nat. Hist. (Amer. Mus. Nat. Hist.) 35 (1): 37-46, unnumbered figs., table.
- Cook, Harold J. 1965. Runningwater Formation, Middle Miocene of Nebraska. Amer. Mus. Novitates 2227: 1-8, Figs. 1-3.
- Green, Morton. 1958. Arikareean rhinoceroses from South Dakota. Jour. Paleont. 32 (3): 587-594, Figs. 1-9, Tables 1-4.
- Holland, W. J., and O. A. Peterson. 1913. The Osteology of the Chalicotheroidea, with special reference to a mounted skeleton of *Moropus elatus* Marsh, now installed in the Carnegie Museum. Mem. Carnegie Mus. 3 (2): i-xvi, 189-411, Figs. 1-115, Pls. 48-77.
- Loomis, Frederic B. 1908. Rhinocerotidae of the Lower Miocene. Amer. Jour. Sci. 176 (ser. 4, 26, whole no. 176), no. 151, art. 4: 51-64, Figs. 1-15.
- Lugn, Alvin L., Sr. 1939. Classification of the Tertiary System in Nebraska. Bull. Geol. Soc. Amer. 50 (8): 1245-1276, Pl. 1, Tables 1-2.
- Macdonald, James R. 1963. The Miocene faunas from the Wounded Knee area of western South Dakota. Bull. Amer. Mus. Nat. Hist. 125 (3): 139-238, Figs. 1-30, Tables 1-2, Maps 1-2.
- Macdonald, James R., and C. Bertrand Schultz. 1956. *Arretotherium fricki*, a new Miocene anthracothere from Nebraska. Bull. Univ. Nebr. State Mus. 4 (3): iii + 52-58, Figs. 1-3.

A New Rhinoceros from the Nebraska Miocene

- McKenna, Malcolm C. 1965. Stratigraphic nomenclature of the Miocene Hemingford Group, Nebraska. *Amer. Mus. Novitate* 2228: 1-21, Fig. 1.
- Marsh, Othniel C. 1875. Notice of new Tertiary mammals, IV. *Amer. Jour. Sci. and Arts* 109 (ser. 3, 9) no. 51: 239-250.
- Matthew, William D. 1931. Critical observations on the phylogeny of the rhinoceroses. *Univ. Calif. Publ., Bull. Dept. Geol. Sci.* 20 (1): 1-9, Figs. 1-2.
- Osborn, Henry F. 1898. The extinct rhinoceroses. *Mem. Amer. Mus. Nat. Hist.*, ser. 1, 1 (3): 75-164 + xviii, Figs. 1-49, Pls. 12A-20.
- . 1899. Frontal horn on *Aceratherium incisivum*. *Science*, new ser. (ser. 2), 9 (214): 161-162, Pl. 1.
- . 1900. Phylogeny of the rhinoceroses of Europe. *Bull. Amer. Mus. Nat. Hist.* 13 (19): 229-267, Figs. 1-16, table.
- . 1904. New Miocene rhinoceroses, with revision of known species. *Ibid.* 20 (27): 307-326, Figs. 1-21.
- Peterson, O. A. 1906a. The Agate Spring fossil quarry. *Ann. Carnegie Mus.*, 3 (4), art. 18: 487-494, 5 figs., 1 plate.
- . 1906b. The Miocene beds of western Nebraska and eastern Wyoming and their vertebrate fauna. *Ibid.*, 4 (1): 21-72, 20 figs.
- . 1906c. Preliminary description of two new species of the genus *Diceratherium* Marsh, from the Agate Spring Fossil Quarry. *Science*, new ser. (ser. 2), 24 (609): 281-283. [Published August 31, 1906; submitted August 15, 1906.]
- . 1920. The American diceratheres. *Mem. Carnegie Mus.* 7 (6): 399-476, Figs. 1-37, Pls. 57-66.
- Roberts, R. J. 1966. The history of Agate Springs. *Nebr. Hist.* 47 (3): 265-293, illus.
- Schultz, C. Bertrand. 1938. The Miocene of western Nebraska. *Amer. Jour. Sci.* 235 (ser. 5, 35), no. 210: 441-444, table.
- . 1942. A review of the Daimonelix problem. *Univ. Nebr. Studies*, new ser., *Studies Sci. Tech.*, no. 2: 1-30, Figs. 1-17.
- . 1966. The Agate Springs Fossil Quarries. *Univ. Nebr. State Mus.*, *Museum Notes* 30: 1-7, illus.
- Schultz, C. Bertrand, and Charles H. Falkenbach. 1949. Promerycochoerinae, a new subfamily of oreodonts. *Bull. Amer. Mus. Nat. Hist.* 93 (3): 69-198, Figs. 1-26, Tables 1-8, Charts 1-6.
- Schultz, C. Bertrand, Charles H. Falkenbach, and Carl F. Vondra. 1967. The Brule-Gering (Oligocene-Miocene) contact in the Wildcat Ridge area of western Nebraska. A guide for the stratigraphic collection of fossil mammals. *Bull. Univ. Nebr. State Mus.* 6 (4): 43-58, Figs. 1-7.

- Schultz, C. Bertrand, and Henry P. Reider. 1943. Modern methods in the preparation of fossil skeletons. *The Compass of Sigma Gamma Epsilon*, 23 (4): 268-291, 22 figs.
- Schultz, C. Bertrand, and Thompson M. Stout. 1961. Field conference on the Tertiary and Pleistocene of Western Nebraska (with contributions by Charles H. Falkenbach and Lloyd G. Tanner; and field conference assistance from Harold J. Cook and A. L. Lugn). *Univ. Nebr. State Mus. Special Publ.* 2: 1-55, Figs. 1-47, Chart, Map. (Also: inserted chart on terrace-fills, and erratum.)
- Simpson, George G. 1945. The principles of classification and a classification of mammals. *Bull. Amer. Mus. Nat. Hist.* 85: i-xvi, 1-350.
- Stecher, Robert M., C. Bertrand Schultz, and Lloyd G. Tanner. 1962. A Middle Miocene rhinoceros quarry in Morrill County, Nebraska (with notes of hip disease in *Diceratherium*). *Bull. Univ. Nebr. State Mus.* 4 (7): 101-111, Figs. 1-6.
- Stout, Thompson M., V. H. Dreeszen, and W. W. Caldwell. 1965. Guidebook for Field Conference "D," Central Great Plains (Guidebooks for the 7th Congress of INQUA: Internatl. Assoc. Quaternary Research, *Nebr. Acad. Sci.*): 1-24, 62 figs.
- Tanner, Lloyd G. 1962. University of Nebraska Trailside Museum at Fort Robinson. *Univ. Nebr. State Mus., Museum Notes* 18: 1-4, illus.
- . 1965a. Additional information regarding the stratigraphic occurrences of the Miocene diceratheres of Nebraska (Abstr.) *Proc. Nebr. Acad. Sci., 75th Ann. Meeting*: 18.
- . 1965b. The geologic history of Nebraska's rhinoceroses. (Abstr.) *Ibid.*, 75th Ann. Meeting: 41-45.
- . 1967. A new species of rhinoceros, *Aphelops kimballensis*, from the Latest Pliocene of Nebraska. *Bull. Univ. Nebr. State Mus.* 6 (1): 1-16, Fig. 1, Pls. 1-5, Tables 1-2.
- Troxell, Edward L. 1921. A study of *Diceratherium* and the diceratheres. *Amer. Jour. Sci.* 202 (ser. 5, 2) no. 10, art. 15: 197-208, Figs. 1-8.
- Wood, Horace E., II. 1927. Some Early Tertiary rhinoceroses and hyracodonts. *Bull. Amer. Paleont.* 13 (whole no. 50): i, 5-104, Pls. 1-7, Tables 1-7.
- . 1929. American Oligocene rhinoceroses; a postscript. *Jour. Mammal.* 10 (1): 63-75.
- . 1933. A fossil rhinoceros (*Diceratherium armatum* Marsh) from Gallatin County, Montana. *Proc. U.S. Natl. Mus.* 82 (7): 1-4, Pls. 1-3.