

3-30-2001

# Do species and functional groups differ in acquisition and use of C, N and water under varying atmospheric CO<sub>2</sub> and N availability regimes? A field test with 16 grassland species

Peter B. Reich

*University of Minnesota, St. Paul, MN*

David Tilman

*University of Minnesota, St. Paul, MN*

Joseph Craine

*University of California, Berkeley, CA*

David Ellsworth

*Brookhaven National Laboratory, Upton, NY*

Mark G. Tjoelker

*University of Minnesota, St. Paul, MN*

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/natrespapers>

 Part of the [Natural Resources and Conservation Commons](#)

---

Reich, Peter B.; Tilman, David; Craine, Joseph; Ellsworth, David; Tjoelker, Mark G.; Knops, Johannes M. H.; Wedin, David A.; Naeem, Shahid; Bahauddin, Dan; Goth, Jenny; Bengtson, Wendy; and Lee, Tali D., "Do species and functional groups differ in acquisition and use of C, N and water under varying atmospheric CO<sub>2</sub> and N availability regimes? A field test with 16 grassland species" (2001). *Papers in Natural Resources*. 111.  
<http://digitalcommons.unl.edu/natrespapers/111>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Natural Resources by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Peter B. Reich, David Tilman, Joseph Craine, David Ellsworth, Mark G. Tjoelker, Johannes M. H. Knops, David A. Wedin, Shahid Naeem, Dan Bahaiddin, Jenny Goth, Wendy Bengtson, and Tali D. Lee

# Do species and functional groups differ in acquisition and use of C, N and water under varying atmospheric CO<sub>2</sub> and N availability regimes? A field test with 16 grassland species

Peter B. Reich<sup>1</sup>, David Tilman<sup>2</sup>, Joseph Craine<sup>3</sup>, David Ellsworth<sup>4</sup>, Mark G. Tjoelker<sup>1</sup>, Jean Knops<sup>5</sup>, David Wedin<sup>6</sup>, Shahid Naeem<sup>7</sup>, Dan Bahauddin<sup>1</sup>, Jenny Goth<sup>1</sup>, Wendy Bengtson<sup>1</sup>, Tali D. Lee<sup>1</sup>

<sup>1</sup>Department of Forest Resources, University of Minnesota, St. Paul, MN 55108 USA; <sup>2</sup>Department of Ecology, Evolution and Behaviour, University of Minnesota, St. Paul, MN 55108 USA; <sup>3</sup>Department of Integrative Biology, University of California, Berkeley, CA 94720 USA; <sup>4</sup>Department of Environmental Science, Brookhaven National Laboratory, Upton, NY 11973 USA; <sup>5</sup>School of Biological Sciences, University of Nebraska, Lincoln, NE 68583 USA; <sup>6</sup>School of Natural Resource Sciences, University of Nebraska, Lincoln, NE 68583 USA; <sup>7</sup>Department of Zoology, University of Washington, Seattle, WA 98195 USA

## Summary

Author for correspondence:  
Peter B. Reich  
Tel: +1 612 624 4270  
Fax: +1 612 625 5212  
Email: preich@forestry.umn.edu

Received: 11 September 2000

Accepted: 11 January 2000

- To evaluate whether functional groups have a similar response to global change, the responses to CO<sub>2</sub> concentration and N availability of grassland species from several functional groups are reported here.
- Sixteen perennial grassland species from four trait-based functional groups (C<sub>3</sub> grasses, C<sub>4</sub> grasses, non-leguminous forbs, legumes) were grown in field monocultures under ambient or elevated (560 μmol mol<sup>-1</sup>) CO<sub>2</sub> using free-air CO<sub>2</sub> enrichment (FACE), in low N (unamended field soil) or high N (field soil +4 g N m<sup>-2</sup> years<sup>-1</sup>) treatments.
- There were no CO<sub>2</sub> × N interactions. Functional groups responded differently to CO<sub>2</sub> and N in terms of biomass, tissue N concentration and soil solution N. Under elevated CO<sub>2</sub>, forbs, legumes and C<sub>3</sub> grasses increased total biomass by 31%, 18%, and 9%, respectively, whereas biomass was reduced in C<sub>4</sub>-grass monocultures. Two of the four legume species increased biomass and total plant N pools under elevated CO<sub>2</sub>, probably due to stimulated N-fixation. Only one species markedly shifted the proportional distribution of below- vs aboveground biomass in response to CO<sub>2</sub> or N.
- Although functional groups varied in responses to CO<sub>2</sub> and N, there was also substantial variation in responses among species within groups. These results suggest that current trait-based functional classifications might be useful, but not sufficient, for understanding plant and ecosystem responses to elevated CO<sub>2</sub> and N availability.

**Key words:** Functional groups, elevated carbon dioxide, nitrogen, N availability, grasses, forbs, legumes, biomass.

© *New Phytologist* (2001) **150**: 435–448

## Introduction

Generalizing about responses of different plant species to elevated atmospheric CO<sub>2</sub> concentrations or N deposition remains an elusive goal in global change biology. Given the wide variety of species, any kind of grouping that simplifies the variation among species while providing predictive power will

significantly advance the field. Based on well-known differences among species in important intrinsic traits, a variety of hypotheses have been developed about potential functional group differences in response to CO<sub>2</sub> and N enrichment.

1) Intrinsic differences in photosynthetic biochemistry should lead to markedly greater responsiveness to elevated CO<sub>2</sub> for C<sub>3</sub> than C<sub>4</sub> plants (H<sub>1</sub>). Although supported by

theory and early field studies (Percy & Ehleringer, 1984; Curtis *et al.*, 1989), more recent studies suggest such differences may be less pronounced (Owensby *et al.*, 1993; Wand *et al.*, 1999), especially during dry periods.

2) Productivity in N-fixing legumes may be stimulated by elevated CO<sub>2</sub> more than in nonfixers (H<sub>2</sub>), because the former should be less N-limited. It has long been known that elevated CO<sub>2</sub> stimulates legume growth and N<sub>2</sub> fixation (Finn & Brun, 1982; Zanetti *et al.*, 1996). N-fixing species have often shown a stronger biomass response to elevated CO<sub>2</sub> than nonfixing species (Soussana & Hartwig, 1996; Clark *et al.*, 1997; Hebeisen *et al.*, 1997; Lüscher & Nösberger, 1997, 1998; Schenk *et al.*, 1997), although little work to date has been on wild species at naturally low levels of N availability.

3, 4) A series of related hypotheses posit that increases in CO<sub>2</sub> or N supply should lead to a more pronounced growth increase in species of given strategies, habitats, or growth rates. For our study we propose the hypothesis (H<sub>3</sub>) that C<sub>3</sub> grasses considered more disturbance adapted and nitrophilic, should respond more to increase in N supply than C<sub>4</sub> grasses (cf Wedin & Tilman, 1996). Moreover, being less N-limited, the legumes should also be less responsive to N addition than the non-fixers (H<sub>4</sub>).

5) In addition to understanding the effects of elevated CO<sub>2</sub> or N singularly, given the potential for the CO<sub>2</sub> fertilization effect to be modulated by N-supply, it is important to study species and functional group responses to combinations of these two elements. Many ecosystem models theorize that CO<sub>2</sub> responses are constrained by N limitation (but see Cannell & Thornley, 1998) and actual evidence is mixed (Larigauderie *et al.*, 1988; Owensby *et al.*, 1994; Leadley & Körner, 1996; Lloyd & Farquhar, 1996; Poorter *et al.*, 1996; Volin & Reich, 1996; Zak *et al.*, 2000). If the CO<sub>2</sub> response of species is generally N-limited, then nonlegumes, without the ability to fix atmospheric N and therefore modulate their own N supply, should demonstrate smaller responses to elevated CO<sub>2</sub> than legumes (see H<sub>2</sub>) and be more likely to show a CO<sub>2</sub> × N interaction (H<sub>5</sub>).

6) For individually grown plants, usually in their first year, root fraction (root biomass as a fraction of total biomass) adjusted for ontogenetic drift is unaffected by CO<sub>2</sub> (Curtis & Wang, 1998; Reich, 2001) and is lower under enhanced N supply (Poorter & Nagel, 2000; Reich, 2001). Data for older plants or stands are rare. If assemblages over multiple years behave similarly as young, individual plants, one might hypothesize that elevated CO<sub>2</sub> would have no effect (H<sub>6a</sub>) and N addition a decreasing effect (H<sub>6b</sub>) on root fraction in our experiment. Since root fraction incorporates both allocation and turnover, neither of which are well documented in the field (Reich, 2001), these hypotheses are proposed as null models.

7) Effects of treatments on biomass and physiology should be reflected in plot-scale resource availability. Assuming that greater CO<sub>2</sub> and N supply both lead to increased biomass, we hypothesize (H<sub>7</sub>) that soil solution N concentration and

percentage soil water should decrease under both treatments. However, increased N supply should compensate for increased N uptake, minimizing the decline in soil N compared with that under elevated CO<sub>2</sub> (H<sub>7a</sub>). Under elevated CO<sub>2</sub>, reduced leaf level water loss could minimize the decline in percentage soil water compared to that experienced in the high N treatment (H<sub>7b</sub>).

Although there are an increasing number of tests of the CO<sub>2</sub> × N interaction hypothesis in general and of the functional group-related hypotheses raised above, few have been done in field settings where both root and shoot processes can be quantified for more than a small number of species. To help fill this gap, we addressed these issues using an experiment comprising 16 grassland species from four functional groups grown in 128 monoculture plots that is a part of a larger experiment (BioCON) designed to test interactions among species diversity, elevated CO<sub>2</sub> and N deposition (Reich *et al.*, 2001). In particular, we assessed whether functional groups, growing in monoculture plots in a free-air CO<sub>2</sub> enrichment (FACE) experiment, differed in their acquisition and use of C, N and water in response to combined treatment with elevated CO<sub>2</sub> and N deposition.

## Materials and Methods

The BioCON (Biodiversity, CO<sub>2</sub> and N) experiment (Reich *et al.*, 2001) (<http://www.swan.lter.umn.edu/biocon/>) is located at the Cedar Creek Natural History area, a National Science Foundation, Long-Term Ecological Research site in Minnesota, USA (lat. 45° N, Long. 93° W). The region has a continental climate with cold winters, warm summers (mean January and July temperatures of -11 and 22°C), and precipitation averaging 660 mm y<sup>-1</sup>. The soils are derived from a glacial outwash sand plain and are sandy and nitrogen poor. Plots were established on a secondary successional grassland after removing the previous vegetation.

Our study included 128 individual monoculture plots (each 2 × 2 m), a subset of all plots, distributed nearly equally among six 20-m diameter experimental areas (rings). In three elevated CO<sub>2</sub> rings, a free-air CO<sub>2</sub> enrichment (FACE) system (Lewin *et al.*, 1994) was used during the 1998 and 1999 growing seasons to maintain the CO<sub>2</sub> concentration at 560 μmol mol<sup>-1</sup>. Three ambient rings (368 μmol mol<sup>-1</sup> CO<sub>2</sub>) were treated identically but without additional CO<sub>2</sub>. The experimental treatments were arranged in complete factorial combination of CO<sub>2</sub> (ambient or elevated), species (a total of 16, four from each of four functional groups), and N level (low and high) for a 2 × 16 × 2 design with two replicates. Each plot was planted in 1997 with 12 g m<sup>-2</sup> of seed. The design consisted of a split-plot arrangement of treatments in a completely randomized design. CO<sub>2</sub> treatment is the whole-plot factor and is replicated three times among the six rings. The subplot factors of species identity and N treatment were randomly assigned and replicated in individual plots among the six rings. CO<sub>2</sub> was added in

elevated treatments during all daylight hours from April 9 to October 16, 1998, and from April 20 to November 9, 1999. Tests found no direct effect of elevated CO<sub>2</sub> on dark respiration (Tjoelker *et al.*, 2001). During CO<sub>2</sub> enrichment periods, 1-min averages were within 10% of the target concentration 94% of the time in 1998 and 95% of the time in 1999. Beginning in 1998, the plots assigned to the high N treatment were amended with 4 g N m<sup>-2</sup> yr<sup>-1</sup>, applied over three dates each year, while the low N treatment soil was unamended. During the two growing seasons of treatments during this study, no severe dry periods occurred.

The 16 perennial species used in this study were all native or naturalized to the Cedar Creek Natural History Area. They include four C<sub>4</sub> grasses (*Andropogon gerardii* Vitman, *Bouteloua gracilis*, *Schizachyrium scoparium* (Michaux) Nash, *Sorghastrum nutans* (L.) (Nash), four C<sub>3</sub> grasses (*Agropyron repens* (L.) Beauv., *Bromus inermis* Leysser, *Koeleria cristata* Pers, *Poa pratensis* L.), four N-fixing legumes (*Amorpha canescens* Pursh, *Lepedeza capitata* Michaux, *Lupinus perennis* L., *Petalostemum villosum* Nutt.) and four nonN-fixing herbaceous species (*Achillea millefolium* L., *Anemone cylindrica* A. Gray, *Asclepias tuberosa* L., *Solidago rigida* L.). Species hereafter are referred to by their genus. Monocultures of all species were replicated twice at all four combinations of CO<sub>2</sub> and N levels. Plots were regularly weeded to remove unwanted species. In June and August of 1998 and 1999 we assessed above- and belowground (0–20 cm) biomass and soil solution N concentrations (extracted using 0.01 mol KCl). For most analyses in this paper we use the mean values per plot from these four harvests. A 10 × 100 cm strip was clipped at just above the soil surface, all matter was collected, sorted to live material and senesced litter, dried and weighed. Roots were sampled at 0–20 cm depth using three 5-cm cores in the area used for the aboveground biomass clipping. Roots were washed, sorted into fine (< 1 mm diameter) and coarse classes and crowns, dried and weighed. Volumetric soil moisture levels (0–20 cm depth) were assessed periodically (on 18 sampling periods) in all plots in 1998 and 1999 using time-domain reflectometry (Baker, 1990). A composite sample was taken from aboveground and belowground biomass from each plot from the August harvests of each year, ground and analysed for N using a CHN analyser (Carlo-Erba Strumatzone, Milan, Italy). To estimate total plant N stocks we multiplied whole plant percentage N (averaged across years) by the mean whole plot biomass (averaged from all harvests).

### Data analysis

ANOVA was used in two complementary ways, including either species or functional group as a treatment effect. In ANOVA all treatment effects were considered fixed. Using *F*-tests, the effect of CO<sub>2</sub> (1 df) was tested against the random effect of ring nested within CO<sub>2</sub> (4 df). The main effects of functional group (3 df) was tested against the random effect of species nested within functional group (12 df). The main

effects of species (15 df), and N (1 df), and interactions between CO<sub>2</sub>, N, and either functional group or species were tested against the residual error. Additionally, to explicitly test hypotheses about preplanned specific functional group contrasts (e.g. C<sub>3</sub> vs C<sub>4</sub>), their effects were partitioned into single-degree-of-freedom contrasts. We also evaluated the proportional distribution of biomass aboveground vs belowground. Since this is a plot-scale measure for assemblages that had been developing in the field for three seasons it should not be taken as a measure of allocation. Instead, it reflects the balance between allocation and turnover (Reich, 2001). We also used separate and same slopes regression to test for slope and intercept differences among treatments in the relationships between soil resources (e.g. % soil water or N) and fine root biomass. We used repeated measures ANOVA to test whether responses to CO<sub>2</sub> or N varied among times of year (June vs August), among years, or among all four harvests. There were no interactions between treatments and time, hence all results are presented averaged across harvests and years. All statistical analyses were made using JMP 4.0.1 software.

### Results

Functional groups differed significantly in shoot, root and total biomass (Tables 1, 2; Fig. 1), with C<sub>3</sub> grass monocultures highest and legumes lowest. For total biomass, averaged across all species or functional groups, there were marginally significant main effects of CO<sub>2</sub> (+11%, mean of enhancement) and N (+7%) (Tables 2, 3; Fig. 1) and there were no significant CO<sub>2</sub> × N interactions (Tables 2, 3). In fact, there were no significant CO<sub>2</sub> × N interactions, or significant three-way interactions involving CO<sub>2</sub> and N, for any of the measured variables in this study (refuting H<sub>5</sub>), and hence responses to CO<sub>2</sub> and N will be presented separately, pooled across levels of the other variable.

#### Biomass response to elevated CO<sub>2</sub>

There were marked differences among functional groups in terms of biomass response to CO<sub>2</sub> (Tables 1, 2; Figs 1, 2). Forbs, legumes, and C<sub>3</sub> grasses increased total biomass by 31%, 18%, and 9%, respectively, under elevated CO<sub>2</sub> whereas C<sub>4</sub> grass monocultures had 3% lower total biomass. The interaction term was significant (*P* < 0.10) for the CO<sub>2</sub>-group interaction and more so (*P* < 0.002) for the C<sub>3</sub> vs C<sub>4</sub> contrast (Table 2), indicating that the C<sub>4</sub> grasses did have less enhancement of biomass than C<sub>3</sub> species in general, supporting (H<sub>1</sub>). There was no evidence that the N-fixing legumes responded more positively to elevated CO<sub>2</sub> than the nonfixing C<sub>3</sub> species (Figs 1, 2; Tables 1, 2), refuting (H<sub>2</sub>). The effects of CO<sub>2</sub> on biomass, and differences in response among functional groups, were largely manifest belowground (Tables 1, 2).

There was substantial variation in response to elevated CO<sub>2</sub> among species, shown by significant species-CO<sub>2</sub> interactions

**Table 1** Means of belowground and aboveground biomass of species under contrasting CO<sub>2</sub> and N treatments for 128 plots. Values shown are for 4 plots per species at each CO<sub>2</sub> or N level, pooled across treatments otherwise and averaged over four harvests in two years. The mean standard errors of the adjusted least squares means (LSM) for belowground and aboveground biomass, respectively, were 59 and 21 g m<sup>-2</sup> for species and 103 and 14 for functional groups


Functional group	Species	Belowground biomass (g m <sup>-2</sup> )				Aboveground biomass (g m <sup>-2</sup> )			
		Ambient CO <sub>2</sub>	Elevated CO <sub>2</sub>	Low N	High N	Ambient CO <sub>2</sub>	Elevated CO <sub>2</sub>	Low N	High N
C <sub>4</sub> grass	<i>Andropogon gerardii</i>	575	525	543	557	261	283	263	282
	<i>Bouteloua gracilis</i>	472	562	471	564	253	200	209	244
	<i>Schizachyrium scoparium</i>	359	349	293	415	177	153	133	196
	<i>Sorghastrum nutans</i>	569	513	614	468	248	221	199	270
	Mean	493	489	480	502	236	215	204	247
C <sub>3</sub> grass	<i>Agropyron repens</i>	806	1114	883	1037	308	307	282	333
	<i>Bromus inermis</i>	828	755	750	834	306	281	253	333
	<i>Koeleria cristata</i>	645	714	666	692	253	225	194	284
	<i>Poa pratensis</i>	991	1058	929	1120	168	200	123	245
	Mean	817	910	807	920	256	252	210	299
Forb	<i>Achillea millefolium</i>	700	1014	944	770	268	293	266	295
	<i>Anemone cylindrica</i>	183	288	324	147	36	53	60	29
	<i>Asclepias tuberosa</i>	97	172	91	179	11	14	12	13
	<i>Solidago rigida</i>	597	690	706	581	315	386	328	372
	Mean	395	541	516	419	160	185	167	179
Legume	<i>Amorpha canescens</i>	172	160	208	124	59	26	59	26
	<i>Lespedeza capitata</i>	274	407	315	366	67	153	143	77
	<i>Lupinus perennis</i>	216	303	270	250	279	383	339	323
	<i>Petalostemum villosum</i>	135	122	118	140	142	39	27	154
	Mean	200	246	226	220	137	151	143	145

**Table 2** ANOVA summary for biomass, N, and soil water measures in 128 plots

Parameter	R <sup>2</sup>	CO <sub>2</sub> × N × Functional group analyses							
		CO <sub>2</sub>	N	Group	CO <sub>2</sub> × Group	CO <sub>2</sub> × (C <sub>3</sub> vs C <sub>4</sub> )	CO <sub>2</sub> × C <sub>3</sub> N-fixer vs C <sub>3</sub> nonfixer	N × group	N vs N-fixer vs nonfixer
Total biomass	0.88	<b>0.05</b>	<b>0.10</b>	<b>0.02</b>	<b>0.10</b>	<b>0.002</b>	<b>0.04</b>	<b>0.001</b>	<b>0.08</b>
Belowground biomass	0.87	<b>0.08</b>	0.77	<b>0.006</b>	0.14	<b>0.005</b>	<b>0.02</b>	<b>0.007</b>	0.67
Aboveground biomass	0.84	0.47	<b>0.0003</b>	0.51	0.30	0.14	0.77	<b>0.01</b>	<b>0.0006</b>
Root fraction	0.65	0.40	<b>0.02</b>	0.39	0.70	0.31	0.72	0.92	0.36
Total plant N pool	0.80	0.27	< <b>0.0001</b>	0.18	0.51	0.18	0.65	<b>0.0006</b>	< <b>0.0001</b>
Belowground percentageN	0.85	0.30	<b>0.03</b>	<b>0.002</b>	0.38	0.19	0.13	<b>0.01</b>	<b>0.0007</b>
Aboveground percentageN	0.89	<b>0.03</b>	<b>0.005</b>	<b>0.07</b>	<b>0.02</b>	< <b>0.0001</b>	<b>0.08</b>	<b>0.002</b>	<b>0.0002</b>
% soil water	0.70	0.60	<b>0.0003</b>	0.11	0.11	0.65	0.11	<b>0.009</b>	0.15
Soil nitrate concentration	0.69	<b>0.08</b>	<b>0.001</b>	<b>0.002</b>	<b>0.03</b>	<b>0.01</b>	0.18	<b>0.009</b>	0.47
Soil N concentration	0.74	0.22	<b>0.0005</b>	<b>0.002</b>	0.28	<b>0.07</b>	0.71	<b>0.03</b>	0.84

R<sup>2</sup> shown for whole model. *P*-values shown for main effects and interactions (bolded when *P* < 0.10). Interactions involving CO<sub>2</sub> × N were not significant at *P* < 0.10) and hence are not shown. Group, functional group contrasts of the four groups; C<sub>3</sub> vs C<sub>4</sub>, C<sub>4</sub> grasses vs all other C<sub>3</sub> species; C<sub>3</sub> N-fixer vs C<sub>3</sub> nonfixer, legumes vs forbs plus C<sub>3</sub> grasses; N-fixer vs nonfixer, legumes vs all other nonfixer species.

**Fig. 1** Total biomass (aboveground plus belowground, 0–20 cm depth) ( $\text{g m}^{-2}$ ) of 16 species as affected by elevated vs ambient  $\text{CO}_2$  treatments (pooled across N treatments) and by high N (addition of  $4 \text{ g N m}^{-2} \text{ yr}^{-1}$ ) vs unamended soil treatments (pooled across  $\text{CO}_2$  treatments). Top panel: grey columns, elevated  $\text{CO}_2$ ; black columns, ambient  $\text{CO}_2$ . Bottom panel: grey columns, low N; white columns, high N. Each value is a mean of four harvests (June and August in each of 1998 and 1999) for 4 plots per species-treatment combination. The mean standard errors of the adjusted least squares means (LSMs) were  $66 \text{ g m}^{-2}$  for species. Statistical details provided in Tables 2 and 3.



**Table 3** ANOVA summary for biomass, N, and soil water measures in 128 plots

Parameter	$\text{CO}_2 \times \text{N} \times \text{Species}$ analyses					
	$R^2$	$\text{CO}_2$	N	Species	$\text{CO}_2 \times \text{Species}$	N $\times$ Species
Total biomass	0.94	<b>0.06</b>	<b>0.07</b>	<b>&lt; 0.0001</b>	<b>0.01</b>	<b>0.01</b>
Belowground biomass	0.93	0.11	0.73	<b>&lt; 0.0001</b>	<b>0.04</b>	<b>0.02</b>
Aboveground biomass	0.94	0.53	<b>&lt; 0.0001</b>	<b>&lt; 0.0001</b>	<b>0.0009</b>	<b>0.0004</b>
Root fraction	0.87	0.33	<b>0.004</b>	<b>&lt; 0.0001</b>	<b>0.02</b>	<b>0.003</b>
Total plant N pool	0.91	0.29	<b>&lt; 0.0001</b>	<b>&lt; 0.0001</b>	<b>0.02</b>	<b>0.0003</b>
Belowground percentageN	0.91	0.34	<b>0.04</b>	<b>&lt; 0.0001</b>	0.98	<b>0.05</b>
Aboveground percentageN	0.96	<b>0.01</b>	<b>0.001</b>	<b>&lt; 0.0001</b>	<b>0.01</b>	<b>&lt; 0.0001</b>
% soil water	0.82	0.58	<b>0.0005</b>	<b>&lt; 0.0001</b>	0.68	0.13
Soil nitrate concentration	0.86	<b>0.09</b>	<b>0.0004</b>	<b>&lt; 0.0001</b>	<b>0.002</b>	0.12
Soil N concentration	0.87	0.15	<b>0.0003</b>	<b>&lt; 0.0001</b>	<b>0.01</b>	0.29

$R^2$  shown for whole model.  $P$ -values shown for main effects and interactions (bolded when  $P < 0.10$ ), except those involving  $\text{CO}_2 \times \text{N}$  which were not significant at  $P < 0.10$ .

for shoot, root and total biomass (Tables 1, 3). Seven species increased total biomass under elevated  $\text{CO}_2$  by at least  $100 \text{ g m}^{-2}$ , including three forbs, two  $\text{C}_3$  grasses and two legumes (Fig. 1). The largest responses to elevated  $\text{CO}_2$  were shown by a forb, *Achillea* ( $+339 \text{ g m}^{-2}$ ,  $+35\%$ ), a  $\text{C}_3$  grass, *Agropyron* ( $+307 \text{ g m}^{-2}$ ,  $+28\%$ ) and two legumes, *Lespedeza* ( $+218 \text{ g m}^{-2}$ ,  $+64\%$ ) and *Lupinus* ( $+190 \text{ g m}^{-2}$ ,  $+38\%$ ), all of which were statistically significant ( $P < 0.05$ ) using post hoc tests. However, some species within all three of these  $\text{C}_3$  functional groups also had modest or minimal responses to elevated  $\text{CO}_2$  (Fig. 1).

### Biomass response to N addition


Functional groups responded differently ( $P < 0.01$ ) to N treatment in terms of both aboveground and belowground biomass (Tables 1, 2). At high N,  $\text{C}_3$  grasses showed the greatest increase in total biomass of all functional groups (Tables 1, 2; Figs 1, 2), supporting ( $\text{H}_3$ ), and the legumes responded less positively in general than the nonfixers, supporting ( $\text{H}_4$ ). However, the forb group also failed to respond to high N with increased biomass production. Moreover, except within

the  $\text{C}_3$  grass group, variation in biomass response to N among species within functional groups was substantial, shown by (Table 1, Fig. 1).

There were significant species–N interactions for shoot, root and total biomass (Tables 1, 3; Fig. 1). Seven species increased total biomass under high N by at least  $140 \text{ g m}^{-2}$ , including all four  $\text{C}_3$  grasses, two  $\text{C}_4$  grasses, and a legume, with *Poa* ( $+313 \text{ g m}^{-2}$ ,  $+30\%$ ), *Agropyron* ( $+206 \text{ g m}^{-2}$ ,  $+18\%$ ) and *Schizachyrium* ( $+163 \text{ g m}^{-2}$ ,  $+37\%$ ) showing the largest increases.

### Variation among functional groups and species

How do differences among functional groups compare with differences among species? We assessed the coefficient of variation (CV) of functional group and species means, and of the responses to treatments, for many variables. Results for total biomass are representative of the general trends. The CV among groups for total biomass at each of the four  $\text{CO}_2$  and N treatment combinations was slightly less than the CV for all species assessed collectively (Table 4). However, the CV among species within groups varied extremely among groups, from as low as 15–20% for  $\text{C}_3$  grasses to as high as 70–80%



**Fig. 2** Proportional response of total biomass, plant N pool, percentage soil water, and soil solution N concentration to  $\text{CO}_2$  and N treatments (otherwise pooled) on average for species in four functional groups. See Table 2 for statistics.

for the forbs. The response of biomass to  $\text{CO}_2$  also had a lower CV among groups (102%) than among all species (187%).

#### Proportional biomass distribution under elevated $\text{CO}_2$ and N treatments

Functional groups did not differ significantly in the proportion of total biomass found belowground, that is root fraction (Tables 2, 5). The  $\text{C}_4$  grasses had root fractions between 67% and 69%, the  $\text{C}_3$  grasses had root fractions between 74% and 86%, and species within other functional groups varied

widely. Among all species, *Lupinus* had the lowest root fraction (43%) but other legumes had intermediate or high root fractions (69–82%).

Root fractions under elevated  $\text{CO}_2$  (0.74 on average) were not significantly different than under ambient  $\text{CO}_2$  (0.72). Root fraction was lower ( $P < 0.005$ ) in the high N (0.71) than the low N treatment (0.75). These results support  $H_6$ . Root fraction was unrelated to total biomass. There were no significant functional group  $\times$  N or group– $\text{CO}_2$  interactions for root fraction (Table 2), but there were significant species  $\times$   $\text{CO}_2$  and species–N interactions (Table 3). Thus, functional groups did not show different biomass distribution response to  $\text{CO}_2$  or N, but species did. Six species, representing all four functional groups, had lower root fraction under high N (Table 5). *Petalostemum* had a root fraction of 0.84 under low N and 0.54 under high N, while another legume species (*Lespedeza*) had greater root fraction under high N (0.81) than low N (0.72). Four species, one from each functional group, had higher root fraction under elevated  $\text{CO}_2$  (Table 5). *Petalostemum* was again the most plastic in root fraction, showing root fraction of 0.55 in ambient  $\text{CO}_2$  and 0.82 under elevated  $\text{CO}_2$ . Again, *Lespedeza* showed an opposite response to most other species, having lower root fraction under elevated than ambient  $\text{CO}_2$ . Thus, species were quite varied in the magnitude of biomass distribution responses, with no relation to functional group membership, and in fact the two species with the most pronounced and opposite behaviour were both legumes.

#### Total plant N and tissue N concentration

Averaged across all functional groups or species, the total plant N pool was significantly higher with N addition and unchanged with elevated  $\text{CO}_2$  (Tables 2, 3). Thus, despite greater root biomass under elevated  $\text{CO}_2$ , total cumulative N uptake did not increase, and hence there was substantial dilution of N in biomass (see below). Species differed markedly ( $P < 0.001$ ) in how plant N pools responded to  $\text{CO}_2$  treatments, however (Table 3, Fig. 3). *Lespedeza* and *Lupinus* had large increases in total plant N (of roughly 40–60%, +3.3–4.0 g N  $\text{m}^{-2}$ ) with increased  $\text{CO}_2$ , whereas most other species had similar total N in the elevated vs ambient  $\text{CO}_2$  treatment. Response of total plant N pools to N treatment differed ( $P < 0.05$ ) by functional groups; with forbs showing minimal increase, whereas both

	Low N		High N	
	Ambient $\text{CO}_2$	Elevated $\text{CO}_2$	Ambient $\text{CO}_2$	Elevated $\text{CO}_2$
$\text{C}_3$ grass spp.	11.5	12.6	12.6	21.8
$\text{C}_4$ grass spp.	25.7	24.4	12.0	20.1
Forb spp.	85.3	70.4	76.4	83.7
Legume spp.	43.3	70.3	41.0	62.8
All species	53.8	51.4	55.1	58.5
Functional groups	39.9	36.8	52.7	46.7


**Table 4** Coefficient of variation of total biomass per plot among species within functional groups, among all species, and among functional groups, for four treatment combinations of  $\text{CO}_2$  and N. Mean values per species or functional group based on averages of all plots in all harvests



**Table 5** Means of root fraction (root biomass/total biomass) of species under contrasting CO<sub>2</sub> and N treatments for 128 plots

Functional group	Species	Root fraction (root dm/total plant dm)			
		Ambient CO <sub>2</sub>	Elevated CO <sub>2</sub>	Ambient N	Enriched N
C <sub>4</sub> grass	<i>Andropogon gerardii</i>	0.69	0.65	0.67	0.67
	<i>Bouteloua gracilis</i>	0.65	0.74	0.69	0.69
	<i>Schizachyrium scoparius</i>	0.68	0.69	0.70	0.68
	<i>Sorghastrum nutans</i>	0.69	0.69	0.75	0.63
	Mean	0.68	0.69	0.70	0.67
C <sub>3</sub> grass	<i>Agropyron repens</i>	0.73	0.79	0.76	0.75
	<i>Bromus inermis</i>	0.74	0.73	0.76	0.71
	<i>Koeleria cristata</i>	0.72	0.76	0.77	0.71
	<i>Poa pratensis</i>	0.86	0.84	0.89	0.82
	Mean	0.76	0.78	0.80	0.75
Forb	<i>Achillea millefolium</i>	0.71	0.77	0.77	0.72
	<i>Anemone cylindrica</i>	0.84	0.85	0.83	0.86
	<i>Asclepias tuberosa</i>	0.91	0.93	0.91	0.92
	<i>Solidago rigida</i>	0.64	0.64	0.68	0.61
	Mean	0.77	0.80	0.80	0.78
Legume	<i>Amorpha canescens</i>	0.81	0.85	0.82	0.83
	<i>Lespedeza capitata</i>	0.80	0.73	0.72	0.81
	<i>Lupinus perennis</i>	0.43	0.43	0.43	0.43
	<i>Petalostemum villosum</i>	0.55	0.82	0.84	0.54
	Mean	0.65	0.71	0.70	0.65

Values shown are for 4 plots per species at each CO<sub>2</sub> or N level, pooled across treatments otherwise and averaged over four harvests in two years. The mean standard errors of the adjusted LSMs were 0.04 for species and 0.06 for functional groups.



**Fig. 3** Means of total plant N pools (g m<sup>-2</sup>) of 16 species under contrasting CO<sub>2</sub> and N treatments. Top panel: grey columns, elevated CO<sub>2</sub>; black columns, ambient CO<sub>2</sub>. Bottom panel: grey columns, low N; white columns, high N. The mean standard errors of the adjusted LSMs were 0.8 g m<sup>-2</sup> for species. See Tables 2 and 3 for statistics.

grass groups and legumes showed substantial increase in total N (Table 2; Figs 2, 3).

Functional groups differed in tissue percentage N (Tables 2, 6). Root and shoot percentage N responded to treatments differently, in that roots had less pronounced treatment effects and there were fewer interactions between functional groups or species and treatments (Tables 2, 3, 6). For aboveground tissue, CO<sub>2</sub> treatment reduced percentage N by 11% on

average and the high N treatment increased percentage N by 8% on average (pooled across all other sources of variation). Both functional groups and species differed significantly (Tables 2, 3) in response of aboveground percentage N to both treatments. The C<sub>4</sub> grasses did not have lower percentage N under elevated CO<sub>2</sub> (Table 6). All C<sub>3</sub> functional groups tended to have substantially lower tissue percentage N under elevated CO<sub>2</sub> (Table 6), but species varied markedly in this respect.

**Table 6** Means of tissue percentage N of species under contrasting CO<sub>2</sub> and N treatments for 128 plots. Values shown are for 4 plots per species at each CO<sub>2</sub> or N level, pooled across treatments otherwise and averaged over August harvests in 1998 and 1999

Functional group	Species	Aboveground % N				Belowground % N				Total % N			
		Amb CO <sub>2</sub>	Elev CO <sub>2</sub>	Low N	High N	Amb CO <sub>2</sub>	Elev CO <sub>2</sub>	Low N	High N	Amb CO <sub>2</sub>	Elev CO <sub>2</sub>	Low N	High N
C <sub>4</sub> grass	<i>Andropogon gerardi</i>	1.09	1.15	0.93	1.31	0.95	1.00	0.88	1.07	0.98	1.05	0.90	1.13
	<i>Bouteloua gracilis</i>	1.41	1.33	1.11	1.63	1.14	1.07	0.94	1.26	1.24	1.16	1.01	1.40
	<i>Schizachyrium scoparium</i>	1.01	1.20	1.04	1.17	0.89	0.90	0.82	0.98	0.94	1.01	0.91	1.05
	<i>Sorghastrum nutans</i>	0.91	0.89	0.80	1.00	0.88	0.87	0.82	0.94	0.89	0.89	0.81	0.97
	Mean	1.10	1.13	0.95	1.28	0.97	0.96	0.87	1.06	1.01	1.02	0.90	1.13
C <sub>3</sub> grass	<i>Agropyron repens</i>	1.26	0.89	0.96	1.19	0.84	0.72	0.72	0.84	0.94	0.73	0.75	0.92
	<i>Bromus inermis</i>	0.91	1.00	0.80	1.10	0.91	0.85	0.87	0.89	0.91	0.89	0.85	0.94
	<i>Koeleria cristata</i>	1.43	1.30	1.20	1.53	0.95	1.03	1.01	0.97	1.12	1.12	1.06	1.18
	<i>Poa pratensis</i>	1.42	1.10	1.17	1.37	0.95	0.86	0.89	0.91	1.03	0.91	0.93	1.01
	Mean	1.27	1.08	1.05	1.30	0.90	0.86	0.87	0.90	1.00	0.92	0.90	1.02
Forb	<i>Achillea millefolium</i>	1.47	1.27	1.35	1.39	0.85	0.71	0.78	0.78	1.06	0.87	0.93	1.00
	<i>Anemone cylindrica</i>	1.89	1.55	1.57	1.88	1.37	1.20	1.19	1.38	1.49	1.30	1.27	1.51
	<i>Asclepias tuberosa</i>	3.44	2.94	3.51	2.86	1.45	1.42	1.49	1.38	1.90	1.65	1.77	1.78
	<i>Solidago rigida</i>	1.07	0.80	0.80	1.08	1.09	0.84	0.79	1.14	1.10	0.79	0.80	1.10
	Mean	1.95	1.65	1.80	1.80	1.20	1.04	1.06	1.18	1.39	1.15	1.19	1.35
Legume	<i>Amorpha canescens</i>	2.70	2.17	2.47	2.40	1.48	1.44	1.40	1.53	1.73	1.55	1.58	1.71
	<i>Lespedeza capitata</i>	2.35	1.95	2.12	2.18	1.89	1.92	1.98	1.83	2.00	1.93	2.02	1.91
	<i>Lupinus perennis</i>	1.52	1.43	1.43	1.52	2.15	2.12	2.29	1.99	1.87	1.78	1.88	1.78
	<i>Petalostemum villosum</i>	2.26	2.25	2.46	2.05	1.35	1.33	1.33	1.35	1.72	1.51	1.61	1.62
	Mean	2.21	1.95	2.12	2.04	1.71	1.70	1.75	1.67	1.83	1.69	1.77	1.75

The mean standard error of the adjusted LSMs was 0.10, 0.09, and 0.08% for aboveground, belowground and total percentage N for species and 0.14, 0.07, and 0.07% for aboveground, belowground and total percentage N for functional groups.

*Agropyron*, *Amorpha*, *Anemone*, *Asclepias*, *Lespedeza*, and *Poa* (equally representing the three C<sub>3</sub> groups) decreased percentage N by the greatest amounts under elevated CO<sub>2</sub>, and most other C<sub>3</sub> species had smaller or negligible decreases. Except for the legumes and *Asclepias*, all other species had modest or marked increases in aboveground percentage N with N addition (Table 6).

### Soil water

On average, plots had lower percentage soil water (% SW) in the high N than low N treatments, as hypothesized (H<sub>7</sub>) (Tables 2, 3, 7; Fig. 2). Moreover, species and functional groups differed marginally in percentage SW and in how percentage SW changed with CO<sub>2</sub> and N. C<sub>3</sub> grasses had the lowest percentage SW, C<sub>4</sub> grass and forbs intermediate and legumes the highest percentage SW (Table 7), likely reflecting the influence of differential root biomass on water uptake. To sum, groups are different in percentage SW, but apparently mostly because they have different fine root biomass. Among

species, however, differences in percentage SW do not neatly follow differences in total or root biomass—*Koeleria* (a C<sub>3</sub> grass) had the lowest percentage SW and *Lupinus*, *Agropyron* and several other species the highest.

On average, C<sub>3</sub> grasses showed the greatest increase in percentage SW with elevated CO<sub>2</sub> and the greatest decrease in percentage SW at high N (Table 7; Fig. 2). Except for the legumes, all species had slightly or substantially higher percentage SW under elevated than ambient CO<sub>2</sub> (Table 7), even when root and total biomass were higher in the latter treatment (e.g. *Achillea*, *Agropyron*, *Anemone*, *Bouteloua*, *Poa*) (supporting H<sub>7b</sub>). *Bromus* had the most pronounced ( $P = 0.05$ ) increase in percentage SW under elevated vs ambient CO<sub>2</sub>, consistent with its slightly lower biomass under elevated CO<sub>2</sub>. For N treatments, *Agropyron* and *Koeleria* (C<sub>3</sub> grasses) both had substantially greater depression of percentage SW (both  $P < 0.05$ ) in high N than other species.

Even accounting for variation in fine root biomass, there were significant differences in percentage SW due to CO<sub>2</sub> and N treatments. When data for all plots were pooled and

**Table 7** Means of percentage soil water and soil solution N concentrations for species under contrasting CO<sub>2</sub> and N treatments for 128 plots. Values shown are for 4 plots per species at each CO<sub>2</sub> or N level, pooled across treatments otherwise and averaged over multiple samplings for soil water, and two harvests for soil N, in each of 1998 and 1999. The mean standard error of the adjusted LSMs for percentage soil water was 0.03% for species and 0.2% for functional groups, and 0.1 mg kg<sup>-1</sup> for soil N for both species and functional groups

Functional group	Species	% soil water				Soil solution N conc. (mg/kg)			
		Amb CO <sub>2</sub>	Elev CO <sub>2</sub>	Low N	High N	Amb CO <sub>2</sub>	Elev CO <sub>2</sub>	Low N	High N
C <sub>4</sub> grass	<i>Andropogon gerardii</i>	7.4	7.5	7.7	7.2	0.42	0.38	0.33	0.47
	<i>Bouteloua gracilis</i>	6.5	7.3	7.1	6.8	0.55	0.45	0.39	0.61
	<i>Schizachyrium scoparium</i>	7.6	7.9	7.8	7.6	0.38	0.41	0.31	0.48
	<i>Sorghastrum nutans</i>	7.4	7.5	7.5	7.3	0.20	0.29	0.25	0.25
	Mean	7.2	7.6	7.6	7.2	0.39	0.38	0.33	0.44
C <sub>3</sub> grass	<i>Agropyron repens</i>	7.8	8.4	8.9	7.2	0.33	0.23	0.27	0.29
	<i>Bromus inermis</i>	6.8	8.1	7.6	7.3	0.29	0.33	0.31	0.31
	<i>Koeleria cristata</i>	6.1	6.3	6.7	5.6	0.22	0.19	0.26	0.14
	<i>Poa pratensis</i>	6.4	6.9	6.9	6.4	0.38	0.27	0.35	0.30
	Mean	6.8	7.4	7.5	6.6	0.29	0.25	0.29	0.26
Forb	<i>Achillea millefolium</i>	7.0	7.3	7.3	7.1	0.19	0.26	0.19	0.26
	<i>Anemone cylindrica</i>	7.6	8.0	7.4	8.2	0.75	0.71	0.37	1.09
	<i>Asclepias tuberosa</i>	7.7	7.8	7.8	7.8	1.84	1.04	1.27	1.61
	<i>Solidago rigida</i>	6.8	7.2	7.1	7.0	0.23	0.25	0.16	0.32
	Mean	7.3	7.6	7.4	7.5	0.77	0.56	0.51	0.82
Legume	<i>Amorpha canescens</i>	8.3	8.0	8.5	7.8	1.39	1.18	0.99	1.58
	<i>Lespedeza capitata</i>	7.8	7.6	8.0	7.4	1.51	0.94	1.11	1.34
	<i>Lupinus perennis</i>	8.7	8.6	8.7	8.5	1.67	2.11	1.57	2.21
	<i>Petalostemum villosum</i>	7.8	8.0	8.1	7.6	1.71	0.80	1.04	1.47
	Mean	8.1	8.1	8.4	7.9	1.57	1.26	1.18	1.65

percentage SW was plotted against fine root biomass, there was significantly higher percentage SW under elevated than ambient CO<sub>2</sub> and lower percentage SW under elevated than ambient N (Fig. 4a,b).


### Soil solution N

The high N treatment increased soil solution nitrate and total N by roughly 40–50% on average (Table 7; Fig. 2). In contrast, and supporting H<sub>7a</sub>, elevated CO<sub>2</sub> reduced soil nitrate and total N by roughly 20–25% (Table 7; Fig. 2), consistent with increased fine root biomass under elevated CO<sub>2</sub>. Functional groups differed in soil solution N, with legumes having the highest and C<sub>3</sub> grasses the lowest levels (Tables 2, 7). These functional group differences were roughly opposite to patterns of fine root biomass. Functional groups differed in the extent of depletion of soil solution N due to CO<sub>2</sub>, with the C<sub>4</sub> group having little effect compared to the C<sub>3</sub> groups in general (Tables 5, 7). For N treatment, legumes had a significantly greater increase in soil solution N than the C<sub>3</sub> grass species,


likely due to the degree of soil solution N depletion associated with differences in fine root biomass.

Accounting for variation in fine root biomass, there were significant differences in soil N concentrations due to N treatments, but not to CO<sub>2</sub>. When data for all plots were pooled and soil solution N was plotted against fine root biomass, plots had significantly higher soil N under high than low N but did not differ under the contrasting CO<sub>2</sub> treatments (Fig. 4c,d).

For nonlegumes, there was an inverse relationship between total plant N pools and soil solution N concentrations ( $P < 0.001$ ), likely because vegetation on plots with higher fine root biomass take up N (which is incorporated into plant tissues) while driving down the soil solution N concentration (Fig. 5). At any soil solution N pool, the N deposition treatment had higher plant N pools. Two of the legumes (*Petalostemum* and *Amorpha*) had slightly higher soil solution N pools at any given plant N pool but fit more or less within the general scatterplot relationship for the non-legumes ( $P < 0.001$ ,  $r^2 = 0.44$  for the 56 plots within each N treatment level) (Fig. 5). By contrast, the other two legumes, *Lespedeza*



**Fig. 4** (a,b) Mean percentage volumetric soil water (averaged over 18 sampling periods over two years) in relation to mean fine root biomass (averaged over 4 harvests in 2 yr) for 16 species monocultures under contrasting  $\text{CO}_2$  and N treatments (pooled across the other treatment). Open circles, ambient  $\text{CO}_2$ ; closed circles, elevated  $\text{CO}_2$ ; open triangles, low N; closed triangles, high N. The slopes were not significantly different among treatments in either case, but the elevation of the line was significantly ( $P < 0.025$ ) different in each case. (c,d) Mean soil solution N concentration ( $\text{mg}/\text{kg}$ ) in relation to mean fine root biomass (both averaged over 4 harvests in 2 yr) for 16 species monocultures under contrasting  $\text{CO}_2$  and N treatments (pooled across the other treatment). Open circles, ambient  $\text{CO}_2$ ; closed circles, elevated  $\text{CO}_2$ ; open triangles, low N; closed triangles, high N. The slopes were not significantly different among treatments in either case, and the elevation of the line was significantly ( $P < 0.025$ ) different for contrasting N treatments but not contrasting  $\text{CO}_2$  treatments.



**Fig. 5** Mean soil solution N concentration ( $\text{mg}/\text{kg}$ ) in relation to mean total plant N ( $\text{g}/\text{m}^2$ ), for plots under low and high N treatments, pooled across  $\text{CO}_2$  treatments. The curves shown are for all nonN-fixers plus *Amorpha* and *Petalostemum*, and the relationship between  $\log_{10}$  soil N and  $\log_{10}$  plant N was significant ( $P < 0.001$ ,  $R^2 = 0.44$ ) in both cases. Closed circles, 14 species; open circles, *Lepedeza* and *Lupinus*.

and *Lupinus*, always had higher soil solution N pools at any given plant N pool and plots for these two species fell far from the relationship for the other 14 species (Fig. 5).

## Discussion

A surprising number of species in this study had modest or negligible increases in biomass production in response to  $\text{CO}_2$  or N fertilization (Fig. 1). Our major hypotheses regarding functional group responses to  $\text{CO}_2$  and N were neither consistently supported nor rejected, and will be discussed below. Overall, these grassland species were less responsive to elevated  $\text{CO}_2$  (see also Lee *et al.*, 2001) than has been generally found elsewhere, although some studies have found weak responses to elevated  $\text{CO}_2$  (Koch & Mooney, 1996; Körner & Bazzaz, 1996; Curtis & Wang, 1998; Poorter, 1998). The mean increase due to elevated  $\text{CO}_2$  in total biomass of all 12  $\text{C}_3$  species in our study was 16%, considerably less than

the 29%, 42% and 44% increases, respectively, reported in reviews of  $\text{C}_3$  woody plants (Curtis & Wang, 1998),  $\text{C}_3$  grasses in general (Poorter, 1993), and  $\text{C}_3$  Poaceae (Wand *et al.*, 1999). The mean response of biomass to elevated  $\text{CO}_2$  for the  $\text{C}_4$  grasses in our study ( $-3\%$ ) is also less than the mean increases of 22% and 33% reported for  $\text{C}_4$  grasses in general (Poorter, 1993) and for  $\text{C}_4$  Poaceae (Wand *et al.*, 1999), respectively. Possible explanations for the limited  $\text{CO}_2$  response in our study compared with previous studies include the relatively low fertility of our site, the general high level of adaptation to infertility among our study species, and the moist conditions during the study years. Alternatively, only a small fraction of the studies used in the reviews cited above were grown under realistic field conditions, and even fewer without chambers. It is important to note that a high degree of photosynthetic acclimation among these species (Lee *et al.*, 2001) can broadly explain the small growth response to elevated  $\text{CO}_2$  at our site, though the correlation between individual species photosynthetic

vs. biomass responses was weak (P. B. Reich *et al.*, unpublished). Testing whether long-term responses to elevated CO<sub>2</sub> of grassland species in the field are generally less than those of shorter-term and more controlled experiments will require a larger data base than is currently available.

### Functional group and species responses to elevated CO<sub>2</sub>

As a group, C<sub>4</sub> grasses did show significantly less biomass enhancement in response to elevated CO<sub>2</sub> than the C<sub>3</sub> functional groups (supporting H<sub>1</sub>) and no C<sub>4</sub> grass species showed a significant increase. These biomass responses are consistent with leaf level photosynthesis data; during the 2-yr study period, the C<sub>4</sub> grass species showed no enhancement of photosynthesis under elevated CO<sub>2</sub>, in contrast to a modest increase on average for the C<sub>3</sub> species (Lee *et al.*, 2001).

Two of the four legumes in our study, *Lupinus* and *Lespedeza*, showed marked production increases under elevated CO<sub>2</sub> but the other two did not. Thus (H<sub>2</sub>) – a hypothesis based on the general tendency of N-fixers to respond more to elevated CO<sub>2</sub> than nonfixers (Soussana & Hartwig, 1996; Clark *et al.*, 1997; Hebeisen *et al.*, 1997; Lüscher *et al.*, 1997, 1998; Schenk *et al.*, 1997) – apparently did not hold. Can we explain this marked difference in response among legume species?

Although we know little about variation in the degree of N-fixation among these four legumes, we can make some indirect inferences based on the data from this study. *Lupinus* and *Lespedeza* plots had somewhat higher soil solution N pools and two to three times as much plant N per plot as *Amorpha* and *Petalostemum*, which we take as evidence of greater N fixation in the former two species, since all else being equal, with greater root biomass they should have otherwise depleted the soil solution N to a level lower than the other two legumes. Moreover, for all nonfixers plus *Amorpha* and *Petalostemum* pooled, there was a significant inverse relationship (Fig. 5) between soil solution N and plant N pools, consistent with earlier studies relating differences among species in root biomass to depletion of soil solution N pools in grasslands at Cedar Creek (Tilman & Wedin, 1991). *Lupinus* and *Lespedeza* did not follow this inverse relationship and had much higher soil solution N for a given plant N pool, suggesting relatively high rates of N fixation.

In addition, for the 12 nonfixer species (plus *Amorpha* and *Petalostemum*) the plant N pool was relatively unchanged under elevated CO<sub>2</sub>. This suggests that increased biomass in response to elevated CO<sub>2</sub> by itself did not result in greater total plant N acquisition in nonfixers. In contrast, *Lupinus* and *Lespedeza* had marked increases in plant N pool in elevated CO<sub>2</sub> (4.0 and 3.3 g m<sup>-2</sup>, respectively). Hence, it is plausible that *Lupinus* and *Lespedeza* more vigorously fix N than the other legumes, and responded to elevated CO<sub>2</sub> by increasing their own N supply, leading to heightened uptake of both C and N. In support of this idea, using the <sup>15</sup>N isotope dilution method for *Lupinus* monocultures, the proportion of

N derived from fixation increased by 38% under elevated CO<sub>2</sub> (T. D. Lee *et al.*, unpublished). Thus, based on indirect evidence from this study, the legumes with greater N-fixation tendencies were more responsive to elevated CO<sub>2</sub> than those with lesser N-fixation tendencies, supporting the concept of N-fixation enhancing CO<sub>2</sub> responses (H<sub>2</sub>), but suggesting that heterogeneity among N-fixers may limit the generalizability of this idea.

### Functional group and species responses to increased N supply

Our hypotheses about biomass responses to increased N supply (H<sub>3</sub> to H<sub>5</sub>) were also supported with mixed results. The C<sub>3</sub> grasses increased biomass the most at high N and the legumes as a group responded little to N addition, as predicted (H<sub>3</sub>, H<sub>4</sub>). However, one of the four legume species, *Petalostemum*, did increase biomass at high N, and perhaps related, this species did not respond positively to elevated CO<sub>2</sub> and may have had a low rate of N fixation. All eight grasses, regardless of photosynthetic pathway, had large increases in aboveground biomass at high N, and 7 of the 8 also had large increases belowground. However, surprisingly, only one of four forbs responded positively to N addition in terms of total biomass. Perhaps these species are poor competitors for soil N in relationship to microbes? If that was so (i.e. microbial uptake depleted the available soil N), soil solution N should be low, yet it was higher for the forbs than for the grasses. Moreover, two of the three forbs which did not increase biomass at high N did have higher tissue percentage N. The failure of these species to respond positively with increased production under high N is difficult to explain.

Finally, there were no CO<sub>2</sub> × N interactions and hence no tendency for any functional group to respond differently to CO<sub>2</sub> as a function of N supply (H<sub>5</sub>). Elevated N supply did not increase monoculture response to CO<sub>2</sub>, unlike a number of other studies (e.g. Owensby *et al.*, 1994; Zanetti *et al.*, 1996; Curtis & Wang, 1998; Zak *et al.*, 2000). Whether this is due to the relatively smaller N addition in this study (4 g N m<sup>-2</sup> yr<sup>-1</sup>) than in some others (e.g. N treatments were as high as 56 g N m<sup>-2</sup> yr<sup>-1</sup> and more closely mimic agricultural N addition rates, Zanetti *et al.*, 1996) can not be answered without a larger number of studies of elevated CO<sub>2</sub> effects under different N addition regimes.

### Controls on percentage soil water

How can we explain CO<sub>2</sub> and N treatment effects on percentage soil water? Biomass per plot had a large influence on percentage SW (Fig. 4), and both species and treatments contributed to variation in biomass, although the former was dominant. For N, even for a given fine root mass, plots under high N had lower percentage SW. There was no effect of N on leaf diffusive conductance (Lee *et al.*, 2001) (and hence on leaf-level water loss), but high N did lead to a lower root


fraction (Tables 3, 5). Hence, high N plots support a higher aboveground biomass and likely a higher leaf area index and transpirational surface, perhaps explaining the lower percentage SW under high N at a given fine root biomass.

By contrast, elevated CO<sub>2</sub> leads to greater biomass which, all else being equal, should lead to greater depletion of soil water. However, individual species actually had lower percentage SW under elevated CO<sub>2</sub> despite greater biomass (Table 7) and overall, percentage SW at any given fine root biomass was higher under elevated CO<sub>2</sub>. Elevated CO<sub>2</sub> did not affect root fraction, so a shift in absorbing (roots) vs transpirational surfaces (canopy) can not be invoked as an explanatory factor. In a companion study (Lee *et al.*, 2001) we found a consistent and roughly 25% decrease in leaf conductance across species, functional groups and years. An analysis (data not shown) that includes functional group and leaf conductance found both factors significantly ( $P < 0.05$ ) associated with percentage SW, with percentage SW negatively related to conductance. Hence, reduced conductance contributes to the tendency for elevated CO<sub>2</sub> plots to have lower percentage SW.

#### Other functional traits and groupings

Variation in a number of other functional traits, including intrinsic growth rate, leaf gas exchange capacity, resource depletion capacity, sink strength, plant strategy, and root symbiont status, has been proposed to explain differential responsiveness to elevated CO<sub>2</sub>, added N and a variety of other global change factors (Reich, 1987; Hunt *et al.*, 1991, 1993; Díaz, 1995; Poorter *et al.*, 1996; Wedin & Tilman, 1996). Such explanations have met with some success with other factors, but less so vis-à-vis CO<sub>2</sub> (Poorter *et al.*, 1996; Volin & Reich, 1996). None of these hypotheses moreover, appear capable of fully explaining differences in responses among the 16 species in this study. The species that were most productive under ambient CO<sub>2</sub> and soil N conditions were not consistently responsive to either elevated CO<sub>2</sub> or N, nor were the less productive species consistently responsive. Similarly, for 10 of the 16 species we also measured relative growth rate of seedlings under controlled conditions (P. B. Reich *et al.*, unpublished) and found no relationship between proportional biomass enhancement due to CO<sub>2</sub> in field plots and individual seedling RGR. Species which reduce the soil N concentration under ambient conditions did generally respond more positively to N enrichment, but with two important exceptions (*Achillea* and *Solidago*).

Among the nonlegumes, tissue percentage N in the low N treatment did not predict which species would respond to N enrichment (data not shown), but it did predict the relative responsiveness to elevated CO<sub>2</sub> (Fig. 6). For C<sub>3</sub> and C<sub>4</sub> species separately, species with greater aboveground tissue percentage N had a greater increase in biomass in response to elevated CO<sub>2</sub> than those with lower percentage N. The slopes were similar for both groups, and the relationship was at a lower



**Fig. 6** The percentage enhancement of total biomass due to elevated CO<sub>2</sub> for all nonN-fixer species, in relation to the aboveground tissue percentage N of plants under ambient CO<sub>2</sub> conditions. Open circles, C<sub>4</sub> species; closed circles, C<sub>3</sub> species. The regression relationships were significant ( $P < 0.001$ ) for C<sub>3</sub> and C<sub>4</sub> species considered separately ( $R^2 = 0.72$  and  $0.99$ , respectively), with a similar slope ( $P > 0.10$ ) but different intercept ( $P < 0.05$ ).

‘elevation’ (i.e. intercept was lower) for the C<sub>4</sub> than C<sub>3</sub> species. Why do these patterns occur? A plausible hypothesis follows from physiological principles. Species with higher tissue percentage N have a higher carboxylation capacity (Evans, 1989) such that if they were ‘operating’ on the steep part of the A-C<sub>i</sub> curve (Farquhar & Sharkey, 1982) they should have a greater enhancement due to elevated CO<sub>2</sub> than species with lower tissue percentage N (this could explain patterns within each group). Moreover, since C<sub>4</sub> plants are already near saturated at ambient CO<sub>2</sub> (i.e. on a shallower part of the curve) they are less responsive as a group to elevated CO<sub>2</sub>, even for a given leaf percentage N (this could explain the lower position of the overall line for this group). With their different C : N dynamics, legumes would not necessarily be expected to follow similar patterns. Although many processes beyond leaf level photosynthesis influence ecosystem scale biomass enhancement to elevated CO<sub>2</sub> (such as canopy architecture, turnover rates, tissue morphology, biomass distribution, and phenology), leaf-level processes provide the starting point for carbon acquisition which leads to biomass accumulation, and which drives increased biomass accumulation under elevated CO<sub>2</sub>. Whether this is a robust general relationship requires further testing in other common garden experiments in the field.

#### Implications and conclusions

There have been many isolated potted plant studies of elevated CO<sub>2</sub> responses. Studies of monocultures in the field with free-air CO<sub>2</sub> enrichment provide information that is closer to natural conditions. Moreover, if we hope to be able to use functional groupings to generalize about, and quantitatively model responses

to elevated CO<sub>2</sub> and N of mixed species communities, then we need to understand variation in species and functional group responses under simpler monoculture conditions. If responses to elevated CO<sub>2</sub> and N deposition are largely a function of changing resource supply then growth in mixed communities is complicated by species interactions that can change as a result of these agents (Owensby *et al.*, 1993, 1994). Competition often involves growing with other species that differentially utilize resources (e.g. soil solution N, Tilman & Wedin, 1991) or supply them (e.g. N-fixers) by dint of species ecophysiological differences. Hence, understanding species responses to enriched CO<sub>2</sub> or N in mixed communities is complicated because competition ensures that any given species does not necessarily have access to increased supplies of a resource that is added to an ecosystem. Nonetheless, responses of species mixtures to elevated CO<sub>2</sub> may be related to their responses in monocultures (Navas *et al.*, 1999). Therefore, interpretation and evaluation of species under interspecific competition (e.g. Warwick *et al.*, 1998; Leadley *et al.*, 1999) will be aided by a better understanding of their responses growing under intraspecific competition, as in the monocultures of this study.

Our results are only somewhat encouraging vis-à-vis the use of functional groups. Functional groups did often respond significantly differently to CO<sub>2</sub> or N, and hence do provide some meaningful information without knowledge about individual species. However, there was also substantial variation in response among species within groups. Alternative classifications based on measured continuous traits were generally no more useful than the *a priori* defined functional groups. These results suggest that current trait-based functional classifications may be useful, but not sufficient for understanding plant and ecosystem responses to elevated CO<sub>2</sub> and N deposition.

## Acknowledgements

We acknowledge funding from the U.S. Department of Energy and the National Science Foundation Cedar Creek Natural History Area, Long-Term Ecological Research site in support of this research. We thank G. Hendrey, K. Lewin, C. Lehman, K. Wrage, S. Jose and many undergraduate interns whose work contributed to myriad phases of this research.

## References

- Baker JM. 1990. Measurement of soil water content. *Remote Sensing Review* 5: 263–279.
- Cannell M, Thornley J. 1998. N-poor ecosystems may respond more to elevated [CO<sub>2</sub>] than N-rich ones in the long term. A model analysis of grassland. *Global Change Biology* 4: 431–442.
- Clark H, Newton PCD, Bell CC, Glasgow EM. 1997. Dry matter yield, leaf growth and population dynamics in *Lolium perenne*/*Trifolium repens*-dominated pasture turves exposed to two levels of elevated CO<sub>2</sub>. *Journal of Applied Ecology* 34: 304–316.
- Curtis PS, Drake BG, Leadley PW, Arp WJ, Whigham DF. 1989. Growth and senescence in plant communities exposed to elevated CO<sub>2</sub> concentrations on an estuarine marsh. *Oecologia* 78: 20–26.
- Curtis PS, Wang X. 1998. A meta-analysis of elevated CO<sub>2</sub> effects on woody plant mass, form, and physiology. *Oecologia* 113: 299–313.
- Díaz S. 1995. Elevated CO<sub>2</sub> responsiveness, interactions at the community level, and plant functional types. *Journal of Biogeography* 22: 289–295.
- Evans JR. 1989. Photosynthesis and nitrogen relationships in leaves of C<sub>3</sub> plants. *Oecologia* 78: 9–19.
- Farquhar GD, Sharkey TD. 1982. Stomatal conductance and photosynthesis. *Annual Review of Plant Physiology* 33: 317–345.
- Finn GA, Brun WA. 1982. Effect of atmospheric CO<sub>2</sub> enrichment on growth, nonstructural carbohydrate content, & root nodule activity in soybean. *Plant Physiology* 69: 327–331.
- Hebeisen T, Luscher A, Zanetti S, Fischer BU, Hartwig UA, Frehner M, Hendrey GR, Blum H, Nösberger J. 1997. Growth response of *Trifolium repens* L. and *Lolium perenne* L. as monocultures and bi-species mixture to free air CO<sub>2</sub> enrichment and management. *Global Change Biology* 3: 149–160.
- Hunt R, Hand DW, Hannah MA, Neal AM. 1991. Response to CO<sub>2</sub> enrichment in 27 herbaceous species. *Functional Ecology* 5: 410–421.
- Hunt R, Hand DW, Hannah MA, Neal AM. 1993. Further response to CO<sub>2</sub> enrichment in British herbaceous species. *Functional Ecology* 7: 661–668.
- Koch GW, Mooney HA. 1996. *Carbon dioxide and terrestrial ecosystem*. San Diego, CA, USA: Academic Press.
- Körner C, Bazzaz FA. 1996. *Carbon dioxide, populations and communities*. San Diego, CA, USA: Academic Press.
- Larigauderie A, Hilbert DW, Oechel WC. 1988. Effect of CO<sub>2</sub> enrichment and nitrogen availability on resource acquisition and resource allocation in a grass, *Bromus mollis*. *Oecologia* 77: 544–549.
- Leadley PW, Körner C. 1996. Effects of elevated CO<sub>2</sub> on plant species dominance in a highly diverse calcareous grassland. In: Körner C, Bazzaz FA, eds. *Carbon dioxide, populations, and communities*. San Diego, CA, USA: Academic Press, Inc., 159–175.
- Leadley PW, Niklaus PA, Stocker R, Körner C. 1999. A field study of the effects of elevated CO<sub>2</sub> on plant biomass and community structure in a calcareous grassland. *Oecologia* 118: 39–49.
- Lee TD, Tjoelker MG, Ellsworth DS, Reich PB. 2001. Leaf gas exchange responses of 13 prairie grassland species in the field under elevated carbon dioxide and increased nitrogen supply. *New Phytologist* 150: 000–000.
- Lewin KF, Hendrey GR, Nagy J, LaMorte R. 1994. Design and application of a free-air carbon dioxide enrichment facility. *Agricultural and Forest Meteorology* 70: 15–29.
- Lloyd J, Farquhar GD. 1996. The CO<sub>2</sub> dependence of photosynthesis, plant growth responses to elevated atmospheric CO<sub>2</sub> concentrations and their interaction with soil nutrient status. I. General principals and forest ecosystems. *Functional Ecology* 10: 4–32.
- Lüscher A, Hendrey GR, Nösberger J. 1998. Long-term responsiveness to free air CO<sub>2</sub> enrichment of functional types, species and genotypes of plants from fertile permanent grassland. *Oecologia* 113: 37–45.
- Lüscher A, Nösberger J. 1997. Interspecific and intraspecific variability in the response of grasses and legumes to free air CO<sub>2</sub> enrichment. *Acta Oecologica* 18: 269–275.
- Navas M-L, Garnier E, Austin MP, Gifford RM. 1999. Effect of competition on the responses of grasses and legumes to elevated atmospheric CO<sub>2</sub> along a nitrogen gradient: differences between isolated plants, monocultures and multi-species mixtures. *New Phytologist* 143: 323–331.
- Owensby CE, Auen LM, Coyne PI. 1994. Biomass production in a nitrogen-fertilized, tallgrass prairie ecosystem exposed to ambient and elevated levels of CO<sub>2</sub>. *Plant and Soil* 165: 105–113.
- Owensby CE, Coyne PI, Ham JM, Auen LM, Knapp AK. 1993. Biomass production in a tallgrass prairie ecosystem exposed to ambient and elevated CO<sub>2</sub>. *Ecological Applications* 3: 644–653.

- Pearcy RW, Ehleringer J. 1984. Comparative ecophysiology of C<sub>3</sub> and C<sub>4</sub> plants. *Plant, Cell & Environment* 7: 1–13.
- Poorter H. 1993. Interspecific variation in the growth response of plants to an elevated ambient CO<sub>2</sub> concentration. *Vegetatio* 104/105: 77–97.
- Poorter H. 1998. Do slow-growing species and nutrient-stressed plants respond relatively strongly to elevated CO<sub>2</sub>. *Global Change Biology* 4: 693–697.
- Poorter H, Nagel O. 2000. The role of biomass allocation in the growth response of plants to different levels of light, CO<sub>2</sub>, nutrients and water: a quantitative review. *Australian Journal of Plant Physiology* 27: 595–607.
- Poorter H, Roumet C, Campbell B. 1996. Interspecific variation in the growth response of plants to elevated CO<sub>2</sub>: a search for functional types. In: Körner C, Bazzaz FA, eds. *Carbon dioxide, populations and communities*. San Diego, CA, USA: Academic Press, 375–412.
- Reich PB. 1987. Quantifying plant response to ozone: a unifying theory. *Tree Physiology* 3: 63–91.
- Reich PB. 2001. Root-shoot relationships: optimality in biomass allocation or the 'Emperor's New Clothes?'. In: Waisel Y, Eschel A, Kafkafi U, eds. *Plant roots: the hidden half*. New York, USA: Marcel Dekker, Inc. (In press.)
- Reich PB, Knops J, Tilman D, Craine J, Ellsworth D, Tjoelker M, Lee T, Naeem S, Wedin D, Bahaeddin D, Hendrey G, Jose S, Wragge K, Goth J, Bengston W. 2001. Plant diversity enhances ecosystem responses to elevated CO<sub>2</sub> and nitrogen deposition. *Nature* (In press.)
- Schenk U, Jager H-J, Weigel H-J. 1997. The response of perennial ryegrass/white clover swards to elevated atmospheric CO<sub>2</sub> concentrations. *New Phytologist* 135: 67–79.
- Soussana JF, Hartwig UA. 1996. The effects of elevated CO<sub>2</sub> on symbiotic N<sub>2</sub> fixation: a link between the carbon and nitrogen cycles in grassland ecosystems. *Plant and Soil* 187: 321–332.
- Tilman D, Wedin D. 1991. Plant traits and resource reduction for five grasses growing on a nitrogen gradient. *Ecology* 72: 685–700.
- Tjoelker MG, Oleksyn J, Lee TD, Reich PB. 2001. Direct inhibition of leaf dark respiration by elevated CO<sub>2</sub> is minor in 12 grassland species. *New Phytologist* 150: 000–000.
- Volin JC, Reich PB. 1996. Interaction of carbon dioxide and ozone on C<sub>3</sub> and C<sub>4</sub> grasses and trees under contrasting nutrient supply. *Physiologia Plantarum* 97: 674–684.
- Ward SJ, Midgley GF, Jones MH, Curtis PS. 1999. Responses of wild C<sub>4</sub> and C<sub>3</sub> (Poaceae) species to elevated atmospheric CO<sub>2</sub> concentration: a meta-analytic test of current theories and perceptions. *Global Change Biology* 5: 723–741.
- Warwick KR, Taylor G, Blum H. 1998. Biomass and compositional changes occur in chalk grassland turves exposed to elevated CO<sub>2</sub> for two seasons in FACE. *Global Change Biology* 4: 375–385.
- Wedin D, Tilman D. 1996. Influence of Nitrogen Loading and Species Composition on the Carbon Balance of Grasslands. *Science* 274: 1720–1723.
- Zak DR, Pregit KS, Curtis PS, Vogel CS, Holmes WE, Lussenhop J. 2000. Atmospheric CO<sub>2</sub>, soil-N availability, and allocation of biomass and nitrogen by *Populus tremuloides*. *Ecological Applications* 10: 34–46.
- Zanetti S, Hartwig UA, Lüscher A, Hebeisen T, Frehner M, Fischer BU, Hendrey GR, Blum H, Nösberger J. 1996. Stimulation of symbiotic N<sub>2</sub> fixation in *Trifolium repens* L. under elevated atmospheric CO<sub>2</sub> in a grassland ecosystem. *Plant Physiology* 112: 575–583.