

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Other Publications in Zoonotics and Wildlife Disease

Wildlife Disease and Zoonotics

1994

SCID mouse spleen does not support scrapie agent replication

Katherine I. O'Rourke

U.S. Department of Agriculture, katherine.orourke@ars.usda.gov

T. P. Huff

U.S. Department of Agriculture

C. W. Leathers

Washington State University

M. M. Robinson

U.S. Department of Agriculture

J. R. Gorham

U.S. Department of Agriculture

Follow this and additional works at: <https://digitalcommons.unl.edu/zoonoticspub>



Part of the [Veterinary Infectious Diseases Commons](#)

O'Rourke, Katherine I.; Huff, T. P.; Leathers, C. W.; Robinson, M. M.; and Gorham, J. R., "SCID mouse spleen does not support scrapie agent replication" (1994). *Other Publications in Zoonotics and Wildlife Disease*. 119.

<https://digitalcommons.unl.edu/zoonoticspub/119>

This Article is brought to you for free and open access by the Wildlife Disease and Zoonotics at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Other Publications in Zoonotics and Wildlife Disease by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

SCID mouse spleen does not support scrapie agent replication

K. I. O'Rourke,^{1*} T. P. Huff,¹ C. W. Leathers,² M. M. Robinson¹ and J. R. Gorham¹

¹ United States Department of Agriculture, Agricultural Research Service, Animal Disease Research Unit, Pullman, Washington 99164-7030 and ² Washington Animal Disease Diagnostic Laboratory, Washington State University, Pullman, Washington 99164-7034, U.S.A.

BALB/c and severe combined immunodeficiency (SCID) mice were inoculated intracerebrally or intraperitoneally with scrapie agent strain ME7 to examine the role of functional lymphocytes and follicular dendritic cells in splenic infectivity and PrP^{Sc} accumulation. Intracerebrally inoculated BALB/c and SCID mice developed the clinical signs and microscopic lesions characteristic

of scrapie. Spleens from terminally affected BALB/c mice contained PrP^{Sc} which was detectable by immunoblot analysis; SCID mouse spleens did not contain detectable PrP^{Sc}. SCID mouse spleens collected during the first 90 days after intraperitoneal infection contained neither infectivity nor PrP^{Sc}.

Sheep scrapie is the archetype of a heterogeneous group of rare transmissible spongiform encephalopathies (TSEs) occurring naturally in humans, cattle, mink, cats and captive deer (Hartsough & Burger, 1965; Williams & Young, 1980; Wells *et al.*, 1987; Leggett *et al.*, 1990; Brown & Gajdusek, 1991). The histopathological lesions of TSEs are restricted to the nervous system (Chandler, 1969). However, immunohistochemical and biochemical analyses detect a host cellular membrane glycoprotein, the prion protein (PrP^C) (Prusiner, 1982; Oesch *et al.*, 1985; Basler *et al.*, 1986), that accumulates in lymphoid and nervous tissues as a relatively protease-resistant isoform [PrP^{Sc} in scrapie-infected animals and PrP^{CJD} in Creutzfeldt-Jakob disease (CJD)-infected animals] (Merz *et al.*, 1987; Doi *et al.*, 1988; Farquhar *et al.*, 1989; Kitamoto *et al.*, 1989; Race *et al.*, 1992). PrP^{Sc} is tightly associated with infectivity and may either contain or be the infectious agent itself (Diringer *et al.*, 1983; McKinley *et al.*, 1983; Merz *et al.*, 1984, 1987). Infectivity and PrP^{Sc} accumulation in the spleen are detectable within 2 weeks of experimental infection in rodents and persist throughout the incubation period (Eklund *et al.*, 1967; Race & Ernst, 1992). Scrapie infectivity in the spleen is associated with post-mitotic non-lymphoid (stromal) cells and is unaffected by thymectomy (McFarlin *et al.*, 1971; Worthington & Clark, 1971; Clarke & Kimberlin, 1984; Robinson & Gorham, 1990). PrP^C in normal mouse spleens and PrP^{Sc}/PrP^{CJD} accumulation in infected mouse spleens is restricted to cells with the morphological characteristics and antigen-trapping capacity of follicular dendritic cells (FDCs) (Kitamoto *et al.*, 1991; McBride *et al.*, 1992). In spite of these similarities, the splenic

replication phase is not obligatory in all rodent TSE models. Prion mRNA is detectable in mouse spleens by Northern blot analysis (Caughey *et al.*, 1988) and PrP^C is detectable by immunoblot and immunofluorescence analysis of hamster leukocytes and spleens (Bendheim *et al.*, 1992).

Severe combined immunodeficiency (SCID) mice, which lack functional FDCs secondary to the absence of lymphocytes (Kapasi *et al.*, 1993), do not develop clinical signs of CJD or accumulation of splenic PrP^{CJD} following intraperitoneal (i.p.) inoculation (Kitamoto *et al.*, 1991), although CJD is readily transmitted to SCID mice by the intracerebral (i.c.) route. To extend this observation, we have examined splenic infectivity and PrP^{Sc} accumulation in SCID mice inoculated with scrapie agent strain ME7.

Homozygous C.B.-17-SCID/SCID female mice, 3 weeks of age, were obtained from Taconic Laboratories and normal 3-week-old female BALB/c mice were obtained from B&K Universal. All mice were housed in groups of two to four animals in microisolator cages and given autoclaved food and water. Scrapie agent strain ME7 was obtained as whole frozen brains from terminally infected C57BL mice. A 10% (w/v) brain suspension was prepared using sterile Tris-buffered saline (TBS; 0.05 M-Tris-HCl pH 7.8, 0.1 M-NaCl) in a sterile glass Dounce homogenizer and frozen at -20 °C until required. Mice were inoculated by the i.c. route with 30 µl or by the i.p. route with 100 µl of freshly thawed 10% brain homogenate or with dilutions made in sterile TBS. Controls included untreated mice and mice inoculated with similar volumes of a 10% homogenate of uninfected C57BL brain or with TBS alone. Mice were

Table 1. Titration of ME7 in BALB/c and SCID mice

Dilution of whole ME7 brain (log ₁₀)	Animals affected/number inoculated [incubation range (days)]	
	BALB/c	SCID
-1	3/3 (146-147)	2/2 (157)
-2	3/3 (161-180)	3/3 (176)
-3	4/4 (161-182)	2/2 (176-179)
-4	4/4 (171-196)	3/3 (184)
-5	4/4 (189-198)	3/3 (198)
-6	3/3 (243)	ND*
-7	4/4 (252)	ND
-8	0/4†	ND

* ND, Not done.

† Animals euthanized at 426 days p.i.

Table 2. Infectivity bioassay of spleens from i.p. inoculated BALB/c and SCID mice

Time p.i. (days)	Animals affected/number inoculated [incubation range (days)]	
	BALB/c donor	SCID donor
30	16/16 (167-196)	0/16
60	15*/15 (164-174)	0/16
90	16/16 (174-201)	0/16

* Excludes one mouse which died within 24 h of inoculation.

observed daily for clinical signs (ruffled fur, lethargy, ataxia) and euthanized either when clearly moribund or 3 weeks after lethargy was first noted. Incubation times are reported as days from inoculation to euthanasia or death. The i.c. infectious titre of the inoculum was 10^9 LD₅₀/ml (Reed & Muench, 1938; Eklund *et al.*, 1967). Inoculation (i.c.) of SCID mice with the five lowest dilutions of the inoculum resulted in incubation periods (Table 1) similar to those observed in BALB/c mice. Histopathological examination of the brains of representative affected BALB/c and SCID mice showed focal spongiosis, astrocytosis and rare intraneuronal vacuolation (not shown).

Infectivity in the spleens of i.p.-inoculated SCID and BALB/c mice was determined at 30, 60 and 90 days post-inoculation (p.i.). Spleens were collected from individual ME7-infected BALB/c or SCID mice and from uninfected control mice. All mice were clinically normal when tissues were collected. Homogenates (10% w/v) were prepared with sterile TBS in disposable tissue homogenizers, clarified at 228 g for 10 min at 4 °C, and the supernatants were kept at -20 °C until used. Freshly thawed suspensions were briefly clarified at 11000 g at 4 °C and bioassayed by i.c. inoculation in groups of four weanling BALB/c mice per donor spleen. Bioassay recipient mice were observed daily beginning at 112 days p.i. for clinical signs of scrapie and surviving animals were observed for 360 days. Tissues from representative

animals were examined for lesions to confirm the clinical diagnosis.

Infectivity was detectable in all spleens collected from ME7-inoculated BALB/c mice. Mean incubation times for recipients of spleens collected at 30, 60 and 90 days p.i. were 187, 168 and 184 days respectively (Table 2), consistent with the observation that splenic infectivity levels reach a plateau within 2 months following subcutaneous (Eklund *et al.*, 1967) or i.c. inoculation (Race & Ernst, 1992). Recipients of inocula from the spleens of ME7-inoculated SCID mice were clinically normal throughout the 360 day observation period. Recipients of inocula from the spleens of uninoculated BALB/c and SCID mice also remained clinically normal. These observations demonstrate that scrapie infectivity, as defined by mouse subinoculation, was not detectable in the spleens of scrapie-inoculated SCID mice during the first 90 days p.i.

PrP^{Sc} accumulation was monitored by immunoblot analysis of tissues using the procedure described by Race & Ernst (1992) with slight modifications. Brains and spleens were collected from i.c. inoculated mice when they were terminally affected and from clinically normal, i.p. inoculated mice at 60 days p.i. Brains from individual mice or 200 to 250 mg aliquots of spleens from groups of three mice were prepared as 10% (w/v) homogenates in sterile 0.01 M-Tris-HCl pH 7.5, 0.005 M-MgCl₂ using disposable homogenizers, digested with DNase (final concentration 1 U/ml) for 1 h at 37 °C, and kept at -20 °C until assayed. Samples were adjusted to 10% Sarkosyl, clarified at 10000 g for 30 min at 20 °C and pelleted at 215000 g for 2 h at 4 °C. Each pellet was resuspended in 1 ml of 0.01 M-Tris-HCl pH 7.5 and incubated with proteinase K (10 µg per g starting tissue) for 30 min at 37 °C. The suspension was adjusted to 5 mM-PMSF and centrifuged at 212000 g for 1 h at 4 °C. Pellets were boiled in SDS-PAGE sample buffer (final SDS concentration 5%) and stored at -20 °C until used for Western immunoblots. PrP^{Sc} preparations (representing 1/2 spleen or 1/4 brain per lane) were electrophoresed on 12% polyacrylamide gels (Pharmacia PhastSystem), transferred to PVDF membranes (Immobilon-P, Millipore) and probed with anti-PrP or control rabbit serum. Anti-PrP serum was prepared by inoculation of rabbits with the pentadecapeptide GQGGGTHNQWNKPSK (Shinagawa *et al.*, 1986) synthesized as a multimeric macromolecule on a heptalysine backbone (serum R2843) (Tam, 1988) or covalently coupled to keyhole limpet haemocyanin (serum R27) (Wiley *et al.*, 1987; Race *et al.*, 1992). Bound rabbit antibody was detected with goat anti-rabbit IgG conjugated to horseradish peroxidase followed by incubation with a chemiluminescent substrate (Amersham) diluted 1:2 in distilled water. Membranes were exposed to ECL




Fig. 1. Immunoblots of PrP^{Sc} preparations from SCID (lane 1) or BALB/c (lane 2) mouse spleen collected 60 days after i.p. inoculation with scrapie agent ME7. Brain (lane 4) or spleen (lane 5) from clinically affected i.c. inoculated SCID mouse, or control brain from an uninoculated BALB/c mouse (lane 3) was probed with rabbit anti-PrP peptide serum (R27) as described in text.

Hyperfilm (Amersham) for intervals varying from 5 to 60 s. Both antisera gave comparable staining patterns, although R27 was 50- to 100-fold more sensitive.

PrP^{Sc} was detected in the brain but not the spleen of terminally affected, i.c. inoculated SCID mice (Fig. 1) and in both the brain and spleen of i.c. inoculated BALB/c mice. PrP^{Sc} was detected in BALB/c mouse spleen but not in SCID mouse spleen collected 60 days after i.p. inoculation. These findings are consistent with the observation that SCID mouse spleen does not accumulate PrP^{CJD} following either i.c. or i.p. inoculation (Kitamoto *et al.*, 1991). In that study, PrP^{CJD} was undetectable in the spleen of a SCID mouse collected 500 days after i.p. inoculation.

In this study, we have demonstrated that the scrapie agent does not replicate in the spleens of SCID mice. Furthermore, PrP^{Sc} does not accumulate in SCID mouse spleen following i.p. or i.c. inoculation with the scrapie agent. Failure of SCID mouse spleen to accumulate PrP^{Sc} or to harbour infectivity may be due to the lymphocyte deficiency, the failure of FDCs to mature, or to other less well characterized defects in SCID mice. Immunohistochemical evidence for PrP^C and PrP^{Sc}/PrP^{CJD} in FDCs is consistent with early experiments using cyclophosphamide-treated mice, which suggested that the tissue-associated non-mitotic cells (Worthington & Clark, 1971), rather than the circulating lymphocytes, are the site of agent replication.

The association of the scrapie agent with antigen-trapping cells such as FDCs is of interest because the agent may replicate in that cell type or it may accumulate there after replication in other tissues and transport via the peripheral circulation (Tenner-Racz *et al.*, 1985; Mori *et al.*, 1991). Pre-clinical diagnostic testing of sheep by identification of PrP^{Sc} in lymphoid tissue has been proposed (Ikegami *et al.*, 1991; Race *et al.*, 1992; Onodera *et al.*, 1993). The feasibility of such testing

increases significantly if a circulating cell type containing infectivity or PrP^{Sc} can be identified. Further discrimination between endogenous and exogenous sources of the FDC-associated scrapie agent will be critical to our understanding of the extraneural pathogenesis of the TSEs.

The authors thank Laurel Mickelsen and Sue Pritchard for technical assistance, Alberta Brassfield for helpful discussions, Julie Kooch for assistance with manuscript preparation and Dr William Davis for critical review of the manuscript. Dr Richard Race provided antiserum R27 and helpful advice on immunoblots. ME7 was kindly provided by the AFRC Neuropathogenesis Unit, Edinburgh, U.K. Mice were maintained by the personnel of the WSU Laboratory Animal Resource Center in accordance with the Public Health Service Policy on Humane Care and Use of Laboratory Animals as specified in the Health Research and Extension Act of 1985. Product names are necessary to report factually on available data; however the USDA neither guarantees nor warrants the standard of the product, and the use of the name by USDA implies no approval of the product to the exclusion of others that may also be suitable.

References

- BASLER, K., OESCH, B., SCOTT, M., WESTAWAY, D., WALCHLI, M., GROTH, D. F., MCKINLEY, M. P., PRUSINER, S. B. & WEISSMANN, C. (1986). Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene. *Cell* **46**, 417–428.
- BENDHEIM, P. E., BROWN, H. R., RUDELLI, R. D., SCALA, L. J., GOLLER, N. L., WEN, G. Y., KASCSAK, R. J., CASHMAN, N. R. & BOLTON, D. C. (1992). Nearly ubiquitous tissue distribution of the scrapie agent precursor protein. *Neurology* **42**, 149–156.
- BROWN, P. & GAJDUSEK, D. C. (1991). The human spongiform encephalopathies: Kuru, Creutzfeldt-Jakob disease, and the Gerstmann-Sträussler-Scheinker syndrome. *Current Topics in Microbiology and Immunology* **172**, 1–20.
- CAUGHEY, B., RACE, R. E. & CHESEBRO, B. (1988). Detection of prion protein mRNA in normal and scrapie-infected tissues and cell lines. *Journal of General Virology* **69**, 711–716.
- CHANDLER, R. L. (1969). Observations on the ultrastructure of the spinal cord, spleen and lymph nodes of mice inoculated with scrapie. *Research in Veterinary Science* **10**, 292–293.
- CLARKE, M. C. & KIMBERLIN, R. H. (1984). Pathogenesis of mouse scrapie: distribution of agent in the pulp and stroma of infected spleens. *Veterinary Microbiology* **9**, 215–225.
- DIRINGER, H., GELDERBLOM, H., HILMERT, H., OZEL, M., EDELBLUTH, C. & KIMBERLIN, R. H. (1983). Scrapie infectivity, fibrils and low molecular weight protein. *Nature, London* **306**, 476–478.
- DOI, S., ITO, M., SHINAGAWA, M., SATO, G., ISOMURA, H. & GOTO, H. (1988). Western blot detection of scrapie-associated fibril protein in tissues outside the central nervous system from preclinical scrapie-infected mice. *Journal of General Virology* **69**, 955–960.
- EKLUND, C. M., KENNEDY, R. C. & HADLOW, W. J. (1967). Pathogenesis of scrapie virus infection in the mouse. *Journal of Infectious Diseases* **117**, 15–22.
- FARQUHAR, C. F., SOMERVILLE, R. A. & RITCHIE, L. A. (1989). Post-mortem immunodiagnosis of scrapie and bovine spongiform encephalopathy. *Journal of Virological Methods* **24**, 215–222.
- HARTSOUGH, G. R. & BURGER, D. (1965). Encephalopathy of mink. 1. Epizootiologic and clinical observations. *Journal of Infectious Diseases* **115**, 387–392.
- IKEGAMI, Y., ITO, M., ISOMURA, H., MOMOTANI, E., SASAKI, K., MURAMATSU, Y., ISHIGURO, N. & SHINAGAWA, M. (1991). Pre-clinical and clinical diagnosis of scrapie by detection of PrP protein in tissues of sheep. *Veterinary Record* **128**, 271–275.
- KAPASI, Z. F., BURTON, G. F., SCHULTZ, L. D., TEW, J. G. & SZAKAL, A. K. (1993). Induction of functional follicular dendritic cell development in severe combined immunodeficiency mice: influence of B and T cells. *Journal of Immunology* **150**, 2648–2658.

- KITAMOTO, T., MOHRI, S. & TATEISHI, J. (1989). Organ distribution of proteinase-resistant prion protein in humans and mice with Creutzfeldt-Jakob disease. *Journal of General Virology* **70**, 3371–3379.
- KITAMOTO, T., MURAMOTO, T., MOHRI, S., DOH-URA, K. & TATEISHI, J. (1991). Abnormal isoform of prion protein accumulates in follicular dendritic cells in mice with Creutzfeldt-Jakob disease. *Journal of Virology* **65**, 6292–6295.
- LEGGETT, M. M., DUKES, J. & PIRIE, H. M. (1990). A spongiform encephalopathy in a cat. *Veterinary Record* **127**, 586–588.
- MCBRIDE, P. A., EICKELENBOOM, P., KRAAL, G., FRASER, H. & BRUCE, M. E. (1992). PrP protein is associated with follicular dendritic cells of spleens and lymph nodes in uninfected and scrapie-infected mice. *Journal of Pathology* **168**, 413–418.
- McFARLIN, D. E., RAFI, M. C., SIMPSON, E. & NEHLSSEN, S. H. (1971). Scrapie in immunologically deficient mice. *Nature, London* **233**, 336.
- MCKINLEY, M. P., BOLTON, D. C. & PRUSINER, S. B. (1983). A protease-resistant protein is a structural component of the scrapie prion. *Cell* **35**, 57–62.
- MERZ, P. A., ROHWER, R. G., KASCSAK, R. J., WISNIEWSKI, H. M., SOMERVILLE, R. A., GIBBS, C. J. & GAJDUSEK, D. C. (1984). Infection-specific particle from the unconventional slow virus diseases. *Science* **224**, 437–440.
- MERZ, P. A., KASCSAK, R. J., RUBENSTEIN, R., CARP, R. I. & WISNIEWSKI, H. M. (1987). Antisera to scrapie-associated fibril protein and prion protein decorate scrapie-associated fibrils. *Journal of Virology* **61**, 42–49.
- MORI, S., WOLFINBARGER, J. B., MIYAZAWA, M. & BLOOM, M. E. (1991). Replication of Aleutian mink disease parvovirus in lymphoid tissues of adult mink: involvement of follicular dendritic cells and macrophages. *Journal of Virology* **65**, 952–956.
- OESCH, B., WESTAWAY, D., WALCHLI, M., MCKINLEY, M. P., KENT, S. B. H., AEBERSOLD, R., BARRY, R. A., TEMPST, P., TEPLow, D. B., HOOD, L. E., PRUSINER, S. B. & WEISSMANN, C. (1985). A cellular gene encodes scrapie PrP²⁷⁻³⁰ protein. *Cell* **40**, 735–746.
- ONODERA, T., IKEDA, T., MURAMATSU, Y. & SHINAGAWA, M. (1993). Isolation of scrapie agent from the placenta of sheep with natural scrapie in Japan. *Microbiology and Immunology* **37**, 311–316.
- PRUSINER, S. B. (1982). Novel proteinaceous infectious particles cause scrapie. *Science* **216**, 136–144.
- RACE, R. E. & ERNST, D. (1992). Detection of proteinase K-resistant prion protein and infectivity in mouse spleen by 2 weeks after scrapie agent inoculation. *Journal of General Virology* **73**, 3319–3323.
- RACE, R., ERNST, D., JENNY, A., TAYLOR, W., SUTTON, D. & CAUGHEY, B. (1992). Diagnostic implications of detection of proteinase K-resistant protein in spleen, lymph nodes, and brain of sheep. *American Journal of Veterinary Research* **53**, 883–889.
- REED, L. J. & MUENCH, H. (1938). A simple method of estimating fifty percent endpoints. *American Journal of Hygiene* **27**, 493–497.
- ROBINSON, M. M. & GORHAM, J. R. (1990). Pathogenesis of hamster scrapie. Adherent splenocytes are associated with relatively high levels of infectivity. *Archives of Virology* **112**, 283–289.
- SHINAGAWA, M., MUNEKATA, E., DOI, S., TAKAHASHI, K., GOTO, H. & SATO, G. (1986). Immunoreactivity of a synthetic pentadecapeptide corresponding to the N-terminal region of the scrapie prion protein. *Journal of General Virology* **67**, 1745–1750.
- TAM, J. P. (1988). Synthetic peptide vaccine design: synthesis and properties of a high-density multiple antigenic peptide system. *Proceedings of the National Academy of Sciences, U.S.A.* **85**, 5409–5413.
- TENNER-RACZ, K., RACZ, P., DIETRICH, M. & KERN, P. (1985). Altered follicular dendritic cells and virus-like particles in AIDS and AIDS-related lymphadenopathy. *Lancet* **i**, 105–106.
- WELLS, G. A. H., SCOTT, A. C., JOHNSON, C. T., GUNNING, R. D., HANCOCK, R. D., JEFFREY, M., DAWSON, M. & BRADLEY, R. (1987). A novel progressive spongiform encephalopathy in cattle. *Veterinary Record* **121**, 419–420.
- WILEY, C. A., BURROLA, P. G., BUCHMEIER, M. J., WOODDELL, M. K., BARRY, R. A., PRUSINER, S. B. & LAMPERT, P. W. (1987). Immunogold localization of prion filaments in scrapie-infected hamster brains. *Laboratory Investigation* **57**, 646–656.
- WILLIAMS, E. W. & YOUNG, S. (1980). Chronic wasting disease of captive mule deer: a spongiform encephalopathy. *Journal of Wildlife Diseases* **16**, 89–98.
- WORTHINGTON, M. & CLARK, R. (1971). Lack of effect of immunosuppression on scrapie infection in mice. *Journal of General Virology* **13**, 349–351.

(Received 22 November 1993; Accepted 4 January 1994)