

2013

## Degree-day accumulation influences annual variability in growth of age-0 walleye

Christopher S. Uphoff

*University of Nebraska at Kearney*, [Christopher.uphoff@state.mn.us](mailto:Christopher.uphoff@state.mn.us)

Casey W. Schoenebeck

*University of Nebraska at Kearney*, [casey.schoenebeck@state.mn.us](mailto:casey.schoenebeck@state.mn.us)

W. Wyatt Hoback

*University of Nebraska at Kearney*, [whoback@okstate.edu](mailto:whoback@okstate.edu)

Keith D. Koupal

*Nebraska Game and Parks Commission, Kearney*, [keith.koupal@nebraska.gov](mailto:keith.koupal@nebraska.gov)

Kevin L. Pope

*University of Nebraska-Lincoln*, [kpope2@unl.edu](mailto:kpope2@unl.edu)

Follow this and additional works at: <http://digitalcommons.unl.edu/ncfwrustaff>

 Part of the [Aquaculture and Fisheries Commons](#), [Environmental Indicators and Impact Assessment Commons](#), [Environmental Monitoring Commons](#), [Natural Resource Economics Commons](#), [Natural Resources and Conservation Commons](#), and the [Water Resource Management Commons](#)

Uphoff, Christopher S.; Schoenebeck, Casey W.; Hoback, W. Wyatt; Koupal, Keith D.; and Pope, Kevin L., "Degree-day accumulation influences annual variability in growth of age-0 walleye" (2013). *Nebraska Cooperative Fish & Wildlife Research Unit -- Staff Publications*. 124.

<http://digitalcommons.unl.edu/ncfwrustaff/124>

This Article is brought to you for free and open access by the Nebraska Cooperative Fish & Wildlife Research Unit at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Cooperative Fish & Wildlife Research Unit -- Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



# Degree-day accumulation influences annual variability in growth of age-0 walleye



Christopher S. Uphoff<sup>a,\*</sup>, Casey W. Schoenebeck<sup>a</sup>, W. Wyatt Hoback<sup>a</sup>,  
Keith D. Koupal<sup>b</sup>, Kevin L. Pope<sup>c</sup>

<sup>a</sup> Department of Biology, University of Nebraska at Kearney, 2401 11th Avenue, Kearney, NE 68849, USA

<sup>b</sup> Nebraska Game and Parks Commission – Fisheries Research Division, 1617 First Avenue, Kearney, NE 68847, USA

<sup>c</sup> U.S. Geological Survey – Nebraska Cooperative Fish and Wildlife Research Unit, and School of Natural Resources, University of Nebraska Lincoln, 424 Hardin Hall, Lincoln, NE 68583, USA

## ARTICLE INFO

### Article history:

Received 11 January 2013

Received in revised form 29 May 2013

Accepted 31 May 2013

### Keywords:

Age-0

Degree-day

Growth

Reservoir

Walleye

## ABSTRACT

The growth of age-0 fishes influences survival, especially in temperate regions where size-dependent over-winter mortality can be substantial. Additional benefits of earlier maturation and greater fecundity may exist for faster growing individuals. This study correlated prey densities, growing-degree days, water-surface elevation, turbidity, and chlorophyll *a* with age-0 walleye *Sander vitreus* growth in a south-central Nebraska irrigation reservoir. Growth of age-0 walleye was variable between 2003 and 2011, with mean lengths ranging from 128 to 231 mm by fall (September 30th–October 15th). A set of a priori candidate models were used to assess the relative support of explanatory variables using Akaike's information criterion (AIC). A temperature model using the growing degree-days metric was the best supported model, describing 65% of the variability in annual mean lengths of age-0 walleye. The second and third best supported models included the variables chlorophyll *a* ( $r^2 = 0.49$ ) and larval freshwater drum density ( $r^2 = 0.45$ ), respectively. There have been mixed results concerning the importance of temperature effects on growth of age-0 walleye. This study supports the hypothesis that temperature is the most important predictor of age-0 walleye growth near the southwestern limits of its natural range.

© 2013 Elsevier B.V. All rights reserved.

## 1. Introduction

Growth of age-0 fishes can be an important indicator of survival (see review by Sogard, 1997), as size attained in the first year can affect over-winter survival and thus, recruitment to the fishery (Chevalier, 1973). Over-winter survival can be size-dependent as smaller age-0 fishes have higher mortality rates due to greater risks of starvation (Biro et al., 2004; Post and Evans, 1989; Post and Parkinson, 2001) and predation (Forney, 1966). Model simulations predicted that early differences in growth changed the probability of larval fish survival by as much as 30 times (Rice et al., 1993). Growth of age-0 fishes can also affect the age at which fish reach harvestable sizes and age at maturity, fecundity, and overall fitness (Beverton and Holt, 1959).

Abiotic factors, in particular, temperature can directly affect the growth of poikilotherms, such as fish (Atkinson, 1994; Kitchell et al., 1977; van der Have and de Jong, 1996) by influencing enzymatic activity, and therefore, metabolic rates (Higley et al., 1986).

Fish experience higher growth rates when their metabolic activity rises with increasing temperature (Clarke and Johnston, 1999). For example, walleye have higher growth rates at higher water temperatures regardless of latitudinal differences among populations (Galarowicz and Wahl, 2003). Other factors include turbidity, which can negatively affect the feeding efficiency of visual predators (Chesney, 1989; Utne-Palm, 2004), as well as primary productivity (inferred from chlorophyll *a* concentrations), which can affect food availability at higher trophic levels (McQueen et al., 1986). Resource availability can affect lower trophic levels, as increased nutrients from resuspension of sediments (Schindler, 1978) can increase phytoplankton biomass that may lead to increases in zooplankton biomass.

Biotic factors including prey size and availability can also affect fish growth (Paloheimo and Dickie, 1966). Walleye typically undergo a series of ontogenetic diet shifts in their first growing season, moving from zooplankton, to macroinvertebrates and finally to fish (Beck et al., 1998; Fox, 1989; Fox and Flowers, 1990; Fox et al., 1989; Johnston and Mathias, 1994a; Mathias and Li, 1982). The complexity of age-0 walleye food habits during the first growing season allows for many different scenarios in which food could be limiting (Colby and Nepszy, 1981).

\* Corresponding author. Tel.: +1 3203847721.

E-mail address: [Christopher.uphoff@state.mn.us](mailto:Christopher.uphoff@state.mn.us) (C.S. Uphoff).

<sup>1</sup> Current address: MN DNR Fisheries, 306 Power Ave N, Hinckley, MN 55037, USA.

The observed growth of age-0 walleye at Harlan County Reservoir in south-central Nebraska has been variable over the last 8 years. As an irrigation reservoir, this system is dynamic and characterized by large fluctuations in water levels and water quality, in particular during a 4-year drought (2003–2006) (Olds et al., 2011). During the drought years, reservoir volume decreased from almost 400 million m<sup>3</sup> to less than 150 million m<sup>3</sup>; during this time turbidity and chlorophyll *a* were significantly higher (Olds et al., 2011). In Harlan County Reservoir, the main prey sources of age-0 walleye have been larger calanoid copepods, age-0 gizzard shad *Dorosoma cepedianum* and some age-0 freshwater drum *Aplodinotus grunniens* (Uphoff, 2012). Densities of age-0 gizzard shad and freshwater drum are highly variable in Harlan County Reservoir (Sullivan et al., 2011a,b). At Harlan County reservoir, age-0 walleye have variable growth and abiotic and biotic factors that have influenced the growth rate of age-0 walleye cohorts in other studies also vary (Forney, 1966; Madenjian, 1991; Quist et al., 2003; Smith and Pycha, 1960; Staggs and Otis, 1996). Therefore, the objective of this study was to elucidate those factors that most strongly influenced growth of age-0 walleye among years.

## 2. Methods

### 2.1. Study site

Harlan County Reservoir is an irrigation/flood-control reservoir located in south-central Nebraska. At conservation pool the reservoir is approximately 5362 surface hectares with a mean depth of 4 m and a maximum depth of 18 m (USACE, 2010). In this polymictic system, long fetch and prevailing high winds allow only weak (i.e., temporary) thermal stratification during summer (Olds et al., 2011). The fishery is primarily managed for walleye and white bass *Morone chrysops* with gizzard shad constituting the dominant prey fish.

### 2.2. Data collection

Annual mean length (TL, mm) and catch-per-unit-effort (CPUE, *n*/min) were determined from age-0 walleye captured during boat electrofishing surveys conducted annually between September 30 and October 15 from 2003 through 2011. Age-0 walleye were collected from Nebraska Game and Parks Commission fixed shoreline stations. Electrofishing surveys were conducted at night and the typical run time at each station was 10 min.

Ten different variables were assessed for effects on annual variability in age-0 walleye growth (Table 1). Water quality and ichthyoplankton sampling protocols were established in 2003, and have been conducted every year. Zooplankton collections and water quality assessments occurred semi-weekly during April–September at 15 fixed sampling locations located throughout the reservoir (Peterson et al., 2005). Assessment occurred semi-weekly during April–September. Zooplankton were collected with an 80- $\mu$ m Wisconsin plankton net (0.5 m<sup>2</sup> opening) towed vertically from the substrate to the surface. Contents from these tows were preserved in sucrose and 4% formalin to prevent osmotic distortion (Haney and Hall, 1973). Zooplankton were identified to lowest possible taxon under a compound microscope and enumerated by taxon group. When possible at least 25 calanoid copepods (both adults and copepodites) were measured to the nearest 0.1 mm for each tow. Chlorophyll *a* concentrations and turbidity readings were taken at each station by collecting water samples every 3 m starting at 1 m with a Van Dorn bottle sampler. All water samples from each site were pooled in a bucket and stirred to assumed homogeneity. A subsample was drawn from the

integrated water sample and analyzed with an Aquafluor(tm) Handheld Fluorometer and Hach DR/890 Colorimeter.

Nighttime collections of larval gizzard shad and freshwater drum were conducted weekly during June and July at 24 fixed sampling stations (Sullivan et al., 2011a). Sampling was accomplished by pushing two ichthyoplankton nets for 5 min at a speed of 0.5 m/s. Each ichthyoplankton net was circular with the larger net having a diameter of 1.0 m (1.8 mm mesh) and the smaller net having a diameter of 0.5 m (0.75 mm mesh). These nets were mounted to the front of a boat at a depth of 0.5-m on a metal frame. Flow meters were placed at the mouth of each net to determine total volume of water sampled.

Air temperature data were obtained for Republican City, Nebraska, which is located approximately 1 mile from Harlan County Reservoir, for 2003–2011 from the National Oceanic and Atmospheric Administration's National Climatic Data Center (NCDC, 2012). Daily air temperature data were then used to calculate growing degree-days (GDD) using the following formula:

$$\text{GDD} = \left[ \frac{T_{\max} + T_{\min}}{2} \right] - T_{\text{base}}$$

where  $T_{\max}$  is the maximum daily temperature,  $T_{\min}$  is the minimum daily temperature, and  $T_{\text{base}}$  is the base temperature at which development is thought to occur. In this case,  $T_{\text{base}}$  was set at 5 °C as this temperature has previously been used to describe the walleye growing season (Kitchell et al., 1977; Neuheimer and Taggart, 2007; Shuter et al., 1983; Venturelli et al., 2010). The GDD were summed from April 1st through September 30th for all days in which the average air temperature was  $\geq 5$  °C. The GDD were used instead of water temperatures because air temperatures are more widely available and water and air temperatures are closely correlated (Livingstone and Padisak, 2007; Quist et al., 2003; Shuter et al., 1983). Daily reservoir elevation data were obtained from the U.S. Bureau of Reclamation (USBR, 2011). Mean reservoir water elevation was calculated based on daily elevations from April through September.

### 2.3. Data analysis

A set of a priori candidate models were developed to assess the relative support of explanatory variables using Akaike's information criterion (AIC). Given that all explanatory variables were available from 2003 to 2007 and 2009 to 2011, the analysis was restricted to those years. Due to small sample size relative to model parameters, second order Akaike's information criterion (AIC<sub>c</sub>) was used to more conservatively rank competing models (Burnham and Anderson, 2002). The most parsimonious models [i.e., lowest difference between model AIC<sub>c</sub> values ( $\Delta_i$ ) and highest model weight ( $W_i$ )] were chosen for model inference.

## 3. Results

The sample size for determining annual mean length ranged from 6 to 54 age-0 walleye, with all but one year having a sample size of 15 or more. No age-0 walleye were collected during 2008, therefore data from 2003 to 2007 and 2009 to 2011 were used in this analysis. Annual mean lengths of age-0 walleye varied throughout the 8 years of this study, with a mean annual length of 199 ( $\pm$ SE 11.8) mm and a range of 128–231 mm. The GDD model was the best supported by the data (i.e., lowest  $\Delta_i$  and highest  $W_i$ ) describing 65% of the variability in mean age-0 walleye length (Table 2). The GDD accumulation from April through September varied from 4391 to 5034 with a mean of 4792 ( $\pm$ SE 75) GDD from 2003 through 2011. Because mean length of age-0 walleye and GDD data were available prior to 2003 (from 1995 through 2011), we

**Table 1**  
Minimum, mean, and maximum values for each variable used to explain annual variations in mean age-0 walleye length from Harlan County Reservoir, Nebraska, during 2003–2011.

Variable	Description	Minimum	Mean	Maximum
Cal.Den	Mean calanoid copepod density (#/L) during May and June	15.7	24.3	41.8
Cal.Length	95th percentile length of calanoid copepods (mm) during May and June	46	54	70
Chlor.a	Mean Chlorophyll <i>a</i> ( $\mu\text{g/L}$ ) values during April–September	17.9	84.2	116.6
Clad.Den	Mean cladoceran density (#/L) during May and June	6.9	17.4	29.7
Drum.Den	Peak larval freshwater drum density (#/L)	0.0	0.8	2.2
GDD	Accumulation of growing degree-days (Base 5°C), from April–September	4391	4792	5034
Res.Elev	Mean reservoir elevation (ft) during April–September	1926.4	1936.6	1947.1
Shad.Den	Peak larval gizzard shad density (#/L)	0.6	2.7	5.6
Turb	Mean turbidity (FAU) values during April–September	17	29	42
WAE.CPUE	Mean CPUE (#/h) of age-0 walleye	5	24	43


**Table 2**  
Coefficient of determination ( $r^2$ ), number of parameters ( $K$ ), AIC<sub>c</sub> values, difference between successive AIC<sub>c</sub> values ( $\Delta_i$ ), and model weights ( $W_i$ ) describing rankings of a priori models based on level of support (i.e., AIC<sub>c</sub>) to evaluate interannual variability in mean age-0 walleye length from Harlan County Reservoir, Nebraska, during 2003–2011.

Model	$r^2$	$K$	AIC <sub>c</sub>	$\Delta_i$	$W_i$
GDD	0.65	3	75.14	0.00	0.623
Chlor.a	0.49	3	78.22	3.07	0.134
Drum.Den	0.45	3	78.82	3.68	0.099
Turb	0.28	3	81.00	5.85	0.033
Res.Elev	0.19	3	82.30	7.16	0.017
GDD, Shad.Den	0.73	4	82.38	7.23	0.017
Clad.Den	0.11	3	82.75	7.61	0.014
Shad.Den	0.11	3	82.75	7.61	0.014
Cal.Den	0.11	3	82.77	7.62	0.014
GDD, Chlor.a	0.70	4	83.43	8.28	0.010
WAE.CPUE	0.00	3	83.67	8.52	0.009
GDD, Cal.Den	0.66	4	84.23	9.09	0.007
Res.Elev, GDD	0.66	4	84.45	9.30	0.006
Res.Elev, Chloro.a	0.50	4	87.41	12.27	0.001
Turb, Chlor.a	0.49	4	87.52	12.38	0.001
Shad.Den, Drum.Den	0.46	4	88.12	12.97	0.001
Cal.Den, Cal.Length	0.18	4	91.43	16.29	0.000
Shad.Den, Cal.Den	0.15	4	91.65	16.50	0.000
Clad.Den, Cal.Den	0.36	4	91.76	16.62	0.000
GDD, Shad.Den, Chloro.a	0.75	5	100.66	25.51	0.000
GDD, Shad.Den, Cal.Den	0.73	5	101.02	25.88	0.000
Clad.Den, Cal.Den, Cal.Length	0.25	5	109.30	34.16	0.000

have included these additional years to show the robustness of the correlation between age-0 walleye growth and GDD over time in Harlan County Reservoir (Fig. 1). The chlorophyll *a* model and the larval freshwater drum density model were the second and third best supported models explaining 49 and 45% of the variability in mean age-0 walleye length, respectively. Other models evaluated were not supported by the data (i.e., high  $\Delta_i$  and low  $W_i$ ; Table 2).

#### 4. Discussion

Growth of age-0 walleye in Harlan County Reservoir was comparable to *Sander* spp. growth in other Great Plains reservoirs (Quist et al., 2003; Sprenkle, 2010). Past studies examining the effects of temperature on walleye growth have had mixed results. Similar to this study, summer temperatures largely explained variability in the growth of age-0 walleye in Lake Winnebago, Wisconsin (Staggs and Otis, 1996) and growth of larval walleye in culture ponds was also positively related to water temperature (Johnston, 1999). Conversely, age-0 walleye growth in a Kansas reservoir and several northern lakes appeared unrelated to temperature (Quist et al., 2003; Smith and Pycha, 1960). Forney (1966) and Madenjian (1991) both reported that temperature had a small effect on growth of



**Fig. 1.** Mean length of age-0 walleye collected from Harlan County Reservoir, Nebraska, from 1995 through 2002 (open symbols) and 2003 through 2011 (closed symbols) during fall (September 30th–October 15th) electrofishing surveys compared to the accumulation of growing degree-days [GDD (base 5°C)] from April through September of that year for the region. Solid line indicates line of best fit from simple linear regression.

age-0 walleye in Oneida Lake and in both cases prey size and availability had a larger effect on growth.

Temperature can directly influence fish growth, and results from this study indicate that the GDD metric was the best indicator of age-0 walleye growth in Harlan County Reservoir. The use of the GDD metric in fisheries science is widely applicable because temperature influences many physiological processes, such as growth due to changes in metabolic rate and consumption (Atkinson, 1994; van der Have and de Jong, 1996). Enzymatic reaction rates that govern metabolic rates, and therefore fish growth, increase with higher temperatures (Clarke and Johnston, 1999; Higley et al., 1986). For example, age-0 tiger muskellunge *Esox masquinongy* × *Esox lucius* grew faster under higher temperatures independent of feeding levels (Chipps et al., 2000). Food consumption rates can also increase with increasing temperatures, as was observed in laboratory experiments with zooplanktivorous age-0 walleye (Johnston and Mathias, 1994b).

Temperature could also be an important regulator of age-0 walleye growth through indirect effects on lake productivity. Similar to the GDD model, there was a positive relationship between chlorophyll *a* values and mean length of age-0 walleye. Both the GDD metric and chlorophyll *a* could be an index of lake productivity and phytoplankton biomass, as increased phytoplankton biomass has been associated with warming temperatures under eutrophic conditions (Tadonleke, 2010). Increased growth rates of age-0 walleye associated with increased chlorophyll *a* concentrations could also



be a result of bottom-up trophic interactions as increases in productivity at the bottom of the food chain could resonate up through the food chain and generate greater prey availability to age-0 walleye (McQueen et al., 1986). However, abundance of larval gizzard shad was not supported as a factor affecting age-0 walleye growth. The strength of the bottom-up effect usually weakens up the food chain (McQueen et al., 1986), although occasions in which bottom-up interactions have affected fish production have been reported (Ware and Thomson, 2005).

The GDD model does not differentiate between a growing season in which daily temperatures are static above the growth threshold of 5 °C and a growing season in which temperatures are variable and high for only a short period. The optimum temperature for walleye growth falls in the range of 20–23 °C (Christie and Regier, 1988; Coutant, 1977), and a growing season in which temperatures are static, within the optimum growth threshold would likely result in greater growth than would a growing season with variable high temperatures. Although the relationship of GDD and annual mean length of age-0 walleye is seemingly linear under the range of conditions during this study, the relationship likely has an upper limit or an asymptote. High daytime temperatures in southern systems may produce water temperatures that limit thermal refugia and possibly stress walleye resulting in reduced activity and slower growth (Kocovsky and Carline, 2001). Indeed, growth of mature walleye was greatest during late summer and autumn in a Kansas reservoir when water temperatures were lower than during mid summer peak when prey availability was greater (Quist et al., 2002). Summer water temperatures of 30 °C or greater that could cause stress in walleye have been observed in Kansas reservoirs and other Great Plains systems (Quist et al., 2002), however water temperatures this high do not typically occur in Harlan County Reservoir (Olds et al., 2011).

Prey abundance can influence walleye growth (Hartman and Margraf, 1992). The only model using a prey variable that was supported in this study was the larval freshwater drum density model. Occasionally age-0 freshwater drum are abundant in Harlan County Reservoir (Sullivan et al., 2011a), and at times age-0 walleye utilize freshwater drum as a prey source as in other systems (Hartman and Margraf, 1992; Uphoff, 2012). A positive relationship between water temperature and larval drum survival was reported in impoundments of the Mississippi River (Butler, 1965). Therefore, in years that age-0 walleye experienced faster growth because of greater GDD accumulation, larval freshwater drum may also experience higher survival, depicting a positive correlation between age-0 walleye mean lengths and larval drum densities.

We used a variety of models to assess the annual variability in age-0 walleye growth; however, most of these models were not supported by the data. Some of the other prey variables, such as larval gizzard shad peak densities and calanoid copepod densities, were not predictors of age-0 walleye growth likely because these prey sources are not limiting in Harlan County Reservoir (Sullivan et al., 2011b). Also, cladoceran densities may not have had an effect on growth because age-0 walleye in Harlan County Reservoir selected for calanoid copepods (Uphoff, 2012), unlike previous studies that reported cladocerans to be an important prey of age-0 walleye (Beck et al., 1998). Relationships between prey variables and age-0 walleye growth would likely become increasingly more important if prey become limiting. Non-limiting prey sources would also explain why age-0 walleye growth was not density-dependent. The *a priori* assessments may not have included the best possible descriptor of age-0 walleye growth in Harlan County Reservoir. Other variables that could influence age-0 walleye growth are mean length of age-0 gizzard shad in October and densities of macroinvertebrates. Overall, this study adds to previous research supporting the importance of temperature on the growth of age-0 walleye, due to direct and indirect influences of

temperature on metabolic rates, consumption, and lake productivity.

## Acknowledgements

We thank the Nebraska Game and Parks Commission for the use of trucks, boats, and equipment and the U.S. Army Corp of Engineers at Harlan County Reservoir for boat and equipment storage. We thank the University of Nebraska at Kearney for their technical support. We also thank Jason Deboer for providing air temperature data and Brian Peterson, Josh Kreitman, Eric Smits, and multiple graduate students, technicians, and volunteers for providing field assistance with this project. This project was funded by Federal Aid in Sport Fish Restoration funds (Federal Aid Project F-160-R-8) administered by the Nebraska Game and Parks Commission. The use of trade names or products does not constitute endorsement by the U.S. Government. The Nebraska Cooperative Fish and Wildlife Research Unit is jointly supported by a cooperative agreement among the U.S. Geological Survey, the Nebraska Game and Parks Commission, the University of Nebraska, the U.S. Fish and Wildlife Service and the Wildlife Management Institute.

## References

- Atkinson, D., 1994. Temperature and organism size – a biological law for ectotherms? *Advances in Ecological Research* 25, 1–58.
- Beck, H.D., Starostka, A.B., Willis, D.W., 1998. Diet overlap of age-0 walleye and white bass in Lake Poinsett, South Dakota. *Journal of Freshwater Ecology* 13, 425–431.
- Beverton, R.J.H., Holt, S.J., 1959. A review of the lifespans and mortality rates of fish in nature and the relation to growth and other physiological characteristics. In: Wolsenholmy, G.E.W., O'Connor, M. (Eds.), *Colloquia on Ageing. The Life Span of Animals*, vol. 5. CIBA Foundation, Churchill, London, pp. 142–177.
- Biro, P.A., Morton, A.E., Post, J.R., Parkinson, E.A., 2004. Over-winter lipid depletion and mortality of age-0 rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences* 61, 1513–1519.
- Burnham, K.P., Anderson, D.R., 2002. *Model Selection and Multimodel Inference: A Practical Information Theoretic Approach*, 2nd edition. Springer Science, New York.
- Butler, R.L., 1965. Freshwater drum, *Aplodinotus grunniens*, in the navigational impoundments of the Mississippi River. *Transactions of the American Fisheries Society* 94, 339–349.
- Chesney, E.J., 1989. Estimating the food requirement of striped bass larvae *Morone saxatilis*: effects of light, turbidity and turbulence. *Marine Ecology Progress Series* 53, 191–200.
- Chevalier, J.R., 1973. Cannibalism as a factor in first year survival of walleye in Oneida Lake. *Transactions of the American Fisheries Society* 102, 739–744.
- Chippis, S.R., Einfalt, L.M., Wahl, D.H., 2000. Growth and food consumption by tiger muskellunge: effects of temperature and ration level on bioenergetic model predictions. *Transactions of the American Fisheries Society* 129, 186–193.
- Christie, G.C., Regier, H.A., 1988. Measures of optimal thermal habitat and their relationship to yields for four commercial fish species. *Canadian Journal of Fisheries and Aquatic Sciences* 45, 301–314.
- Clarke, A., Johnston, N.M., 1999. Scaling of metabolic rate with body mass and temperature in teleost fish. *Journal of Animal Ecology* 68, 893–905.
- Colby, P.J., Nepszy, S.J., 1981. Variation among stocks of walleye (*Stizostedion vitreum vitreum*): management implications. *Canadian Journal of Fisheries and Aquatic Sciences* 38, 1814–1831.
- Coutant, C.C., 1977. Compilation of temperature preference data. *Journal of the Fisheries Research Board of Canada* 34, 739–745.
- Forney, J.L., 1966. Factors affecting first-year growth of walleyes in Oneida Lake, New York. *New York Fish Game Journal* 13, 146–167.
- Fox, M.G., 1989. Effect of prey density and prey size on growth and survival of juvenile walleye (*Stizostedion vitreum vitreum*). *Canadian Journal of Fisheries and Aquatic Sciences* 46, 1323–1328.
- Fox, M.G., Flowers, D.D., 1990. Effect of fish density on growth, survival, and food consumption by juvenile walleyes in rearing ponds. *Transactions of the American Fisheries Society* 119, 112–121.
- Fox, M.G., Keast, J.A., Swainson, R.J., 1989. The effect of fertilization regime on juvenile walleye growth and prey utilization in rearing ponds. *Environmental Biology of Fishes* 26, 129–142.
- Galarowicz, T.L., Wahl, D.H., 2003. Differences in growth, consumption, and metabolism among walleyes from different latitudes. *Transactions of the American Fisheries Society* 132, 425–437.
- Haney, J.F., Hall, D.J., 1973. Sugar-coated *Daphnia*: a preservation technique for Cladocera. *Limnology and Oceanography* 29, 941–948.
- Hartman, K.J., Margraf, F.J., 1992. Effects of prey and predator abundances on prey consumption and growth of walleyes in western Lake Erie. *Transactions of the American Fisheries Society* 121, 245–260.

- Higley, L.G., Pedigo, L.P., Ostlie, K.R., 1986. DEGDAY: a program for calculating degree-days and assumptions behind the degree-day approach. *Environmental Entomology* 15, 999–1016.
- Johnston, T.A., Mathias, J.A., 1994a. Feeding ecology of walleye, *Stizostedion vitreum*, larvae: effects of body size, zooplankton abundance and zooplankton community composition. *Canadian Journal of Fisheries and Aquatic Sciences* 51, 2077–2089.
- Johnston, T.A., Mathias, J.A., 1994b. The effects of temperature on feeding in zooplanktivorous walleye, *Stizostedion vitreum*, larvae. *Environmental Biology of Fishes* 40, 189–198.
- Johnston, T.A., 1999. Factors affecting food consumption and growth of walleye, *Stizostedion vitreum*, larvae in culture ponds. *Hydrobiologia* 400, 129–140.
- Kitchell, J.F., Stewart, D.J., Weininger, D., 1977. Applications of a bioenergetics model to yellow perch (*Perca flavescens*) and walleye (*Stizostedion vitreum vitreum*). *Journal of the Fisheries Research Board of Canada* 34, 1922–1935.
- Kocovsky, P.M., Carline, R.F., 2001. Influence of extreme temperatures on consumption and condition of walleyes in Pymatuning Sanctuary, Pennsylvania. *North American Journal of Fisheries Management* 21, 198–207.
- Livingstone, D.M., Padisak, J., 2007. Large-scale coherence in the response of lake surface-water temperatures to synoptic-scale climate forcing during summer. *Limnology and Oceanography* 52, 896–902.
- Madenjian, C.P., 1991. Limits to growth of young-of-the-year walleye (*Stizostedion vitreum vitreum*): an individual-based model perspective. *Canadian Journal of Fisheries and Aquatic Sciences* 48, 1492–1499.
- Mathias, J.A., Li, S., 1982. Feeding habits of walleye larvae and juveniles: comparative laboratory and field studies. *Transactions of the American Fisheries Society* 111, 722–735.
- McQueen, D.J., Post, J.R., Mills, E.L., 1986. Trophic relationships in freshwater pelagic ecosystems. *Canadian Journal of Fisheries and Aquatic Sciences* 43, 1571–1581.
- NCDC (National Climatic Data Center), 2012. Record of Climatological Observations, <http://www.ncdc.noaa.gov/oa/ncdc.html> (accessed 03.11.12).
- Neuheimer, A.B., Taggart, C.T., 2007. The growing degree-day and fish size-at-age: the overlooked metric. *Canadian Journal of Fisheries and Aquatic Sciences* 64, 375–385.
- Olds, B.P., Peterson, B.C., Koupal, K.D., Farnsworth-Hoback, K.M., Schoenebeck, C.W., Hoback, W.W., 2011. Water quality parameters of a Nebraska reservoir differ between drought and normal conditions. *Lake and Reservoir Management* 27, 229–234.
- Paloheimo, J.E., Dickie, L.M., 1966. Food and growth of fishes. II. Effects of food and temperature on the relation between metabolism and body weight. *Journal of the Fisheries Research Board of Canada* 23, 869–908.
- Peterson, B.C., Fryda, N.J., Koupal, K.D., Hoback, W.W., 2005. Daphnia lumholzi, an exotic zooplankton, invading a Nebraska Reservoir. *Prairie Naturalist* 37, 11–19.
- Post, J.R., Evans, D.O., 1989. Size-dependent overwinter mortality of young-of-the-year yellow perch (*Perca flavescens*): laboratory, in situ enclosure, and field experiments. *Canadian Journal of Fisheries and Aquatic Sciences* 46, 1958–1968.
- Post, J.R., Parkinson, E.A., 2001. Energy allocation strategy in young fish: allometry and survival. *Ecology* 82, 1040–1051.
- Quist, M.C., Guy, C.S., Bernot, R.J., Stephen, J.L., 2002. Seasonal variation in condition, growth and food habits of walleye in a Great Plains reservoir and simulated effects of an altered thermal regime. *Journal of Fish Biology* 61, 1329–1344.
- Quist, M.C., Guy, C.S., Shultz, R.D., Stephen, J.L., 2003. Latitudinal comparisons of walleye growth in North America and factors influencing growth of walleyes in Kansas Reservoirs. *North American Journal of Fisheries Management* 23, 677–692.
- Rice, J.A., Miller, T.J., Rose, K.A., Crowder, L.B., Marschall, E.A., Trebitz, A.S., DeAngelis, D.L., 1993. Growth-rate variation and larval survival: inferences from an individual-based size-dependent predation model. *Canadian Journal of Fisheries and Aquatic Sciences* 50, 133–142.
- Schindler, D.W., 1978. Factors regulating phytoplankton production and standing crop in the world's lakes. *Limnology and Oceanography* 23, 478–486.
- Shuter, B.J., Schlesinger, D.A., Zimmerman, A.P., 1983. Empirical predictors of annual surface water temperature cycles in North American lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 40, 1838–1845.
- Smith Jr., L.L., Pycha, R.L., 1960. First-year growth and survival in the walleye, *Stizostedion vitreum vitreum* (Mitchill), and associated factors in the Red Lakes, Minnesota. *Limnology and Oceanography* 5, 281–290.
- Sogard, S.M., 1997. Size-selective mortality in the juvenile stage of teleost fishes: a review. *Bulletin of Marine Science* 60, 1129–1157.
- Sprenkle, E., 2010. Tuttle Creek Reservoir Progress and Management Report. Kansas Department of Wildlife and Parks.
- Staggs, M.S., Otis, K.J., 1996. Factors affecting first-year growth of fishes in Lake Winnebago, Wisconsin. *North American Journal of Fisheries Management* 16, 608–618.
- Sullivan, C.L., Koupal, K.D., Hoback, W.W., Peterson, B.C., Schoenebeck, C.W., 2011a. Food habits and abundance of larval freshwater drum in a South Central Nebraska irrigation reservoir. *Journal of Freshwater Ecology* 27, 111–121.
- Sullivan, C.L., Schoenebeck, C.W., Koupal, K.D., Hoback, W.W., Peterson, B.C., 2011b. Patterns of age-0 gizzard shad abundance and food habits in a Nebraska irrigation reservoir. *Prairie Naturalist* 43, 110–116.
- Tadonleke, R.D., 2010. Evidence of warming effects on phytoplankton productivity rates and their dependence on eutrophication status. *Limnology and Oceanography* 55, 973–982.
- Uphoff, C.S., 2012. Seasonal food habits and variability in length of age-0 walleye in Harlan County Reservoir, Nebraska. University of Nebraska at Kearney, Kearney, NE (thesis).
- USACE (United States Army Corps of Engineers), 2010. <http://www.nwk.usace.army.mil/hc/> (accessed 10.02.10).
- USBR (United States Bureau of Reclamation), 2011. Harlan County Reservoir, [http://www.usbr.gov/gp-bin/arc050\\_form.pl?HCNE](http://www.usbr.gov/gp-bin/arc050_form.pl?HCNE) (accessed 07.11.11).
- Utne-Palm, A.C., 2004. Effects of larvae ontogeny, turbidity, and turbulence on prey attack rate and swimming activity of Atlantic herring larvae. *Journal of Experimental Marine Biology and Ecology* 314, 147–161.
- van der Have, T.M., de Jong, G., 1996. Adult size in ectotherms: temperature effects on growth and differentiation. *Journal of Theoretical Biology* 183, 329–340.
- Venturelli, P.A., Lester, N.P., Marshall, T.R., Shuter, B.J., 2010. Consistent patterns of maturity and density dependent growth among populations of walleye (*Sander vitreus*): application of the growing degree-day metric. *Canadian Journal of Fisheries and Aquatic Sciences* 67, 1057–1067.
- Ware, D.M., Thomson, R.E., 2005. Bottom-up ecosystem trophic dynamics determine fish production in the northeast Pacific. *Science* 308, 1280–1284.