

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications in Food Science and  
Technology

Food Science and Technology Department

---

2013

## Legume allergy: clinical threshold doses in legume-allergic individuals

Benjamin C. Remington

*University of Nebraska-Lincoln*, ben.c.remington@gmail.com

WM Blom

*TNO, Zeist, Germany*

GF Houben

*TNO, Zeist, Germany*

A Knulst

*UMC Utrecht*, a.c.knulst@umcutrecht.nl

J. L. Baumert

*University of Nebraska-Lincoln*, jbaumert2@unl.edu

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/foodsciefacpub>

---

Remington, Benjamin C.; Blom, WM; Houben, GF; Knulst, A; Baumert, J. L.; and Taylor, S. L., "Legume allergy: clinical threshold doses in legume-allergic individuals" (2013). *Faculty Publications in Food Science and Technology*. 124.

<https://digitalcommons.unl.edu/foodsciefacpub/124>

This Article is brought to you for free and open access by the Food Science and Technology Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in Food Science and Technology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Benjamin C. Remington, WM Blom, GF Houben, A Knulst, J. L. Baumert, and S. L. Taylor



POSTER PRESENTATION

Open Access

# Legume allergy: clinical threshold doses in legume-allergic individuals

BC Remington<sup>1</sup>, WM Blom<sup>2\*</sup>, GF Houben<sup>2,3</sup>, AC Knulst<sup>3</sup>, JL Baumert<sup>1</sup>, SL Taylor<sup>1</sup>

From Food Allergy and Anaphylaxis Meeting (FAAM 2013)  
Nice, France. 7-9 February 2013

## Background

Soybean and peanut, members of the Legume family, are recognized as common allergenic foods by the FAO Codex Alimentarius. EU directive 2003/89/EC and US Food Allergen Labeling and Consumer Protection Act prescribe labelling of food products for these two major allergenic foods. EU directive 2007/68/EC requires labelling for lupin, another member of the Legume family. The reported prevalence and severity of allergic reactions to soybeans is lower than peanut and most other major allergenic foods. For lupin only limited clinical data are available. Our previous work with peanut has used clinical results from double-blind, placebo-controlled food challenges (DBPCFC) to estimate a population threshold. Further exploration of soy and lupin thresholds would benefit all stakeholders.

## Methods

Publications were screened for DBPCFCs providing lowest observed adverse effect level (LOAEL) and/or no observed adverse effect level (NOAEL) for soy- or lupin-allergic individuals. Unpublished data were retrieved from diagnostic DBPCFCs from the UMCU outpatient population. LOAELs were based upon elicitation of objective symptoms. All doses were converted to mg protein to standardize challenge materials across studies. Individual thresholds were analyzed using Interval-Censoring Survival Analysis and fitted to parametric models (Log-Normal, Log-Logistic, and Weibul) using LIFEREG procedure (SAS v9.2). The eliciting dose predicted to provoke reactions in a proportion of the allergic population (ED<sub>p</sub>) was estimated and compared to the known peanut-allergic population.

## Results

Public literature revealed clinical thresholds of 43 soy-allergic individuals in 4 studies and 9 lupin-allergic individuals in 3 studies. An additional 15 unpublished lupin DBPCFC were retrieved. Comparison of the ED doses indicate that soy and lupin are less potent allergenic foods than peanut. Using the Log-Normal distributions, the ED<sub>05</sub> for peanut is 1.3 mg peanut protein while the ED<sub>05</sub> for lupin is 19.1 mg lupin protein and soybean is 24.5 mg soy protein. However, the lowest eliciting dose observed for each group of allergic individuals is 36 mg lupin protein and 88 mg soy protein, considerably higher than the predicted ED<sub>05</sub>.

## Conclusion

The eliciting dose for allergic reactions of three members of the Legume family showed a different potency for peanut, lupin, and soybean. Additional clinical data from soy and/or lupin challenges would be valuable to increase the statistical confidence in the population threshold estimates.

## Disclosure of interest

None declared.

## Author details

<sup>1</sup>Food Allergy Research & Resource Program, Department of Food Science & Technology, University of Nebraska-Lincoln, Lincoln, NE, USA. <sup>2</sup>Food and Nutrition, TNO, Zeist, Germany. <sup>3</sup>Department of Dermatology/Allergology, University Medical Center Utrecht, Utrecht, the Netherlands.

Published: 25 July 2013

doi:10.1186/2045-7022-3-S3-P141

**Cite this article as:** Remington et al.: Legume allergy: clinical threshold doses in legume-allergic individuals. *Clinical and Translational Allergy* 2013 **3**(Suppl 3):P141.

<sup>2</sup>Food and Nutrition, TNO, Zeist, Germany  
Full list of author information is available at the end of the article