

2010

EVALUATING CHEMICAL DETERRENCE AT TWO SPATIAL SCALES: THE EFFECTIVENESS OF CHEMICAL DETERRENCE FOR SANDHILL CRANES IN CORNFIELDS

ANNE E. LACY

International Crane Foundation

JEB A. BARZEN

International Crane Foundation

Follow this and additional works at: <http://digitalcommons.unl.edu/nacwgproc>

 Part of the [Behavior and Ethology Commons](#), [Biodiversity Commons](#), [Ornithology Commons](#), [Population Biology Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

LACY, ANNE E. and BARZEN, JEB A., "EVALUATING CHEMICAL DETERRENCE AT TWO SPATIAL SCALES: THE EFFECTIVENESS OF CHEMICAL DETERRENCE FOR SANDHILL CRANES IN CORNFIELDS" (2010). *North American Crane Workshop Proceedings*. 131.

<http://digitalcommons.unl.edu/nacwgproc/131>

This Article is brought to you for free and open access by the North American Crane Working Group at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in North American Crane Workshop Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

EVALUATING CHEMICAL DETERRENCE AT TWO SPATIAL SCALES: THE EFFECTIVENESS OF CHEMICAL DETERRENCE FOR SANDHILL CRANES IN CORNFIELDS

ANNE E. LACY, International Crane Foundation, E-11376 Shady Lane Road, Baraboo, Wisconsin 53913, USA
JEB A. BARZEN, International Crane Foundation, E-11376 Shady Lane Road, Baraboo, WI 53913, USA

Abstract: From 2006 through 2008, 9,10 anthraquinone (sold as Avitec™) was used as a deterrent on planted corn seed in Minnesota, Wisconsin, and Michigan. ICF conducted field trials in Wisconsin to determine efficacy of Avitec™ to repel sandhill cranes (*Grus canadensis*) from germinating corn. We assessed crane use at 2 levels: between and within habitats by crane population surveys to determine crane use of fields, and corn density surveys to assess possible damage within fields. In addition, corn seed samples were taken to assess amount of active ingredient on treated corn seeds in the ground. In 2008 the concentrations of Avitec™ on seed obtained from powder treatments (as compared to liquid treated) were generally lower. Where concentration of Avitec™ on the corn seeds was adequate (liquid or powder), it successfully deterred crane herbivory even though crane use of the fields remained high. Non-treated fields had higher damage as crane use increased, whereas treated fields had low or no damage, even with increased crane use. An effective deterrent is a win-win situation for both cranes and farmers. Its use protects a valuable crop while allowing cranes to access critical food items in cultivated fields, which also confers a benefit to the farmer (i.e., consumption of crop pests). Farmers can solve the problem more economically on their own without handling toxic seed treatments. Successful solutions such as this example are critical for advancing wildlife conservation on private lands.

PROCEEDINGS OF THE NORTH AMERICAN CRANE WORKSHOP 11:207

Key words: anthraquinone, chemical deterrent, corn, *Grus canadensis*, sandhill crane, spatial scale.
