

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Transactions of the Nebraska Academy of
Sciences and Affiliated Societies

Nebraska Academy of Sciences

1992

Pollination Characteristics of Blowout Penstemon (*Penstemon haydenii* S. Watson)

Theresa R. Flessner
USDA- Soil Conservation Service

James L. Stubbendieck
University of Nebraska - Lincoln, jstubbendieck@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/tnas>

 Part of the [Life Sciences Commons](#)

Flessner, Theresa R. and Stubbendieck, James L., "Pollination Characteristics of Blowout Penstemon (*Penstemon haydenii* S. Watson)" (1992). *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*. 131.

<https://digitalcommons.unl.edu/tnas/131>

This Article is brought to you for free and open access by the Nebraska Academy of Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Transactions of the Nebraska Academy of Sciences and Affiliated Societies by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

**POLLINATION CHARACTERISTICS OF
BLOWOUT PENSTEMON (*PENSTEMON HAYDENII* S. WATSON)***

Theresa R. Flessner

USDA-Soil Conservation Service
Plant Materials Center, 3415 N.W. Granger Avenue
Corvallis, Oregon 97330

James Stubbendieck

Department of Agronomy
University of Nebraska–Lincoln
Lincoln, Nebraska 68583-0915

*Journal Series No. 9711, Agricultural Research Division, University of Nebraska–Lincoln

ABSTRACT

The pollination characteristics of blowout penstemon (*Penstemon haydenii* S. Wats.), the only officially endangered plant species in Nebraska, were investigated. Cross-pollination was determined to be the major method of pollination, but self-pollination also occurs. Insects increased cross-pollination. The natural habitat of blowout penstemon has greatly decreased during this century, resulting in only a few isolated populations. The opportunity for cross-pollination with other populations no longer exists. Without cross-pollination, lowered genetic diversity of the species may result in less vigorous plants and declining populations. Introduction of new genetic material into declining populations may help to assure the continued existence of the species.

† † †

Blowout penstemon (*Penstemon haydenii* S. Wats.) is a rare, multistemmed, perennial forb that occurs naturally only in a few Sandhills blowouts, sites of active wind erosion (Stubbendieck et al., 1989). It is endemic to the Nebraska Sandhills and is the only plant species in Nebraska classified as endangered and protected under the federal Endangered Species Act (Fish and Wildlife Service, 1987; Jobman, 1989).

Blowout penstemon was once a common plant in blowouts (Pool, 1914). Populations rapidly declined until it was thought to be extinct by mid-century. Since its rediscovery in 1968 (Sutherland, 1988), extensive searching led to locating about 6,000 plants in only a few isolated blowouts (Fritz et al., 1991). Isolated

populations may have a higher vulnerability to extinction because of low genetic variability restricting adaptation to changing environments (Hardy et al., 1989; Soule, 1983). Hardy et al. (1989) stated that decreased vigor, viability, and fecundity associated with inbreeding can be the result of genetic deterioration often occurring in small out-crossing populations. The objective of this study was to determine the pollination characteristics of blowout penstemon.

MATERIALS AND METHODS

The study was conducted in a natural population of blowout penstemon in Cherry County in 1987 and 1988. Four blowout penstemon plants, each having at least six inflorescences, were selected for this study in 1987, and another four plants were selected in 1988. Each plant was considered to be a block. Treatments designed to determine the breeding system of blowout penstemon were randomly applied to inflorescences of each plant.

Before applying a treatment, all immature flower buds, opened flowers, and fruits of the inflorescence were removed with forceps. To standardize the inflorescence, the number of mature (but unopened) flower buds was reduced to 10 per inflorescence. Treatments were two levels of emasculation (not emasculated and emasculated), two levels of bagging (unbagged and bagged), and hand-self- and cross-pollination by hand. Emasculation involved removal of the anthers within the bud with a forceps. Bagging was accomplished by enclosing the inflorescence within a tightly woven cot-

ton bag and securely tying the open end around the base of the inflorescence with a small wire. The mesh was sufficiently fine to exclude all potential insect visitors and pollen.

Hand-pollination included emasculation, the application of pollen with a fine camel's hair brush to the stigma, and enclosing the treated flower stalk with a cloth bag (Hartmann and Kester, 1975). To self-pollinate flowers, pollen collected from flowers of the same plant just prior to hand-pollination was used. To cross-pollinate flowers, pollen collected from a blowout penstemon plant growing in a display garden on the University of Nebraska-Lincoln campus 10 days prior to hand-pollination was used. The display plant was propagated from seed collected in Cherry County. Pollen collection involved removing the anthers from unopened buds, placing them on a sheet of paper, and allowing them to open in a germinator at 30°C. Upon dehiscence, pollen and anthers were removed and stored in a glass vial over CaCl₂ at 0°C. Pollen was transported to Cherry County.

Thus, each of the 6 treatments was replicated 4 times in a randomized complete block design. The experimental unit consisted of a standardized inflorescence. In mid-August, 1987, and mid-August, 1988, all treated inflorescences were harvested, placed in sepa-

rate paper bags, and dried at 25–35°C. After drying, the number of developed fruits, number of seeds, and seed weight were recorded per treated inflorescence. Statistical analysis was performed on data using SAS, PROC GLM for balanced designs (SAS Institute Inc., 1985). Univariate analysis of variance (ANOVA) with preplanned contrasts was used to determine the effect of bagging, emasculation, and hand-pollination on number of fruits, number of seeds, and seed weight (Steel and Torrie, 1980). Mean separation was based upon the 0.05 level of probability. Each year was analyzed separately.

RESULTS AND DISCUSSION

In both 1987 and 1988, inflorescences with emasculated flowers produced nearly the same mean number of fruits, seeds, and seed weight as inflorescences with intact flowers, indicating that cross-pollination was the major method of pollination, but self-pollination also occurred (Table I). Unbagged inflorescences with emasculated flowers produced significantly more seed than unbagged inflorescences with intact flowers in 1988, indicating that cross-pollination was more effective that year (Table I).

In both trials, the bagged inflorescences produced fewer fruits and seeds and lower seed weight on the

Table I. Mean number of fruits and seeds, and mean seed weight (g) produced per treated blowout penstemon inflorescence in 1987 and 1988. Treatment contrasts describe the effects of various treatments on fruit and seed production.

Treatment	1987			1988		
	# fruits	# seeds	seed (g)	# fruits	# seeds	seed (g)
1. Unbagged intact	7	96	0.15	4	12	0.30
2. Unbagged emasculated	8	103	0.19	6	53	0.42
3. Bagged intact	3	0	0.00	0	0	0.00
4. Bagged emasculated	0	0	0.00	0	0	0.00
5. Hand-pollinated, self	<1	<1	<0.01	<1	<1	0.02
6. Hand-pollinated, crossed	2	4	0.01	6	49	0.29
Treatment Contrasts ¹	Pr>F					
Unbagged vs bagged	0.01	<0.01	<0.01	<0.01	0.02	<0.01
Intact vs. emasculated	0.92	0.54	0.80	0.58	0.93	0.99
Natural crossing vs. hand	0.03	0.01	0.02	0.70	0.66	0.51
Natural selfing vs. hand	0.24	0.99	0.97	0.87	0.99	0.87
Self- vs. cross (hand)	0.97	0.61	0.49	<0.01	0.03	0.07

¹Treatment contrasts were 1, 2 vs 3, 4; 1, 3 vs 2, 4; 2 vs 6; 3 vs 5; and 5 vs 6.

average than unbagged inflorescences, indicating the importance of insects for cross-pollination (Table I). Lower fruit and seed production in 1988 than 1987 is probably a reflection of lower precipitation recorded during the growing season of 1988.

In 1987, the hand-pollinated, bagged inflorescences set fewer seeds than naturally-pollinated inflorescences, indicating inadequate pollination techniques (Table I). In 1988, artificially cross-pollinated inflorescences produced as many fruits, seeds, and seed weights on the average as unbagged inflorescences with emasculated flowers (Table I). The fruit and seed production in artificially self-pollinated inflorescences were limited but not different from naturally self-pollinated inflorescences which did not produce fruit or seed in either year. Pollination techniques presumably improved with practice, and ambient conditions may have influenced success. Fruit and seed production were also greater in artificially cross-pollinated inflorescences than in artificially self-pollinated inflorescences, again indicating that blowout penstemon is largely an outcrosser (Table I).

Many penstemon species exhibit both self- and cross-pollination. For example, Ramstetter and Peterson (1984) reported that *Penstemon lemhiensis* (Keck) Keck & Cronq. was largely an outcrosser, but that many of the plants are self-pollinated as well. Baurer (1983) reported that cross-pollination was important, but not essential, for fruit production of little-flower penstemon (*Penstemon procerus* Dougl.).

Penstemon spp. pollinators include hummingbirds, coleopterans, carpenter bees, honeybees, bumblebees, solitary bees, masarid wasps, bee flies, syrphid flies, moths, hymenopterans, dipterans, and lepidopterans (Schmid, 1976). A wasp *Pseudomasaris vespoides* (Cresson), which provisions its nest exclusively with pollen and nectar, utilizes various *Penstemon* spp. a majority of the time (Tepedino, 1979; Ramstetter and Peterson, 1984). This wasp and many *Penstemon* spp. appear to have coevolved many adaptive features to ensure successful cross-pollination (Straw, 1956; Torchio, 1974).

Several pollen collectors and other insects have been reported to visit blowout penstemon flowers (Flessner, 1988; Lawson et al., 1989). Many different insects were seen removing pollen and nectar from the flowers of blowout penstemon during this study. Thus, it is hypothesized that blowout penstemon pollination is not restricted by inadequate numbers and species of pollinators. Additional research determining the phenology and insect pollinators of blowout penstemon is needed to describe further the pollination characteristics of this species.

Since the natural habitat (the blowout) of blowout penstemon has greatly decreased with improved range management practices and wildfire control, the populations of blowout penstemon have become more isolated (Stubbenieck et al., 1989). Thus, the potential for cross-pollination with other populations has also decreased. Perhaps, a reduction in dispersal and cross-pollination has lowered the genetic diversity of this species, resulting in less vigorous plants followed by declining populations. Introduction of new genetic material into declining populations should be considered as a way to improve vigor of resultant seedlings. Further research on the vigor and reproductive capability of seedlings produced from crosses might aid in the development of vigorous blowout penstemon seedlings for transplant and, ultimately, its recovery.

ACKNOWLEDGMENTS

Research reported here is a part of a project on the status, propagation, and ecology of blowout penstemon funded by the Nebraska Game and Parks Commission.

LITERATURE CITED

- Baurer, P. J. 1983. Bumble bee pollination relationships on the Beartooth Plateau tundra of southern Montana, USA. *American Journal of Botany* 70: 134-144.
- Fish and Wildlife Service. 1987. Final rule to determine *Penstemon haydenii* (blowout penstemon) to be an endangered species. *Federal Register* 52: 32926-32929.
- Flessner, T. R. 1988. Propagation, establishment, and ecological characteristics of *Penstemon haydenii* S. Wats. Master of Science Thesis. University of Nebraska-Lincoln.
- Fritz, M., J. Stubbenieck, and W. Jobman. 1991. *Blowout penstemon recovery plan* (draft). U.S. Fish and Wildlife Service, Denver.
- Hardy, J. P., R. R. Weedon, and W. R. Bowlin. 1989. Blowout penstemon: Description and present situation. In: T. B. Bragg and J. Stubbenieck (eds.), *Proceedings of the Eleventh North American Prairie Conference*. University of Nebraska-Lincoln: 227-231.
- Hartmann, H. T., and D. E. Kester. 1975. *Plant propagation: principles and practices*. Englewood Cliffs, New Jersey, Prentice Hall, Inc: 662 pp.
- Jobman, W. G. 1989. Determining federally listed endangered and threatened species, emphasizing blowout penstemon. In: T. B. Bragg and J. Stubbenieck (eds.), *Proceedings of the Eleventh North American Prairie Conference*. University of Nebraska-Lincoln: 241-245.

- Lawson, H. R., V. J. Tepedino, and T. L. Griswold. 1989. Pollen collectors and other insect visitors to *Penstemon haydenii* S. Wats. In: T. B. Bragg and J. Stubbendieck (eds.), *Proceedings of the Eleventh North American Prairie Conference*. University of Nebraska—Lincoln: 233–235.
- Pool, R. J. 1914. A study of the vegetation of the Sandhills of Nebraska. *Minnesota Botanical Studies* 4: 188–312.
- Ramstetter, J., and K. M. Peterson. 1984. Pollination biology of *Penstemon lemhiensis*. *American Journal of Botany* 71: 87.
- SAS Institute Inc. 1985. *SAS User's Guide: Statistics, version 5 edition*. Cary, North Carolina, SAS Institute Inc.: 956 pp.
- Schmid, R. 1976. Fly pollination of *Penstemon davidsonii* and *P. procerus* (Scrophulariaceae). *Madroño* 23: 401–402.
- Soule, M. E. 1983. What should we do about extinction? In: C. M. Schonewald-Cox (ed.) *Genetics and Conservation*. Menlo Park, California, Benjamin/Cummings: 111–124.
- Steel, R. G. D., and J. H. Torrie. 1980. *Principles and procedures of statistics*. New York, McGraw-Hill: 633 pp.
- Straw, R. M. 1956. Adaptive morphology of the *Penstemon* flower. *Phytomorphology* 6: 112–119.
- Stubbendieck, J., T. R. Flessner, and R. R. Weedon. 1989. Blowouts in the Nebraska Sandhills: The habitat of *Penstemon haydenii*. In: T. B. Bragg and J. Stubbendieck (eds.), *Proceedings of the Eleventh North American Prairie Conference*. University of Nebraska—Lincoln: 223–225.
- Sutherland, D. M. 1988. Historical notes on collections and taxonomy of *Penstemon haydenii* S. Wats. (blowout penstemon), Nebraska's only endemic plant species. *Transactions of the Nebraska Academy of Sciences* 16: 191–194.
- Tepedino, V. J. 1979. Notes on the flower visiting habits of *Pseudomasaris vespoides* (Hymenoptera: Masaridae). *Southwestern Naturalist* 24: 380–381.
- Torchio, P. V. 1974. Mechanisms involved in the pollination of *Penstemon* visited by the masarid wasp, *Pseudomasaris vespoides* (Cresson). *The Pan-Pacific Entomologist* 50: 226–234.