

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Harold W. Manter Laboratory: Library Materials

Parasitology, Harold W. Manter Laboratory of

6-22-2023

***Rickettsia africae* and other unclassified *Rickettsia* species of the spotted fever group in ticks of the Western Ghats, India**

N. Naren Babu

Anup Jayaram

Amogh Milind Auti

Yuvraj Bhandari

Ujwal Shetty

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/manterlibrary>



Part of the [Parasitology Commons](#)

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Harold W. Manter Laboratory: Library Materials by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

N. Naren Babu, Anup Jayaram, Amogh Milind Auti, Yuvraj Bhandari, Ujwal Shetty, and Govindakarnavar Arunkumar



Rickettsia africae and other unclassified *Rickettsia* species of the spotted fever group in ticks of the Western Ghats, India

N Naren Babu¹ · Anup Jayaram¹ · Amogh Milind Auti^{1,2} · Yuvraj Bhandari¹ · Ujwal Shetty¹ · Govindakarnavar Arunkumar³

Received: 13 August 2022 / Accepted: 19 June 2023 / Published online: 22 June 2023

© The Author(s) 2023

Abstract

The spotted fever group (SFG) of *Rickettsia* are zoonotic disease-causing pathogens, commonly transmitted by hard ticks to a wide range of hosts, including humans. *Rickettsia conorii* is the common SFG recognised in India, whereas most of the infections due to other group species go undifferentiated at the species level. Hence, this study was conducted to screen host-seeking ticks in the Western Ghats region, India, for the DNA of SFG *Rickettsia*. The ticks were collected from Kerala, Goa, and Maharashtra states of India during a survey conducted between November 2017 and January 2018. In total, 288 tick pools were screened for *Rickettsia* spp. DNA using pan-*Rickettsia* real-time PCR, and conventional PCR targeting the *gltA*, *OmpA* and 17-kDa protein-coding genes. Nucleotide sequences were subjected to phylogenetic analysis using the NCBI BLAST tool to identify submitted sequences with higher homology. Neighbour-joining trees were constructed using the reference sequences of the GenBank database. Overall, *Rickettsia* spp. DNA was detected in 27.2% (62/228 pools) of host-seeking ticks across the Western Ghats region, with an estimated minimum infection rate of 0.057. Upon phylogenetic analysis, it was identified that the detected sequences were highly similar (>99% sequence homology) to *R. africae*, *Candidatus R. laeensis* and an un-categorised *Rickettsia* species, and they were widely carried by *Haemaphysalis* ticks. The current study is the first report of *R. africae* and *Candidatus R. laeensis* in ticks in India. Although the pathogenicity of these species is not well documented, they may pose a potential threat to both animal and the human population in this geographical region.

Keywords *Rickettsiae* · Ixodid ticks · Zoonosis · PCR · Western ghats · India

Introduction

Rickettsia is an obligate intracellular Gram-negative bacterium that causes zoonotic diseases in a wide range of hosts (Raoult and Roux 1997; Weinert et al. 2009). *Rickettsia* species cause mild to severe infections in humans (Raoult and Roux 1997; Parola and Raoult 2001) – Rocky Mountain spotted fever (RMSF), Mediterranean spotted fever (MSF), scrub

Extended author information available on the last page of the article

typhus, and epidemic typhus are a few of the widespread *Rickettsia* infections in humans (Raoult and Roux 1997; Parola and Raoult 2001; Parola et al. 2005, 2013). *Rickettsia prowazekii* and *R. rickettsii* are few among the highly pathogenic rickettsia species that may cause mortality (20–60%) among untreated human cases (Raoult and Roux 1997; Parola and Raoult 2001; Parola et al. 2013).

Many species of Rickettsiae are known for their endosymbiotic relationship with arthropods (Raoult and Roux 1997; Weinert et al. 2009). Among the arthropods carrying *Rickettsia*, ticks were strongly associated with the spotted fever group Rickettsiae (SFGR) (Reháček 1989; Parola and Raoult 2001; Parola et al. 2005, 2013; Socolovschi et al. 2009). While several novel SFGRs are regularly being detected in ticks around the globe (Parola et al. 2013), many are known to cause rickettsiosis in humans (Raoult and Roux 1997; Parola et al. 2013). Some members of the SFGR, such as *R. africae* and *R. massiliae* are a few common tick-borne rickettsioses with high global prevalence. *Rickettsia africae* causes African tick-bite fever (ATBF) transmitted predominantly by *Amblyomma variegatum* among natives and travellers to Africa (Kelly et al. 1996; Raoult and Roux 1997; Jensenius et al. 2004; Parola et al. 2013; Binder and Gupta 2015). *Rickettsia massiliae* causes a Mediterranean spotted fever-like disease through *Rhipicephalus* spp. bite (Raoult and Roux 1997; Cascio et al. 2013; Parola et al. 2013). Both infections may present with fever, rashes, eschars and rarely with regional lymphadenopathy (Kelly et al. 1996; Cascio et al. 2013).

In India, *Rickettsia* infections have been recorded since World War II, and their reports have constantly increased in number over the last decade (Rathi and Rathi 2010; Dasari et al. 2014; Rahi et al. 2015). Infections of scrub typhus group (STG) and spotted fever group (SFG) *Rickettsia* are common across the country, especially in the sub-Himalayan region, Maharashtra, Rajasthan, Punjab and southern states of India (Mahajan et al. 2006; Batra 2007; Rathi and Rathi 2010; Dasari et al. 2014; Rahi et al. 2015). *Rickettsia conorii* transmitted by *Rhipicephalus sanguineus* sensu lato (s.l.) is the common SFGR in India, and it is prevalent in many states of the country (Rathi and Rathi 2010; Dasari et al. 2014), whereas records on other species are scanty. The current study assessed the presence and prevalence of SFGR carried by the host-seeking ixodid ticks of the Western Ghats region in India.

Materials and methods

Study area

Tick pools

The ticks collected from various parts of the Western Ghats (a highly diverged ecosystem of India) during a previous study (Naren Babu et al. 2019) were used for *Rickettsial* screening. In total, 8373 host-seeking ticks were collected (4474 larvae [53.4%], 3719 nymphs [44.4%], and 180 adults [2.2%]) through the flagging method from October 2017 to January 2018 (see details Naren Babu et al. 2019). Ticks of various genera, including *Haemaphysalis*, *Dermacentor*, *Amblyomma*, and *Rhipicephalus*, were identified morphologically. Ticks were collected from Sattari taluk, Goa (n=1020), Dodamarg taluk, Maharashtra (n=411) and Sultan-Bathery taluk, Kerala (n=377). The ticks were pooled based on the site of collection, species and life stage, with a maximum of 15 ticks per pool. Pooled tick sam-

ples were homogenised using TissueLyser II (Qiagen, Valencia, CA, USA) and DNA was extracted using the QIAamp DNA Blood Mini Kit (Qiagen, Hilden, Germany) as per the manufacturer's instructions.

Real-time PCR

The DNA extracted was screened for the presence of *Rickettsia* spp. using pan-*Rickettsia* real-time PCR. The primer, probe, reaction mix and cycling condition were optimised to amplify the 23 S rRNA region of *Rickettsia* as described by a previous study (Kato et al. 2013). The amplification was performed using an ABI 7500 real-time PCR machine (Applied Biosystems, Carlsbad, CA, USA). The reaction sets were validated using at least one internal positive control (IPC), and 2–3 non-template negative controls (using nuclease-free water). Cycle threshold values ≤ 40 were considered a positive cut-off for *Rickettsia*.

Conventional PCR

Further, the positive samples were amplified for *OmpA*, *gltA* and 17-kDa protein-coding genes as previously described (Anderson et al. 1988; Roux et al. 1997; Fournier et al. 1998; Stenos et al. 1998; Chmielewski et al. 2009). The reactions were carried out under the conditions described in Table 1, with standard enzyme activation (95 °C, for 7 min), extension (68 °C, 1 min) and final extension (72 °C, 7 min) using the ProFlex PCR system (Applied Biosystems). All the reaction cycles were conducted in a 25- μ L reaction volume: which includes 10 μ M of primer (1 μ L each, forward and reverse), 10 μ L buffer mix (Ambion Life Technologies, Carlsbad, CA, USA), 1.0 μ L enzyme mix (Ambion Life Technologies) and 7.0 μ L of nuclease-free water (NFW) and 5 μ L of extracted query DNA. Each reaction set was validated using at least one synthetic positive control (referring *R. conorii* str. Malish 7), and 2–3 non-template negative controls (using nuclease-free water). The amplified prod-

Table 1 List of primers and cycling conditions used for the amplification of different *Rickettsia* spp. genes

Gene	Primer name	Primer sequence (5'→3')	Dena- turation (°C/s)	An- nealing (°C/s)	Frag- ment length (bp)
<i>OmpA</i> (Fournier et al. 1998)	Rr. 190.70	ATGGCGAATATTTCTCAAAA	95/20	56/30	630
	Rr. 190.701	GTTCCGTTAATGGCAGCATCT			
<i>gltA</i> (Roux et al. 1997)	RpCS.409d	CCTATGGCTATTATGCTTGC	95/30	45/30	750
	RpCS.1258n	ATTGCAAAAAGTACAGTGAACA			
17-kDa protein- coding gene (Anderson et al. 1988; Stenos et al. 1998)	Rr17.61p	GCTCTTGCAACTTCTATGTT	95/30	50/45	434
	Rr17.492n	CATTGTTTCGTCAGGTTGGCG			
23 S rRNA Pan-Rickett- sia real-time PCR (Kato et al. 2013)	PanR8_F	AGCTTGCTTTTGATCATTGG	95/15	60/60	221
	PanR8_R	TTCTTGCCCTTTTCATACATC TAG T			
	PanR8_P	FI-CCTGCTTCTATTTGTCTT GCAGTAACACGCCA-BHQ1			

ucts were resolved in 1.2% agarose gel, along with a molecular marker of 1 kb, to determine positive samples (Table 1).

Sequencing

The PCR products were excised from the agarose gel post-electrophoresis and purified using the GenElute Gel Extraction Kit (Sigma-Aldrich, St. Louis, MO, USA). Purified products were sequenced with the same primer sets using Big Dye Terminator Cycle Sequencing Kit v.3.1 and 3500 Genetic Analyzer (both Applied Biosystems).

Phylogenetic analysis

The sequences were assembled from raw files using Sequencher v.5.4.6 (Gene Codes Corporation, Ann Arbor, MI, USA) and analysed through the NCBI BLAST tool (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). Further, phylogenetic trees were constructed with the query and *Rickettsia* reference sequences of *OmpA* and 17-kDa protein-coding genes using the Neighbour-joining method and Kimura-2 parameter at 1000 replicates of bootstrap. All the analyses were performed in MEGA v.11.

Statistical analysis

To understand the distribution of *Rickettsia* among the tick population, a minimum infection rate (MIR) and estimated pooled prevalence (EPP) were estimated for each tick species at the site level. The minimum infection rate was estimated as a ratio of the number of positive pools to the total number of ticks tested (Gu et al. 2003). Estimated pooled prevalence was estimated using a Bayesian approach and a Gibbs sampler iterative model by assuming pool size is 15 and an assay of perfect sensitivity and specificity (Cowling et al. 1999).

Results

In total, 1808 ticks belonging to the genera *Haemaphysalis*, *Dermacentor*, *Amblyomma* and *Rhipicephalus* were grouped into 228 pools and screened for *Rickettsia* spp. DNA. Overall, 27.2% (62/228) of the pools were positive for pan-*Rickettsia* real-time PCR (Table 2). Among the *Rickettsia*-positive pools, 74.2% (46/62) were the ticks of immature life stages (i.e., 24 larval and 22 nymphal pools). The minimum infection rate (MIR) of *Rickettsia* in the tick pools was estimated at 0.057, with an adjusted prevalence rate of 0.022 (EPP). In general, *H. turturries* ticks had higher MIR in all the sites (Sattari, nymphs: 0.055; Dodamarg, adults: 0.333; Sultan-Bathery, adults: 0.285, nymphs: 0.375). However, the Bayesian adjusted prevalence estimation (EPP) revealed that *H. bispinosa* nymphs had higher *Rickettsia* spp. prevalence in Sultan-Bathery. Amongst the surveyed sites, Sultan-Bathery ticks showed as high as 0.09 MIR with 0.043 EPP (Table 2). In Sultan-Bathery, 66.7% of the larval population carried *Rickettsia* spp. DNA, and recorded the highest prevalence rate (0.13 EPP) amongst the tick population of all the sites.

Out of the 62 *Rickettsia*-positive pools by real-time PCR, 13 were sequenced for *OmpA*, *gltA* and 17-kDa protein-coding genes, and the nucleotide sequences were submitted to

GenBank under the following accession numbers: MK905239–MK905251 (*OmpA*), MN557213–MN557224 (*gltA*) and MN557225–MN557235 (17-kDa protein-coding gene). Further, the NCBI BLAST analysis identified the two query sequences extracted from *Haemaphysalis* spp. larvae of the population of Sattari taluk, Goa had homologous sequences to *R. africae* (>99% identity). Two other detected *Rickettsia* sequences of Sultan-Bathery taluk, Kerala were homologous to the *Candidatus R. laoensis* (>99% identity). The rest of the nine detected *Rickettsia* (four from Sattari, Goa, and five from Dodamarg, Maharashtra taluks) had homology with an unclassified *Rickettsia* species ZJ43/2007 (>99% identity). Similar *Rickettsia* species have been reported earlier in the Kerala state of India in 2014 as an endosymbiont of *Rhipicephalus haemaphysaloides* (>97% identity).

Phylogenetic analysis showed that sequences in this study, LG07 and LG09 share the same clade with *R. africae*. LW05 and LW15 were found homologous with *Candidatus R. laoensis* (closely formed with *R. raoulti*) and seven other detected *Rickettsia* were found homologous with unclassified *Rickettsia* such as ZJ43/2007 and LOI69 (Figs. 1 and 2) (*OmpA* genes closely formed with *Rickettsia* endosymbiont of *R. haemaphysaloides* and *R. massiliae*).

Further, an additional phylogenetic tree was constructed using the *OmpA* genes of *R. africae* (Fig. S1). The tree formed two distinct clades; most of the sequences of clade-1 contain *R. africae* sequences extracted from *Amblyomma* ticks, whereas clade-2 contains sequences from diverse genera of vector species except *Amblyomma*. The query sequences were grouped in the clade-2, with close identity to an Egyptian strain.

Discussion

The current study identified that SFGR is widely prevalent amongst the host-seeking ixodid ticks across the Western Ghats region in India. The ticks of the Sultan-Bathery site showed the highest MIR of 0.09, whereas the overall MIR was 0.057. Sultan-Bathery hosts half of the positive tick pools in this study. Higher *Rickettsia* positivity among larval tick pools of *Haemaphysalis* spp. is suggestive of the possible transovarial maintenance of these organisms in the tick population of the Western Ghats, India.

The current study provides the first evidence of the presence of *R. africae* in the larval stage of *Haemaphysalis* ticks, the widely prevalent cattle tick of the Western Ghats (Naren Babu et al. 2019). Sequence analysis revealed that the *Rickettsia* detected in this study is closest to the non-*Amblyomma* tick isolate of *R. africae*, and is closely related to the Egyptian strain isolated from *Hyalomma marginatum* (Fig. S1). In concurrence with other studies, *R. africae* is found to be prevalent among cattle ticks (Pillay et al. 2022) and occurs at a higher rate among larval tick population, inferring to possible transovarial maintenance of the species (Mazhetese et al. 2022). Only a few other reports from China had a similar prevalence of *R. africae* among the *Haemaphysalis* tick population (Fig. S1).

Candidatus R. laoensis (an *R. massiliae*-like species) is being reported in the larvae of *Haemaphysalis* spp. ticks of the Wayanad district for the first time in India. *Candidatus R. laoensis* was first and only described from the Nakai District of Laos in a *Haemaphysalis* nymph (Taylor et al. 2016). Additionally, an uncharacterised *Rickettsia* spp. (a novel *R. massiliae*-like species) was identified at multiple sites in the Western Ghats region. Similar species have been reported earlier from *Amblyomma testudinarium* in Laos (Satjanadum-

Table 2 Site-wise positivity (%) of *Rickettsia* spp. among the tick population screened in the Western Ghats, India

Site of tick collection	Tick species and life-stage	Pools tested ¹	Positive pools ²	Positivity (%) ³	MIR ⁴	EPP ⁵ (mean±SD)	
Sattari taluk, Goa	All stages	98	21	21.42	0.02	0.017±0.0036	
	Adult						
		<i>H. spinigera</i>	5	-	-	-	-
		<i>H. turturries</i>	6	-	-	-	-
		<i>H. bispinosa</i>	5	-	-	-	-
		<i>H. minuta</i>	1	-	-	-	-
		<i>H. leachii</i>	2	-	-	-	-
		<i>Rhipicephalus</i> spp.	2	-	-	-	-
	Nymph						
		<i>H. spinigera</i>	24	6	25	0.02	0.022±0.0085
		<i>H. turturries</i>	3	1	33	0.055	0.071±0.15
		<i>H. bispinosa</i>	3	-	-	-	-
		<i>Amblyomma</i> spp.	1	-	-	-	-
		<i>Haemaphysalis</i> spp.	42	14	33.33	0.023	0.029±0.0076
	<i>Amblyomma</i> spp.	4	0	-	-	-	
Dodamarg taluk, Maharashtra	All stages	54	6	11.11	0.014	0.0092±0.0035	
	Adult						
		<i>H. spinigera</i>	7	2	28.57	0.142	0.033±0.02
		<i>H. turturries</i>	4	2	33	0.333	0.1±0.18
		<i>H. bispinosa</i>	2	-	-	-	-
		<i>H. leachii</i>	1	-	-	-	-
		<i>Dermacentor</i> spp.	8	-	-	-	-
		<i>Rhipicephalus</i> spp.	2	-	-	-	-
		<i>Amblyomma</i> spp.	1	-	-	-	-
	Nymph						
		<i>H. spinigera</i>	12	-	-	-	-
		<i>H. intermedia</i>	2	-	-	-	-
	Larvae						
		<i>Haemaphysalis</i> spp.	14	2	14.28	0.006	0.015±0.0091
	<i>Amblyomma</i> spp.	1	-	-	-	-	

Table 2 (continued)

Site of tick collection	Tick species and life-stage	Pools tested ¹	Positive pools ²	Positivity (%) ³	MIR ⁴	EPP ⁵ (mean±SD)	
Sultan-Bathery taluk, Kerala	All stages	76	35	46.05	0.09	0.043±0.0074	
	Adult	<i>H. bispinosa</i>	21	10	50	0.344	0.048±0.015
		<i>H. turturries</i>	5	2	40	0.285	0.051±0.066
	<i>H. intermedia</i>	3	-	-	-	-	
	<i>H. aculeata</i>	1	-	-	-	-	
	<i>H. minuta</i>	1	-	-	-	-	
	<i>H. centropi</i>	1	-	-	-	-	
	<i>H. wellingtoni</i>	1	-	-	-	-	
	Nymph	<i>H. spinigera</i>	5	-	-	-	-
		<i>H. turturries</i>	7	3	42.85	0.375	0.057±0.13
		<i>H. bispinosa</i>	16	12	75	0.088	0.12±0.11
		<i>H. intermedia</i>	2	-	-	-	-
	Larvae	<i>H. cuspidata</i>	1	-	-	-	-
		<i>Haemaphysalis</i> spp.	12	8	66.66	0.044	0.13±0.2
Total	All stages	228	62	27.19	0.057	0.022±0.028	

¹No. tick pools tested for pan-*Rickettsia* real-time PCR.

²No. tick pools producing cycle threshold (1) value of < 35 for pan-*Rickettsia* real-time PCR.

³Percentage of tick pools producing cycle threshold (1) value of < 35 for pan-*Rickettsia* real-time PCR of total tested

⁴Minimum infection rate (ratio of the number of positive pools to the total number of ticks tested)

⁵Estimated pooled prevalence using a Bayesian approach and a Gibbs sampler iterative model (assumed pool size is 15 and an assay of perfect sensitivity and specificity)




Fig. 2 Phylogeny of *Rickettsia* species 17-kDa protein-coding gene sequences (432 bp) extracted from ticks of India (highlighted in bold) in comparison to the reference sequences. The phylogenetic tree was constructed using the neighbor-joining method based on the Kimura 2-parameter, analyzed at 1000 bootstraps (bootstrap values of > 70% are displayed at the nodes). The bar represents the divergence

rong et al. 2019), *R. haemaphysaloides* in Taiwan (Satjanadumrong et al. 2019), and in India. Even though both the rickettsial species are widely prevalent across the ticks of the Western Ghats, further studies are required to determine their pathogenicity to mammals, humans and birds.

The diagnosis of acute febrile illnesses (AFI) has become increasingly challenging due to the constant spill-over of novel zoonotic pathogens into the human population (Parola and

Raoult 2001; Weinert et al. 2009). In Asia, a vast proportion of AFI cases (8–80%) remain undiagnosed, leading to non-specific treatment (Susilawati and McBride 2014; Joshi and Kalantri 2015). Despite the SFGR infections frequently being detected in humans via the Weil-Felix test in India, the infecting species are mostly unknown (Rathi and Rathi 2010; Dasari et al. 2014; Rahi et al. 2015; Narvencar et al. 2017). Knowledge of the prevalence and distribution of *Rickettsial* pathogens is critical for early diagnoses, prompt treatment and disease control (Rathi and Rathi 2010; Dasari et al. 2014). The current report provides evidence for the prevalence of *R. africae*, *Candidatus R. laoensis* and another novel SFG *Rickettsia* among the tick population of India. This is suggestive of the potential risk of transmission of these SFGR to animals and/or humans through tick-bite. These SFGRs might also be the contributing cause for some proportion of pyrexia of unknown origin (PUO) in India. Therefore, accurate identification of the locally prevalent *Rickettsia* species in the tick population could help in improving the diagnosis of PUO.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10493-023-00814-2>.

Acknowledgements The authors acknowledge their sincere gratitude to all the state and district Government health officials of Wayanad in Kerala, North Goa in Goa and Sindhudurg in Maharashtra for their immense support in executing this study. We would also like to extend our gratitude to all the AFI project administrators, site managers, specialists, research assistants, technicians and field support staff for their support in this study. The views expressed in this manuscript are those of the authors and do not necessarily represent the official position of the Manipal Academy of Higher Education.

Author contributions Mr Naren Babu N and Dr Govindakarnavar Arunkumar contributed to the study conception and design. Material preparation, data collection and analysis were performed by Mr Naren Babu N, Mr Anup Jayaram, Mr Ujwal Shetty, Mr Amogh Milind Auti, and Mr Yuvraj Bhandari. The first draft of the manuscript was written by Mr Naren Babu N and Mr Anup Jayaram. Dr Govindakarnavar Arunkumar supervised all work, reviewed and edited the manuscript. All authors read and approved the final manuscript.

Funding This work was supported by the research funds of Manipal Institute of Virology, Manipal Academy of Higher Education, Manipal, India. Open access funding provided by Manipal Academy of Higher Education, Manipal

Data Availability The nucleotide sequences generated during the current study are available in the GenBank repository [GenBank ID for *OmpA*: MK905239 - MK905251, *gltA*: MN557213 -MN557224 and 17-kDa protein-coding gene: MN557225 - MN557235].

Declarations

Competing interests The authors declare no competing interests.

Ethics approval Not Applicable.

Consent to participate Not Applicable.

Consent for publication Not Applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material.

If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Anderson BE, Baumstark BR, Bellini WJ (1988) Expression of the gene encoding the 17-kilodalton antigen from *Rickettsia rickettsii*: transcription and posttranslational modification. *J Bacteriol* 170(10):4493–4500
- Batra HV (2007) Spotted fevers & typhus fever in Tamil Nadu. *Indian J Med Res* 126(2):101–103
- Binder WD, Gupta R (2015) African tick-bite fever in a returning traveler. *J Emerg Med* 48(5):562–565. <https://doi.org/10.1016/j.jemermed.2014.12.044>
- Cascio A, Torina A, Valenzise M, Blanda V, Camarda N, Bombaci S, Iaria C, De Luca F, Wasniewska M (2013) Scalp eschar and neck lymphadenopathy caused by *Rickettsia massiliae*. *Emerg Infect Dis* 19(5):836–837. <https://doi.org/10.3201/eid1905.121169>
- Chmielewski T, Podsiadly E, Karbowski G, Tylewska-Wierzbanowska S (2009) *Rickettsia* spp. in ticks, Poland. *Emerg Infect Dis* 15(3):486–488. <https://doi.org/10.3201/eid1503.080711>
- Cowling DW, Gardner IA, Johnson WO (1999) Comparison of methods for estimation of individual-level prevalence based on pooled samples. *Prev Vet Med* 39(3):211–225. [https://doi.org/10.1016/s0167-5877\(98\)00131-7](https://doi.org/10.1016/s0167-5877(98)00131-7)
- Dasari V, Kaur P, Murhekar MV (2014) Rickettsial disease outbreaks in India: a review. *Ann Trop Med Public Health* 7(6):249. <https://doi.org/10.4103/1755-6783.155018>
- Fournier PE, Roux V, Raoult D (1998) Phylogenetic analysis of spotted fever group rickettsiae by study of the outer surface protein rOmpA. *Int J Syst Bacteriol* 48 Pt 3:839–849. <https://doi.org/10.1099/00207713-48-3-839>
- Gu W, Lampman R, Novak RJ (2003) Problems in estimating mosquito infection rates using minimum infection rate. *J Med Entomol* 40(5):595–596. <https://doi.org/10.1603/0022-2585-40.5.595>
- Jensenius M, Fournier P-E, Raoult D (2004) Rickettsioses and the international traveler. *Clin Infect Dis Off Publ Infect Dis Soc Am* 39(10):1493–1499. <https://doi.org/10.1086/425365>
- Joshi R, Kalantri SP (2015) Acute undifferentiated fever: management algorithm. *Update on Tropical Fever*, pp 1–14
- Kato CY, Chung IH, Robinson LK, Austin AL, Dasch GA, Massung RF (2013) Assessment of real-time PCR assay for detection of *Rickettsia* spp. and *Rickettsia rickettsii* in banked clinical samples. *J Clin Microbiol* 51(1):314–317. <https://doi.org/10.1128/JCM.01723-12>
- Kelly PJ, Beati L, Mason PR, Matthewman LA, Roux V, Raoult D (1996) *Rickettsia africae* sp. nov., the etiological agent of african tick bite fever. *Int J Syst Bacteriol* 46(2):611–614. <https://doi.org/10.1099/00207713-46-2-611>
- Mahajan SK, Kashyap R, Kanga A, Sharma V, Prasher BS, Pal LS (2006) Relevance of Weil-Felix test in diagnosis of scrub typhus in India. *J Assoc Physicians India* 54:619–621
- Mazhetese E, Lukanji Z, Byaruhanga C, Neves L, Morar-Leather D (2022) *Rickettsia africae* infection rates and transovarial transmission in *Amblyomma hebraeum* ticks in Mnisi, Bushbuckridge, South Africa. *Exp Appl Acarol* 86:407–418. <https://doi.org/10.1007/s10493-022-00696-w>
- Naren Babu N, Jayaram A, Hemanth Kumar H, Pareet P, Pattanaik S, Auti AM, Abdulmajeed J, Maity H, Devadiga S, Bhandari Y, Agre Deepchand H, Shakir M, Kumar N, Arunkumar G (2019) Spatial distribution of *Haemaphysalis* species ticks and human Kyasanur Forest Disease cases along the western ghats of India, 2017–2018. *Exp Appl Acarol* 77(3):435–447. <https://doi.org/10.1007/s10493-019-00345-9>
- Narvencar K, Kaur G, Rodrigues S (2017) Rickettsial infections in Goa-Not just scrub Typhus! *J Assoc Physicians India* 65(8):24–27
- Parola P, Raoult D (2001) Ticks and tickborne bacterial diseases in humans: an emerging infectious threat. *Clin Infect Dis Off Publ Infect Dis Soc Am* 32(6):897–928. <https://doi.org/10.1086/319347>
- Parola P, Paddock CD, Raoult D (2005) Tick-borne rickettsioses around the world: emerging diseases challenging old concepts. *Clin Microbiol Rev* 18(4):719–756. <https://doi.org/10.1128/CMR.18.4.719-756.2005>
- Parola P, Paddock CD, Socolovschi C, Labruna MB, Mediannikov O, Kernif T, Abdad MY, Stenos J, Bitam I, Fournier P-E, Raoult D (2013) Update on tick-borne rickettsioses around the world: a geographic approach. *Clin Microbiol Rev* 26(4):657–702. <https://doi.org/10.1128/CMR.00032-13>
- Pillay A, Manyangadze T, Mukaratirwa S (2022) Prevalence of *Rickettsia africae* in tick vectors collected from mammalian hosts in sub-saharan Africa: a systematic review and meta-analysis. *Ticks Tick Borne Dis* 13(4):101960. <https://doi.org/10.1016/j.ttbdis.2022.101960>

- Rahi M, Gupte MD, Bhargava A, Varghese GM, Arora R (2015) DHR-ICMR guidelines for diagnosis & management of rickettsial diseases in India. *Indian J Med Res* 141(4):417–422. <https://doi.org/10.4103/0971-5916.159279>
- Raoult D, Roux V (1997) Rickettsioses as paradigms of new or emerging infectious diseases. *Clin Microbiol Rev* 10(4):694–719
- Rathi N, Rathi A (2010) Rickettsial infections: indian perspective. *Indian Pediatr* 47(2):157–164. <https://doi.org/10.1007/s13312-010-0024-3>
- Reháček J (1989) Ecological relationships between ticks and rickettsiae. *Eur J Epidemiol* 5(4):407–413. <https://doi.org/10.1007/bf00140130>
- Roux V, Rydkina E, Eremeeva M, Raoult D (1997) Citrate synthase gene comparison, a new tool for phylogenetic analysis, and its application for the rickettsiae. *Int J Syst Bacteriol* 47(2):252–261. <https://doi.org/10.1099/00207713-47-2-252>
- Satjanadumrong J, Robinson MT, Hughes T, Blacksell SD (2019) Distribution and ecological drivers of spotted Fever Group Rickettsia in Asia. *EcoHealth* 16(4):611–626. <https://doi.org/10.1007/s10393-019-01409-3>
- Socolovschi C, Mediannikov O, Raoult D, Parola P (2009) The relationship between spotted fever group Rickettsiae and ixodid ticks. *Vet Res* 40(2):34. <https://doi.org/10.1051/vetres/2009017>
- Stenos J, Roux V, Walker D, Raoult D (1998) Rickettsia honei sp. nov., the aetiological agent of Flinders Island spotted fever in Australia. *Int J Syst Bacteriol* 48 Pt 4:1399–1404. <https://doi.org/10.1099/00207713-48-4-1399>
- Susilawati TN, McBride WJH (2014) Acute undifferentiated fever in Asia: a review of the literature. *Southeast Asian J Trop Med Public Health* 45(3):719–726
- Taylor AJ, Vongphayloth K, Vongsouvath M, Grandadam M, Brey PT, Newton PN, Sutherland IW, Dittrich S (2016) Large-scale survey for tickborne Bacteria, Khammouan Province, Laos. *Emerg Infect Dis* 22(9):1635–1639. <https://doi.org/10.3201/eid2209.151969>
- Weinert LA, Werren JH, Aebi A, Stone GN, Jiggins FM (2009) Evolution and diversity of Rickettsia bacteria. *BMC Biol* 7:6. <https://doi.org/10.1186/1741-7007-7-6>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

N Naren Babu¹ · Anup Jayaram¹ · Amogh Milind Auti^{1,2} · Yuvraj Bhandari¹ · Ujwal Shetty¹ · Govindakarnavar Arunkumar³

✉ N Naren Babu
naren.babu@manipal.edu

✉ Govindakarnavar Arunkumar
arunviro@gmail.com

¹ Manipal Institute of Virology, Manipal Academy of Higher Education (Institution of Eminence Deemed to be University), Manipal, India

² Present address: Department of Precision Medicine, University of Campania 'Luigi vanvitelli', Naples, Italy

³ II-49, Vaikathu, Marotithota Road, Mooduathrady, Udipi District, Athrady 576107, Karnataka, India