

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Anthony F. Starace Publications

Research Papers in Physics and Astronomy

---

8-21-2001

## Static-electric-field behavior in negative ion detachment by an intense, high-frequency laser field

M. V. Frolov

*Voronezh State University, Russia*

N. L. Manakov

*Voronezh State University, manakov@phys.vsu.ru*

B. Borca

*University of Nebraska - Lincoln*

Anthony F. Starace

*University of Nebraska-Lincoln, astarace1@unl.edu*

Follow this and additional works at: <https://digitalcommons.unl.edu/physicsstarace>



Part of the [Physics Commons](#)

---

Frolov, M. V.; Manakov, N. L.; Borca, B.; and Starace, Anthony F., "Static-electric-field behavior in negative ion detachment by an intense, high-frequency laser field" (2001). *Anthony F. Starace Publications*. 150.  
<https://digitalcommons.unl.edu/physicsstarace/150>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Anthony F. Starace Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Submitted June 12, 2001; revised August 6, 2001; published August 21, 2001.

LETTER TO THE EDITOR

## Static-electric-field behavior in negative ion detachment by an intense, high-frequency laser field

M. V. Frolov,<sup>1</sup> N. L. Manakov,<sup>1</sup> B. Borca,<sup>2</sup> and  
A. F. Starace<sup>2</sup>

<sup>1</sup> Department of Physics, Voronezh State University, Voronezh  
394693, Russia

<sup>2</sup> Department of Physics and Astronomy, University of Nebraska-  
Lincoln, Lincoln, NE 68588-0111, USA

### Abstract

Based upon the exact numerical solution of the complex quasienergy problem for a 3-dimensional short-range potential as well as upon analytical evaluations, we demonstrate for any finite frequency  $\omega$  that the action of an ultra-intense laser field (with electric vector  $F(\omega t)$ ) on a weakly bound atomic system may be described by the cycle-averaging of results for an instantaneous static electric field of strength  $|F(\omega t)|$ .

The accurate description of the intensity dependence of the decay rate of a bound level over a broad interval of laser frequencies is one of the challenging problems of strong field laser-atom physics. Existing qualitative results obtained from nonperturbative (in the intensity) analyses of atomic decay rates in a laser field depend significantly on the relation between the laser frequency  $\omega$  and  $\omega_0 = |E_0|/\hbar$  (where  $E_0$  is the binding energy), as well as that between the laser amplitude  $F$  (see (2) below) and the characteristic internal atomic field strength,  $F_0 = (2m|E_0|^3/|e|\hbar)^{1/2}$ . (Below we use the following scaled units: energies,  $\omega$  and  $F$  are measured in units  $|E_0|$ ,  $\omega_0$ , and  $F_0$ , respectively.) For small frequencies,  $\omega \ll 1$ , and for field strengths  $F \geq \omega$  (or equivalently for  $\gamma_K < 1$ , where  $\gamma_K = \omega/F$  is the well-known Keldysh parameter), the tunneling mechanism for the decay is realized, which is valid only for weak (although nonperturbative) fields,  $F \ll 1$  (see [1] and the improved analyses in [2]). The tunneling mechanism for the decay has been confirmed by many experiments for frequencies up to  $\omega \sim (0.1\text{--}0.2)$ , particularly for the rare gases [3]. For the case of ground state atomic hydrogen, H(1s), Pont *et al.* [4, 5] performed a low-frequency analysis of the decay rate  $\Gamma$  beyond the Keldysh approach (up to  $F \leq 0.2$ ) using the  $\omega^2$  expansion of the complex quasienergy using the basis of quasistationary states of the hydrogen atom in a static electric field (whose magnitude equals that of the instantaneous laser field, see below). For  $\omega = 0.134$  ( $\lambda = 616$  nm), a comparison of the  $F$  dependence of these “static-field-based” results with the exact ones shows a reasonable agreement (which becomes better for stronger  $F$ ) except for the structure seen in the exact  $\Gamma(F)$  which is due to Rydberg levels shifting in and out of resonance as the intensity varies. With increasing  $F$  (e.g. for  $F \geq 0.2$  in the case of H(1s)), over-barrier ionization becomes important. Recently, the over-barrier decay rate  $\Gamma$  in the low-frequency limit,  $\omega \ll 1$ ,

has been obtained by Popov [6] using an adiabatic cycle-averaging of the Stark width  $\Gamma_{\text{stat}}$  for a strong static electric field. It demonstrates a surprisingly linear dependence of  $\Gamma$  on  $F$  (the “intermediate” asymptotic regime [6]),

$$\Gamma \approx k(F - F_{\text{cr}}) \quad (1)$$


where the fitting parameters  $k$  and  $F_{\text{cr}}$  do not depend on  $F$  over a wide fitting interval (e.g.  $0.6 < F < 2$  for H(1s)) and are smooth functions of the laser ellipticity. For above-threshold frequencies,  $\omega > 1$ , and in the strongly nonperturbative regime, the concept of quasistationary stabilization of atomic decay rates is conventionally understood to be applicable, in which case  $\Gamma(F)$  has a decreasing trend with increasing  $F$  (see reviews [7, 8] on the recent status of this problem). However, for a Coulomb potential the existence of a stabilization regime for decay rates in the ultra-strong field limit is still an open question.

The analysis of  $\Gamma(F, \omega)$  is simplified for the case of negative ion detachment, for which simple analytic, short-range binding potentials can be utilized. One of them is the 3-dimensional zero-range potential (ZRP) that has been widely used for the description of a weakly bound electron, as for example in the  $\text{H}^-$  negative ion. The use of a quasistationary quasienergy state (QQES) approach [9] for the ZRP model essentially permits exact predictions of  $\Gamma(F, \omega)$  (which is determined by the imaginary part of the complex quasienergy,  $\epsilon = \text{Re } \epsilon - i\Gamma/2$ ) for laser intensities extending from the perturbative to the ultrahigh intensity regime and for frequencies extending from the tunneling to the multiphoton regimes. In particular, recently we have demonstrated [10] that, for the ZRP model, the stabilization-like behavior of  $\Gamma(F)$  in a high-frequency field only exists for a limited interval of  $F$ , up to the closing of the direct photoionization channel caused by the ponderomotive shift. Moreover, for the particular case of circular polarization, the strong field behavior of  $\Gamma(F)$  was found to be similar to that for a strong static electric field, both for  $\omega < 1$  as well as for the post-stabilization regime at  $\omega > 1$ .

In this letter we present a global analysis of the dependence of  $\Gamma$  (for the ZRP model) on  $F$ ,  $\omega$ , and on the polarization state of a laser field described by the electric vector

$$\mathbf{F}(\omega t) = \frac{F}{\sqrt{1 + \eta^2}} \{\cos \omega t, \eta \sin \omega t, 0\}, \quad -1 \leq \eta \leq +1. \quad (2)$$

(Instead of the ellipticity,  $\eta$ , it is more convenient in what follows to use the related degree of linear polarization,  $l = (1 - \eta^2)/(1 + \eta^2)$ .) For details concerning the exact numerical calculations of the complex quasienergy  $\epsilon$  for the ZRP in the nonperturbative regime, see [10, 11]. The method we employ gives results that are in agreement with those of other authors who employ the ZRP, e.g. [12]. Results of exact numerical calculations for  $\Gamma(F)$  are presented in Figure 1 for four different values of  $l$  and for  $\omega = 1.5$ , which corresponds to the case of  $\text{H}^-$  irradiated by a Nd:YAG laser. (These results cover a much greater range of  $F$  and  $l$  than in [10].) One observes that as  $F$  increases, the perturbative regime, in which  $\Gamma \sim F^2$ , evolves smoothly into a stabilization-like behavior, which breaks up at the closure of the one-photon ionization channel, i.e. at  $F = F_{\text{th}}^{(1)}$ . Note that the finite value of  $\Gamma$  at  $F = F_{\text{th}}^{(1)}$  results from the contributions of partial rates  $\Gamma^{(n)}$  for  $n$ -photon (above-threshold) detachment with  $n = 2, 3, 4, \dots$ , whose  $F$ -dependence (for  $n > 2$ ) is essentially perturbative for  $F \sim F_{\text{th}}^{(1)}$ . It is also seen that the threshold structure of  $\Gamma(F)$  at higher thresholds is significantly different from that for  $n = 1$  and depends sensitively on the laser polarization. The frequency dependence of  $\Gamma$  in the interval  $0.15 < \omega < 2$  is presented in Figure 2 for  $l = 0.72$  for four different values of  $F$ . For weak  $F$ ,  $\Gamma(\omega)$  exhibits the typical perturbative behavior, i.e. the step-like increase as  $\omega$  increases that results from the sequential contributions of the partial



**Figure 1.**  $F$ -dependence of the total detachment rate  $\Gamma$  for  $\omega = 1.5$ . Full curve, QUES results for four different values of  $l$ , as indicated in the figure; open circles, the lowest-order perturbation theory (PT) result for  $\Gamma^{(1)} \sim F^2$ ; full circles, PT result for  $\Gamma^{(1)} + \Gamma^{(2)}$  (including terms up to the order of  $F^4$ ) for  $l = 0.72$ .

rates,  $\Gamma^{(n)} \sim F^{2n}$ , with  $n$  becoming smaller as  $\omega$  increases. As  $F$  increases, the stair-step behavior gradually disappears as  $\Gamma(\omega)$  nearly becomes insensitive to  $\omega$  for essentially nonperturbative values of  $F$ . This unusual behaviour of  $\Gamma(\omega)$  at high  $F$  allows us to assume that in the strong field limit the decay mechanism itself becomes essentially independent of the frequency, even in the  $\omega > 1$  domain.

To analyze the strong field regime in more detail, instead of the conventional representation for a quasienergy state,  $\Psi_\epsilon(r, t) = \Phi_\epsilon(r, t)\exp(-i\epsilon t)$ , we use the following one:


$$\Psi_\epsilon(r, t) = \chi(r, t) \exp \left( -i \int^t \mathcal{E}(t') dt' \right) \tag{3}$$

where  $\Psi_\epsilon(r, t)$  is the solution of the Schrödinger equation for a Hamiltonian  $H(r, t) = H_{\text{at}}(r) + V(r, t)$ , where  $H_{\text{at}}(r)$  describes the atom and  $V(r, t) = r \cdot F(\omega t)$ . The periodic functions  $\chi(r, t)$  and  $\mathcal{E}(t)$  satisfy the following equation

$$\left( H_{\text{at}}(r) + r \cdot F(\omega t) - \mathcal{E}(t) - i \frac{\partial}{\partial t} \right) \chi(r, t) = 0. \tag{4}$$

One may easily verify that the quasienergy  $\epsilon$  is the cycle-average of  $\mathcal{E}(t)$ ,

$$\epsilon = \frac{1}{T} \int_0^T \mathcal{E}(t) dt, \quad T = 2\pi/\omega. \tag{5}$$



**Figure 2.**  $\omega$ -dependence of the total detachment rate  $\Gamma$  for four values of  $F$ , as indicated in the figure, and for  $l=0.72$ .

Equations (3)–(5) are very general and were used by Langhoff *et al.* [13] in their analyses of so-called “secular terms” in higher orders of perturbation theory (in  $V$ ), and by Pont *et al.* [5] in the low-frequency analysis of the ionization of H(1s). In [5] the formal development of a perturbation theory in  $W = -i\omega\partial/\partial\tau$ , where  $\tau = \omega t$ , is presented for calculations of  $\chi(r, t)$  and  $\mathcal{E}(t)$  based on the instantaneous state of an atom in a static electric field of strength  $\mathcal{F} = |F(\omega t)|$ ,  $\chi^{(0)}(r, t)$ , with energy  $\mathcal{E}^{(0)}(t)$ . In what follows, we employ such an approach to analyze the frequency dependence of  $\epsilon$  for the ZRP model in the strong field limit. Since we do not assume that  $\omega$  is small compared to the binding energy  $|E_0|$ , the key issue is to calculate the next order correction,  $\mathcal{E}^{(2)}(t) \sim \omega^2$ , to  $\mathcal{E}^{(0)}(t)$  in order to estimate the accuracy of the expansion of  $\epsilon$  in a power series in  $\omega^2$ , which is generally an asymptotic expansion [5].

The general result for  $\mathcal{E}^{(2)}(t)$  is [5],

$$\mathcal{E}^{(2)}(t) = \omega^2 \left\langle \frac{\partial \tilde{\chi}^{(0)}(\mathbf{r}, t)}{\partial \tau} \left| \mathcal{G}'_{\mathcal{E}^{(0)}(t)}(\mathbf{r}, \mathbf{r}') \right| \frac{\partial \chi^{(0)}(\mathbf{r}', t)}{\partial \tau} \right\rangle \quad (6)$$

where  $\mathcal{G}'_{\mathcal{E}^{(0)}(t)}(\mathbf{r}, \mathbf{r}')$  is the reduced Green function of an atom in a static electric field and  $\tilde{\chi}^{(0)}(\mathbf{r}, t)$  is the “dual” function,  $\tilde{\chi}^{(0)}(\mathbf{r}, t) = \chi^{(0)*}(\mathbf{r}, -t)$ , which is necessary to provide a proper normalization of the quasistationary (resonance) state  $\chi^{(0)}(\mathbf{r}, t)$  [5, 14]. In the ZRP model (see the review [14] for details),  $\mathcal{E}^{(0)}(t)$  at any fixed  $t$  can be obtained as the root of the transcendental equation:

$$1 + \pi \mathcal{F}^{1/3} J(\xi) = 0 \quad (7)$$

where  $\xi = -\mathcal{E}^{(0)}(t) \mathcal{F}^{-2/3}$ ,  $\mathcal{F} \equiv |F(\omega t)| = F [(1 + l \cos 2\omega t)/2]^{1/2}$ , and  $J(\xi)$  is a combination of regular ( $Ai$ ) and irregular ( $Bi$ ) Airy functions and their derivatives:

$$J = Ai'(\xi) Bi'(\xi) - \xi Ai(\xi) Bi(\xi) + i \left[ Ai'^2(\xi) - \xi Ai^2(\xi) \right].$$

Using the explicit forms of  $\chi_0(r, t)$  and  $\mathcal{G}'_{\mathcal{E}^{(0)}(t)}(r, r')$ , the matrix element in (6) is calculated analytically (some details regarding the calculation of the integrals that occur can be found in [14]):

$$\begin{aligned} \mathcal{E}^{(2)}(\tau) = & -\omega^2 \left[ \left( \frac{\partial \mathbf{F}(\tau)}{\partial \tau} \right)^2 I^{(4)} \right] \left( 360 \mathcal{F}^{8/3} I \right)^{-1} + \frac{\omega^2}{8 \mathcal{F}^2 I} \left( \frac{\partial \mathbf{F}(\tau)}{\partial \tau} \cdot \mathbf{F}(\tau) \right)^2 \\ & \times \left\{ \frac{I^{(1)}}{I^2} \left( I^{(1)} f(\tau) + \frac{I^{(3)}}{6 \mathcal{F}^{4/3}} \right)^2 - \frac{1}{I} \left( I^{(1)} f(\tau) + \frac{I^{(3)}}{6 \mathcal{F}^{4/3}} \right) \left( \frac{4}{3} I^{(2)} f(\tau) + \frac{I^{(4)}}{5 \mathcal{F}^{4/3}} \right) \right. \\ & \left. + \frac{1}{3} I^{(3)} f^2(\tau) + \frac{4 I^{(5)}}{45 \mathcal{F}^{4/3}} f(\tau) + \frac{I^{(7)}}{180 \mathcal{F}^{8/3}} \right\} \end{aligned} \quad (8)$$

where

$$I(\xi) = -J'(\xi) = Ai(\xi)(Bi(\xi) + iAi(\xi)), \quad I^{(n)} = \frac{\partial^n I(\xi)}{\partial \xi^n}.$$

The function  $f(\tau)$  is connected with the derivative of  $\mathcal{E}^{(0)}(t)$ , which is calculated with the use of (7):

$$\frac{\partial \mathcal{E}^{(0)}}{\partial \tau} = \left( \frac{\partial \mathbf{F}(\tau)}{\partial \tau} \cdot \mathbf{F}(\tau) \right) f(\tau), \quad f(\tau) = \frac{1}{3 \mathbf{F}^2(\tau)} \left[ 2\mathcal{E}^{(0)} + \left( \frac{\pi I}{\mathcal{F}^{1/3}} \right)^{-1} \right].$$

The result (8) simplifies for the case of a circularly polarized laser field. In this case,

$$\left( \frac{\partial \mathbf{F}}{\partial \tau} \right)^2 = \frac{F^2}{2} \quad \text{and} \quad \left( \frac{\partial \mathbf{F}}{\partial \tau} \cdot \mathbf{F} \right) = -\left( \frac{1}{2} \right) l F^2 \sin(2\tau) = 0.$$

Thus,  $\mathcal{E}(t)$  is time-independent and the correction  $\epsilon^{(2)}$  is


$$\epsilon_{\text{circ}}^{(2)} = -\frac{\omega^2}{360 \mathcal{F}^{2/3}} \frac{I^{(4)}}{I}. \quad (9)$$

This result coincides with that obtained by an alternative approach in [10], in which the calculations are carried out in a coordinate frame rotating with frequency  $\omega$  (see also the similar calculations for H(1s) in [4]). In [10] an analytical result for the asymptotic behavior of  $\epsilon_{\text{circ}}$  in ultra-strong fields,  $F \gg 1$ , has been obtained. In the weak field limit ( $F \ll 1$ ), neglecting exponentially small (tunneling) terms, we obtain the following result for  $\epsilon = \epsilon^{(0)} + \epsilon^{(2)}$ :

$$\epsilon = -1 - \frac{F^2}{32} \left( 1 + \frac{3F^2}{4} \left( 1 + \frac{l^2}{2} \right) + \frac{7}{24} \omega^2 \left[ 1 + \frac{13}{2} F^2 \left( 1 + \frac{25}{28} l^2 \right) \right] \right). \quad (10)$$

Note that the Stark-shift in this equation coincides exactly with the first two terms of the . expansion for the known dynamic polarizability and hyperpolarizability of a weakly bound particle in the ZRP model [15]. Thus, for weak fields, the “zero approximation,”  $\epsilon \simeq \epsilon^{(0)}$ , is valid for  $\omega \ll 1$  and is equivalent to the standard adiabatic approach. To establish the accuracy of the term  $\epsilon^{(0)}$  for the strong field regime, in Figure 3 we present numerical results for real and imaginary parts of the ratio of  $\epsilon^{(2)}$  to  $\epsilon^{(0)}$  for a number of values of  $l$  at fixed  $\omega = 1.5$ . One observes that with increasing  $F$  the two-term approximation,  $\epsilon^{(0)} + \epsilon^{(2)}$  (which we call the AA result), is applicable over a wide interval of  $\omega$  including the above-threshold region,  $\omega > 1$ .

To check both the relation between the AA results and exact numerical results for  $\epsilon$  and also the applicability of the ZRP model for real negative ions in a strong laser field, in Table 1 we compare our numerical (QQES) and approximate (AA) results for the detachment of  $\text{H}^-$  by linearly polarized  $\text{CO}_2$  laser radiation (for which  $\omega = 0.155$ , and the scaled unit of intensity for  $\text{H}^-$  is  $1.494 \times 10^{12} \text{ W cm}^{-2}$ ) with existing theoretical predictions in [16–19]. The comparison shows the excellent agreement of the exact



**Figure 3.**  $F$  and  $l$  dependences of the real (a) and imaginary (b) parts of the ratio  $\epsilon^{(2)}/\epsilon^{(0)}$  for  $\omega = 1.5$ . Full curve,  $l = 0$ ; long-dashed curve,  $l = 0.5$ ; chain curve,  $l = 0.7$ ; short-dashed curve,  $l = 0.9$ .

ZRP results with more refined (and time consuming) calculations and also the high accuracy of the AA results for nonperturbative intensities  $I \geq 5 \times 10^{10} \text{ W cm}^{-2}$ , when  $F \geq \omega$  (in scaled units).


Comparisons of  $\epsilon_{\text{QES}}$  and  $\epsilon_{\text{AA}}$  as functions of  $F$  are presented in Figure 4 for  $\omega < 1$  and in Figure 5 for  $\omega > 1$ . The AA and QES results for  $l = 0$  and  $\omega < 1$  are almost indistinguishable: for  $\omega = 0.36$  and  $F > 0.3$ , the difference between  $\epsilon_{\text{QES}}$  and  $\epsilon_{\text{AA}}$  is less than 3%; for  $\omega = 0.56$  and  $F > 0.4$ , it is less than 2%; and for  $\omega = 0.77$  and  $F > 0.5$ , the difference is less than 4%. Generally, the AA results accurately describe the trends of the position and the width of a quasistationary level but fail to describe the threshold related peculiarities, which are lost by using the  $\omega^2$  expansion for the iterative solution of equation (4). These peculiarities are most pronounced for the case of linear polarization and they are exhibited at the points of non-analyticity of the function  $\epsilon(F)$ , which correspond to the closure of partial detachment channels with increasing  $F$  (at  $F = F_{\text{th}}^{(n)}$ ). These points are branch points of the type  $(\epsilon + U_p + n\omega)^{k+1/2}$  (where  $U_p$  is the ponderomotive shift,  $U_p = F^2/(2\omega^2)$ ) and as  $F$  increases (and  $\text{Im } \epsilon$  becomes impor-

**Table 1.** Detachment rates for  $H^-$  in the field of a  $CO_2$  laser having linear polarization ( $(n) \equiv 10^n$ ).


Intensity (W cm <sup>-2</sup> )	Detachment rates (au)					
	[16]	[17]	[18]	[19]	AA	QQES
1.0(10)	(1.04 ± 0.12)(-9)	0.97(-9)	0.91(-9)		0.32(-9)	0.97(-9)
1.12(10)	(2.04 ± 0.11)(-9)			2.7(-9) <sup>a</sup>	0.73(-9)	2.28(-9)
				2.1(-9) <sup>b</sup>		
2.52(10)	(1.12 ± 0.08)(-7)			1.4(-7) <sup>a</sup>	0.88(-7)	1.14(-7)
				1.0(-7) <sup>b</sup>		
5.0(10)	(1.81 ± 0.06)(-6)	1.67(-6)	1.76(-6)		1.64(-6)	1.79(-6)
1.0(11)	(1.68 ± 0.03)(-5)	1.61(-5)	1.61(-5)		1.62(-5)	1.66(-5)
1.6(11)	(5.91 ± 0.02)(-5)				5.74(-5)	6.12(-5)
2.0(11)	(9.97 ± 0.01)(-5)				9.75(-5)	9.87(-5)

<sup>a</sup>Floquet calculations with a parametrized one-electron potential.

<sup>b</sup>Faisal-Reiss formulas with a Hylleraas ground state wavefunction.



**Figure 4.**  $F$ -dependence of the real (a) and imaginary (b) parts of the complex quasienergy ( $\epsilon = \text{Re } \epsilon - i\Gamma/2$ ) for  $\omega = 0.36$  and  $l = 1$ . Full curve, the exact QQES result; dashed curve, the AA result.



**Figure 5.**  $F$ -dependence of  $\Gamma$  for  $\omega = 1.5$ , and  $l = 0$  (a) and  $l = 1$  (b). Full curve, the exact QQES result; dashed curve, the AA result.



tant) they are shifted to the complex  $F$  plane. Thus, in strong fields the peculiarities of  $\epsilon(F)$  on the real  $F$  axis become smoother. As Figures 4 and 5 demonstrate, in the strong field limit, the behavior of the exact results for  $\epsilon(F)$  (when averaged over the threshold peculiarities) show surprisingly close coincidence with the AA results, even in the high-frequency domain,  $\omega > 1$ . Moreover, over a wide interval of nonperturbative values of  $F$  the  $F$ -dependence of  $\Gamma$  (averaged over threshold peculiarities) is close to linear, which is similar to the “intermediate” asymptotic (1) found for the hydrogen atom in the low-frequency limit. For instance, at  $\omega = \omega_{\text{CO}_2}$  (see Table 1), the parameters for this linear dependence are  $F_{\text{cr}} = 0.86$  and  $k = 0.12$  for  $l = 1$ , and results obtained from formula (1) are in reasonable agreement with the exact ones beginning from  $F > 1.5$  (or for  $I > 2.25 \times 10^{12} \text{ W cm}^{-2}$  for  $\text{H}^-$ ). Unlike the adiabatic case ( $\omega \ll 1$ ), for a finite frequency the interval of the applicability of the asymptotic (1) depends on  $\omega$ : as  $\omega$  increases, the result (1) becomes applicable at stronger fields. Namely, for  $\omega = 1.5$  (when the parameters  $k$  and  $F_{\text{cr}}$  in (1) are  $F_{\text{cr}} = 0.84$ ,  $k = 0.13$  for  $l = 0$ , and  $F_{\text{cr}} = 0.89$ ,  $k = 0.1165$  for  $l = 1$ ) the linear in  $F$  regime is realized with an accuracy of about 5% over the interval  $2.5 < F < 10$ .

In conclusion, the results presented in this letter justify our key conceptual statement, namely, that the decay of a weakly bound atomic system in a strong laser field  $F(\omega t)$  with any frequency and polarization state may be described by cycle-averaging the results for an instantaneous static electric field of strength  $|F(\omega t)|$ .

#### Acknowledgments

This work was supported by RFBR Grant number 00-02-17843, by Grant number E00-3.2-515 of the Russian Ministry of Education and by NSF Grant number PHY-0070980.

#### References

- [1] Keldysh L V 1964 *Zh. Exp. Teor. Fiz.* **47** 1945 (Engl. transl. 1964 *Sov. Phys.-JETP* 20 1307)
- [2] Nikishov A I and Ritus V I 1966 *Zh. Exp. Teor. Fiz.* **50** 255 (Engl. transl. 1966 *Sov. Phys.-JETP* **23** 168)  
Perelomov A M, Popov V S and Terent'ev M V 1966 *Zh. Exp. Teor. Fiz.* **50** 1393 (Engl. transl. 1966 *Sov. Phys.-JETP* **23** 1924)
- [3] Delone N B and Krainov V P 1994 *Multiphoton Processes in Atoms* (Berlin: Springer)
- [4] Pont M, Shakeshaft R, and Potvliege R M 1990 *Phys. Rev. A* **42** 6969
- [5] Pont M, Potvliege R M, Shakeshaft R, and Teng Z-J 1992 *Phys. Rev. A* **45** 8235
- [6] Popov V S 2000 *Zh. Exp. Teor. Fiz.* **118** 56 (Engl. transl. 2000 *Sov. Phys.-JETP* **91** 48)
- [7] Gavrilin M 2000 *8th Int. Conf. on Multiphoton Processes* (AIP Conf. Proceedings 525) ed L F DiMauro, R R Freeman, and K C Kulander (New York: Melville) p. 103
- [8] Fedorov M V 1999 *Laser Phys.* **9** 209
- [9] Manakov N L, Ovsiannikov V D, and Rapoport L P 1986 *Phys. Rep.* **141** 319  
Manakov N L and Fainstein A G 1980 *Zh. Exp. Teor. Fiz.* **79** 751 (Engl. transl. 1980 *Sov. Phys.-JETP* **52** 382)
- [10] Manakov N L, Frolov M V, Borca B, and Starace A F 2000 *Pis'ma ZhETF* **72** 294 (Engl. transl. 2000 *JETP Lett.* **72** 426)  
Manakov N L, Frolov M V, Borca B, and Starace A F 2001 *Super-Intense Laser-Atom Physics* (NATO Sci. Ser. II: Math., Phys. and Chemistry, vol 12) ed. P Piraux and K Rzażewski (Dordrecht: Kluwer) p. 295
- [11] Manakov N L, Frolov M V, Borca B, and Starace A F 2001 to be published
- [12] Krstić D S, Milošević D B, and Janev R C 1991 *Phys. Rev. A* **44** 3089
- [13] Langhoff P W, Epstein S T, and Karplus M 1972 *Rev. Mod. Phys.* **44** 602
- [14] Manakov N L, Frolov M V, Starace A F, and Fabrikant I I 2000 *J. Phys. B: At. Mol. Opt. Phys.* **33** R141
- [15] Manakov N L, Preobrazhenskii M A, Rapoport L P, and Fainshtein A G 1978 *Zh. Eksp. Teor. Fiz.* **75** 1243 (Engl. transl. 1978 *Sov. Phys.-JETP* **48** 626)
- [16] Haritos C, Mercouris Th, and Nicolaides C A 2001 *Phys. Rev. A* **63** 013410
- [17] Telnov D A and Chu S I 1994 *Phys. Rev. A* **50** 4099
- [18] Gribakin G F and Kuchiev M Yu 1997 *Phys. Rev. A* **55** 3760
- [19] Dörr M, Potvliege R, Proulx D, and Shakeshaft R 1990 *Phys. Rev. A* **42** 4138