

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Other Publications in Zoonotics and Wildlife
Disease

Wildlife Disease and Zoonotics

2002

Pregnancy status and fetal prion genetics determine PrP^{Sc} accumulation in placentomes of scrapieinfected sheep

Wenbin Tuo

U.S. Department of Agriculture

Katherine I. O'Rourke

U.S. Department of Agriculture, katherine.orourke@ars.usda.gov

Dongyue Zhuang

Washington State University

William P. Cheevers

Washington State University

Terry R. Spraker

Colorado State University - Fort Collins

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/zoonoticspub>



Part of the [Veterinary Infectious Diseases Commons](#)

Tuo, Wenbin; O'Rourke, Katherine I.; Zhuang, Dongyue; Cheevers, William P.; Spraker, Terry R.; and Knowles, Donald P., "Pregnancy status and fetal prion genetics determine PrP^{Sc} accumulation in placentomes of scrapieinfected sheep" (2002). *Other Publications in Zoonotics and Wildlife Disease*. 150.
<https://digitalcommons.unl.edu/zoonoticspub/150>

This Article is brought to you for free and open access by the Wildlife Disease and Zoonotics at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Other Publications in Zoonotics and Wildlife Disease by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Wenbin Tuo, Katherine I. O'Rourke, Dongyue Zhuang, William P. Cheevers, Terry R. Spraker, and Donald P. Knowles

Pregnancy status and fetal prion genetics determine PrP^{Sc} accumulation in placentomes of scrapie-infected sheep

Wenbin Tuo^{*†‡}, Katherine I. O'Rourke^{*}, Dongyue Zhuang[§], William P. Cheevers[§], Terry R. Spraker[¶], and Donald P. Knowles^{*§}

^{*}Animal Disease Research Unit, Agricultural Research Service, U.S. Department of Agriculture, Pullman, WA 99164; [†]Center for Reproductive Biology and [§]Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, WA 99164; and [¶]Colorado State Diagnostic Laboratory, Colorado State University, Fort Collins, CO 80523

Communicated by Janice M. Miller, Agricultural Research Service, U.S. Department of Agriculture, Ames, IA, February 6, 2002 (received for review November 12, 2001)

Ovine scrapie is a fatal neurodegenerative disorder that may be transmitted through exposure to infected uterine and placental tissues. Susceptibility to scrapie is primarily controlled by polymorphisms in the prion protein (PrP) gene. Scrapie in the U.S. Suffolk breed and in many breeds in Europe occurs in sheep homozygous for glutamine (171QQ), but rarely in sheep heterozygous for glutamine and arginine (171QR) or homozygous for arginine (171RR) at codon 171 of the PrP gene. This study demonstrated that accumulation of PrP^{Sc} in uterine-placental epithelial cells in the placentome was determined by fetal PrP genotype and the pregnancy status of scrapie-infected ewes. PrP^{Sc} was detected in 171QQ placentomes of infected ewes, but not in placentomes of infected ewes pregnant with 171QR conceptuses or in the non-pregnant uterus of infected ewes. The distribution of PrP^{Sc} plaques in placentomes was temporally associated with stage of gestation. There was a tendency toward increased size and number of placentomal PrP^{Sc} plaques from the endometrial stalk (maternal side) to chorionic plate (fetal side). These results indicate that accumulation of PrP^{Sc} is eliminated or reduced to undetectable levels in reproductive and placental tissues if infected ewes are not pregnant or conceive conceptuses with a resistant PrP genotype.

Ovine scrapie is a fatal neurodegenerative disorder presumably caused by the aggregated proteinase K (PK) resistant form of the sheep prion (PrP^{Sc}), which is derived from an endogenous, PK-sensitive cellular precursor (PrP^C) through conformational alteration (1–6). This host protein is encoded by a single copy gene (2) and expressed by many cell and tissue types (7–10). In several species, individuals with certain coding mutations in the prion gene are predisposed to prion disease (3, 11, 12). Susceptibility to scrapie infection in Suffolk sheep and Suffolk crosses is determined primarily by codon 171 of the prion gene (11, 13). Sheep homozygous for glutamine at codon 171 (171QQ) are susceptible to scrapie, whereas sheep heterozygous for glutamine and arginine (171QR) or homozygous for arginine (171RR) are usually resistant to scrapie (11, 13).

In sheep, PrP^{Sc} or scrapie infectivity is present primarily in tissues of the central and peripheral nervous system, lymphoreticular system, and placenta (9, 14–21). Recent studies of reproductive and placental tissues from scrapie-infected pregnant sheep demonstrated that PrP^{Sc} accumulates at the fetal–maternal interface (the placentome) and proposed that scrapie is not transmitted *in utero*, but may be transmitted from infected dams to their scrapie susceptible lambs during the perinatal period by exposure to infectious placental tissues (9). The ovine placentome is a natural chimera consisting of interdigitating uterine endometrial septae and placental chorionic villi (Fig. 1), in which the uterine cells carry the maternal genotype and the placental chorionic cells carry the fetal genotype. The unique structures of the placentome provide an applicable model for the study of the effect of prion genotype on scrapie transmission

between tissues with intimate physical contact within a natural host for scrapie. The advantages of this model are that the placentomal PrP^{Sc} in the infected ewes is abundant for detection and PrP genotypes of the placentomal components can be controlled by breeding.

This study determined the effects of pregnancy stage, fetal PrP genotype, and pregnancy status on PrP^{Sc} accumulation at the fetal–maternal interface of scrapie-infected ewes. This study showed that accumulation of PrP^{Sc} in both uterine epithelial and placental trophoblast cells is determined by PrP genotype of the fetus and the pregnancy status of the infected ewe, and that the pattern of PrP^{Sc} plaque distribution in placentomes is associated with gestational stage. These results suggest that scrapie transmission through uterine-placental tissues may be reduced or eliminated if the infected ewes are not pregnant or conceive conceptuses with a resistant genotype.

Materials and Methods

Animals, Tissue Collection, and Processing. All ewes ($n = 61$) in this study were either uninfected or naturally infected 171QQ Suffolk or Suffolk \times Hampshire sheep. The scrapie status of the fetal, young, and adult ewes was determined before euthanasia, at necropsy, or at lambing by immunohistochemistry (22). Of 32 peripheral lymphoid tissue [live animal test of the third-eyelid lymphoid tissue (22)] PrP^{Sc}-positive ewes used in this study, 19 ewes exhibited clinical symptoms including pruritis, wool loss, weight loss, and/or ataxia. Eleven scrapie-infected ewes were mated to a 171QQ ram, 6 scrapie-infected ewes were mated to a 171RR ram, 25 uninfected ewes were mated to a 171QQ ram, and 15 scrapie-infected and 4 uninfected ewes were not mated. Infected ewes pregnant with 171QQ conceptuses were euthanized either at early (days 40–60; $n = 8$), mid- (days 80–100; $n = 1$), or term (Days 140–150; $n = 2$) pregnancy. All infected ewes pregnant with 171QR conceptuses were euthanized at term pregnancy, except for one that was euthanized at mid-pregnancy. All uninfected pregnant ewes were euthanized between days 40 and 140 of pregnancy. The uninfected ewes had no reported exposure to scrapie, and PrP^{Sc} was not detected in brain or tonsil by Western blotting or immunohistochemistry. The genotypes of fetuses/lambs and placentae were determined by sequencing of the prion gene.

Brainstem, tonsil, placentomes, interplacentomal endometri-

Abbreviations: PrP, prion protein; PrP^C, cellular PrP; PrP^{Sc}, scrapie PrP; 171QQ, -RR, or -QR, homozygous for glutamine or arginine, or heterozygous for glutamine and arginine at codon 171 of the PrP gene; PK, proteinase K.

[†]To whom reprint requests should be addressed at: Animal Disease Research Unit, Room 3003 ABDF, Agricultural Research Service, U.S. Department of Agriculture, Washington State University, Pullman, WA 99164-6630. E-mail: wtuo@vetmed.wsu.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.




Fig. 1. A schematic illustration of a medial cross section of an ovine placentome. The placentome is a natural chimera, consisting of interdigitating uterine (placentomal or caruncular endometrium) and placental (placentomal or cotyledonary chorioallantois) tissues. The medial cross section of a placentome is divided into three zones: zones A, B, and C. Zone A contains maternal endometrial crypts and the distal ends of the cotyledonary villous tree and is located closest to the endometrial stalk and myometrium, zone B is the intermediate area, and zone C is the distal (relative to myometrium) portion of the placentome containing the distal ends of endometrial septae and the chorionic plate. Zone C is typically characterized by structures known as arcade system or hematomata. The placentomal endometrium consists of the uterine superficial epithelium and associated stroma, but no glands. Interplacentomal endometrium consists of loosely attached intercotyledonary chorioallantois and intercaruncular endometrium, which is rich in glands. Pla. villus, placental villus; m-septum, maternal endometrial septum; m-crypt, maternal crypt; end. stalk, endometrial stalk.

al-chorioallantois, ovary, oviduct, mammary gland, and bladder collected at euthanasia were fixed in 10% buffered formalin and processed for immunohistochemistry as described previously (22). Maternal brainstem, tonsil, dissected uterine caruncular (caruncles) and interplacentomal (intercaruncular) endometria, placental cotyledonary (cotyledons) and interplacentomal (intercotyledonary) chorioallantois, ovary, oviduct, mammary gland, and bladder of pregnant ewes, and fetal brain, kidney and bladder at different stages of pregnancy were collected and snap-frozen in liquid nitrogen and stored at -80°C until use. For each fetal unit, 2–3 placentomes and dissected caruncles and cotyledons from 3–6 placentomes were collected from each of the three areas occupied by the head, mid-body and tail of a fetus. A schematic illustration of the ovine placentome is shown in Fig. 1. Animal Care and Use Protocols were approved by the Institutional Animal Care and Use Committee of Washington State University.

Tissue Homogenate Preparation and Western Blot Analysis. Tissue homogenates were prepared with (for PrP^{Sc} detection) or without (for PrP^C and/or PrP^{Sc} detection) the use of 10% sarkosyl and ultracentrifugation as described (9). Briefly, tissue was homogenized and incubated for 30 min at 37°C in 10 mM Tris-HCl (pH 7.5), 5 mM MgCl₂, and DNase (200 $\mu\text{g}/\text{g}$ wet tissue), followed by 30 min incubation at room temperature in sarkosyl solution (10 mM Tris-HCl, pH 7.5/20% sarkosyl). The homogenate was centrifuged at $6,000 \times g$ for 10 min, and the resultant supernatant was centrifuged at $348,000 \times g$ for 50 min at room temperature. The pellet was dissolved in 10 mM Tris-HCl (pH 7.5) and treated with or without PK (20 $\mu\text{g}/\text{ml}$) at 37°C for 30 min. The PK-treated mixture was centrifuged at $279,000 \times g$ for 30 min. The pellet was resuspended in 10 mM Tris-HCl and boiled in sample buffer before Western blotting.

Western blot procedure was performed (9, 23). Briefly, tissue homogenates were analyzed by 12% SDS/PAGE (Invitrogen), followed by transfer to poly(vinylidene difluoride) (PVDF) membranes (Amersham Pharmacia) before detection by anti-prion mAb F99/97.6.1 (22) and goat anti-mouse IgG-horseradish peroxidase (HRPO) (Southern Biotechnology Associates), and developed with a chemiluminescence substrate (Amersham Pharmacia).

Immunohistochemistry. Immunohistochemistry was performed as previously described (22). Briefly, tissues were fixed in 10% neutral buffered formalin for 2–7 days, treated with 95% formic acid for 1 h, rinsed with water, and fixed for an additional 24 h before embedding in paraffin. Tissue sections placed on glass slides were deparaffinized, hydrated, and autoclaved in Target Retrieval Solution (pH 6; DAKO) at 121°C for 30 min. Slides were stained by using an automated stainer (Ventana Medical Systems, Tucson, AZ). Anti-prion mAbs F99/97.6.1 and F89/160.1.5 (22, 23) were used as primary reagents, followed by the application of a biotinylated secondary antibody, streptavidin-alkaline phosphatase complex and alkaline phosphatase substrate (Ventana Medical Systems). Controls included isotype-matched irrelevant IgG and blocking of the mAbs with synthetic peptides to which the mAbs were raised. All slides were counterstained with hematoxylin. Photomicrographs were taken with a microscope equipped with a digital camera.

The medial cross section of a placentome was divided into three zones in parallel to the myometrium (Fig. 1). Zone A is proximal to the myometrium and composed of endometrial-chorioallantoic interdigitation at the base of the placentome; zone C is composed of the distal portion of the interdigitating tissue, which contains the arcade systems or hematomata; and zone B is the intermediate area. All stained PrP^{Sc} plaques within each zone of the entire medial sections of placentomes of early, mid-, and term pregnant scrapie-infected ewes were enumerated by using a bright field microscope.

Statistical Analysis. Percent total PrP^{Sc} plaques from different zones of placentomes of infected ewes with 171QQ conceptuses during early and term pregnancies were analyzed by one-way ANOVA with a Student-Newman Keuls multiple comparisons test. Probability values of $P < 0.05$ were used.

Results

Detection Of PrP^{Sc} in Placentomes Consisting of 171QQ Uterine Caruncular Endometrium and 171QQ Placental Cotyledonary Chorioallantois from Scrapie-Infected Ewes. Scrapie-infected sheep were defined by PrP^{Sc} accumulation in brain and/or lymphoid tissue by using an immunohistochemistry assay (22). Uterine and placental tissues collected during various stages of pregnancy were assayed for PrP^{Sc} by immunohistochemistry. PrP^{Sc} was detected in placentomes consisting of 171QQ uterine (caruncular endometrium) and 171QQ placental (cotyledonary chorioallantois) tissues collected from 10 of 11 of the pregnant (day 40–term post mating) scrapie-infected ewes (Fig. 2 A and D). PrP^{Sc} was not detected in 171QQ placentomes of 1 of 11 pregnant scrapie infected ewes (Table 1). This placentomal PrP^{Sc}-negative sheep was bred before arriving at this U.S. Department of Agriculture/Agricultural Research Service station; therefore, the gestational stage at euthanasia was unknown. However, we determined that this sheep was at an early stage of pregnancy when euthanized (data not shown). PrP^{Sc} was not detected in maternal intercaruncular endometrium, intercotyledonary chorioallantois, amniotic membrane, myometrium, ovary, oviduct, bladder, or mammary gland, fetal bladder, kidney, and brain of these pregnant scrapie-infected ewes (data not shown), or 171QQ placentomes of uninfected ewes ($n = 25$; Table 1). Interestingly, no PrP^{Sc} was detected in placentomes of




Fig. 2. Immunohistochemical analysis of PrP^{Sc} in placentomes of scrapie-infected ewes. The presence of pink staining is indicative of the presence of PrP^{Sc}. TRO/END, trophoblast-endometrium interdigitation; TRO, trophoblast; END, endometrium; m-crypt, maternal crypt; MNC, multinucleated cells; RBC, maternal erythrocytes; hemat, hematoma; D-END, degenerating endometrium; ALL, allantoic cavity; BNC, binucleate cells; SYN, endometrial syncytium. (A) PrP^{Sc} plaques in the 171QQ placentome of a scrapie-infected ewe at day 40 of pregnancy ($\times 100$). PrP^{Sc} was seen only in endometrial/placental interdigitation, not in endometrium alone. The yellow line represents the border between maternal endometrium and trophoblast/endometrium interdigitation at the fetal-maternal interface at the bottom of the maternal crypt. (B) PrP^{Sc} in cells in the maternal crypt (m-crypt) in the 171QQ placentome of an early pregnant scrapie-infected ewe ($\times 200$). (C) PrP^{Sc} in a multinucleated cell (MNC) located in the maternal crypt in the 171QQ placentome of an early pregnant scrapie-infected ewe ($\times 400$). (D) A PrP^{Sc} plaque in zone C of a 171QQ placentome of a term pregnant scrapie-infected ewe ($\times 100$). The yellow circle defines the PrP^{Sc} plaque. (E) PrP^{Sc} in maternal endometrial epithelial cells in the 171QQ placentome of a term pregnant scrapie-infected ewe ($\times 400$). (F) PrP^{Sc} in both endometrial epithelial cells (END) and trophoblast cells (TRO) in the 171QQ placentome of a term pregnant scrapie-infected ewe ($\times 200$). (G) PrP^{Sc} in placental trophoblast cells (TRO) located along the hematoma (hemat) ($\times 400$). Phagocytosed RBCs (RBC and arrow) were apparent in the placental trophoblast cells. The yellow line defines part of the hematoma containing maternal blood (RBC) and degenerating endometrium (D-END). (H) A placental cross section near the endometrial stalk stained with an isotype-matched mAb ($\times 400$).

171QQ uterine and 171QR placental tissues from scrapie-infected ewes ($n = 6$; Table 1).

All ewes studied were between 2 and 5 years old and had PrP^{Sc} in the brain and lymphoid tissue. Eight of these 18 ewes exhibited clinical signs of scrapie infection at euthanasia. However, no correlation was found between the presence of PrP^{Sc} in placentomes and age, stage of pregnancy, or clinical status of the dams (Table 1).

We confirmed the presence of PrP^{Sc} by Western blotting in 171QQ placental endometrium and chorioallantois collected at term pregnancy (Fig. 3C, lanes 4 and 6). PrP^{Sc} was not detected in uterine or placental tissues of uninfected ewes (Fig. 3A, lanes 5 and 8) or in scrapie-infected ewes pregnant with

171QR conceptuses (Fig. 3D, lanes 4 and 6). PrP^C in brain, uterine, and placental tissues of uninfected sheep (Fig. 3A, lanes 1, 3, and 6) and PrP^{Sc} in brain of scrapie-infected sheep (Fig. 3B–D, lane 2) were readily detectable by Western blotting using direct tissue homogenates without further enrichment (9). However, PrP^{Sc} was not consistently detectable in the uterine and placental tissues unless prepared by ultracentrifugation in the presence of sarkosyl (9, 17). Furthermore, PrP^{Sc} was not detectable by Western blotting in uterine and placental tissues of early pregnant (day 40–60 post mating) scrapie-infected ewes (data not shown). Therefore, uterine and placental tissues collected from term pregnant ewes were prepared for Western blotting by ultracentrifugation in the presence of sarkosyl.

PrP^{Sc} Was Not Detected in Reproductive Tissues of Non-Pregnant, 171QQ, Scrapie-Infected Ewes. PrP^{Sc} accumulation in the reproductive tissues in 15 scrapie-infected and 4 uninfected non-pregnant 171QQ ewes was determined (Table 1). All scrapie-infected ewes had PrP^{Sc} accumulation in brain and lymphoid tissue and some (11 of 15) exhibited clinical signs of scrapie (Table 1). No PrP^{Sc} was detected in brain or lymphoid tissues of uninfected non-pregnant ewes (data not shown). Homogenates of tissues from infected ewes were prepared with or without ultracentrifugation in the presence of sarkosyl. PrP^{Sc} was not detected by either Western blotting (Fig. 3B, lanes 5 and 8) or immunohistochemistry (data not shown) in brain, maternal caruncular endometrium and myometrium, intercaruncular endometrium, ovary, oviduct, bladder, or mammary gland of non-pregnant scrapie-infected ewes, nor in any of the reproductive tissues of non-pregnant uninfected ewes (data not shown).

Distribution of PrP^{Sc} Plaques in 171QQ Placentomes of Scrapie-Infected Ewes Is Associated with Stage of Pregnancy. Ten scrapie-infected ewes tested positive for PrP^{Sc} in placentomes collected between day 40 post mating and term (Table 1). PrP^{Sc} was localized as plaques in placentomes of these ewes by immunohistochemistry (Fig. 2A and D), and the size and location of PrP^{Sc} plaques in these placentomes varied depending on gestational stage (Table 2). PrP^{Sc} plaques were seen in placental areas where both placental villi and endometrial septae were present, but not in placental regions with endometrial tissue alone (Fig. 2A). PrP^{Sc} plaques in early pregnant placentomes (days 40–60 post mating; Fig. 2A) were small relative to the plaques in term placentomes (Fig. 2D) and were located in the area of maternal endometrial crypts of zone A (Fig. 4; Table 2). PrP^{Sc} staining in term pregnant placentomes was a mixture of small and large plaques located primarily in zone C near the chorionic plate of the placentome (Table 2; Fig. 4). The PrP^{Sc} plaques in zone C (75% of total plaques) were more abundant ($P < 0.001$) than those in zone B (16% of total plaques) and/or zone A (7% of total plaques) of term pregnant placentomes of scrapie-infected ewes (Fig. 4). Zone B of the placentome had primarily small PrP^{Sc} plaques (data not shown).

Cellular Localization of PrP^{Sc} in Uterine Caruncular Endometrial Epithelium and Placental Cotyledonary Chorioallantoic Trophoblast of 171QQ Placentomes. Cellular localization of PrP^{Sc} in placentomes of scrapie-infected ewes is shown in Fig. 2. Both mAbs used in the present study bound with identical patterns; the results obtained by using mAb F99/97.6.1 are shown. As shown in Fig. 2B and C for placentomes collected from early pregnant scrapie-infected ewes, PrP^{Sc} staining was primarily associated with uterine endometrial epithelium, trophoblast cells, and multinucleate cells in zone A of the maternal endometrial crypts. Very little mAb binding was found in zones B and C of placentomes from early pregnant scrapie-infected ewes. PrP^{Sc} was not detected in the stroma or blood vessels (data not shown).

Table 1. Correlation between the presence of PrP^{Sc} in placentomes and fetal prion genotypes or pregnancy status of the scrapie infected ewes

Scrapie status	Genotype		No. of dams	Clinical disease	Pregnancy status	PrP ^{Sc} accumulation in tissues		
	Dam	Fetus				Bra + lym	Plact	Np-ute
Uninfected	QQ	QQ	25	0/25	Px	0/25	0/25	—
	QQ	—	4	0/4	NP	0/4	—	0/4
Infected	QQ	QQ	11	6/11	Px	11/11	10/11	—
	QQ	QR	6	2/6	Px	6/6	0/6	—
	QQ	—	15	11/15	NP	15/15	—	0/15

Uninfected, Animals that had no reported exposure to scrapie and disease status confirmed by Western blotting and immunohistochemistry of postmortem brain and lymphoid tissues; Infected, animals that either exhibited clinical signs of scrapie or had detectable PrP^{Sc} in the lymphoid tissue (based on a live animal third eyelid test) prior to experiment. Genotype, Allelic coding amino acid variation at codon 171 of the prion gene; QQ, homozygous for glutamine; QR, heterozygous for glutamine and arginine; RR, homozygous for arginine. Clinical disease, The number of sheep with clinical scrapie vs. total number of infected sheep used (preclinical sheep had PrP^{Sc} in brain and/or lymphoid tissues but no clinical signs, which include rubbing and wool loss, progressive weight loss, and ataxia). Pregnancy status, Px, pregnant; NP, non-pregnant. Bra + lym, PrP^{Sc} detected in postmortem brain and lymphoid tissue; plact, placentome; np-ute, non-pregnant uterus.

No particular uterine or placental structures were associated with this distribution pattern.

In term pregnant scrapie-infected ewes, PrP^{Sc} was clearly




Fig. 3. Western blot of PrP^{Sc} in tissue homogenates of brainstem, uterine caruncular endometrium, and placental cotyledonary chorioallantois of pregnant and non-pregnant ewes with or without scrapie. Protein from 0.5 mg brainstem or 30 mg endometrium, chorioallantois, and other tissues was loaded in each lane. PK (20 μg/ml) treatment at 37°C for 30 min of each tissue homogenate is indicated at the bottom of each panel (+). (A) Western blot of tissue homogenates prepared from brainstem (lanes 1 and 2), uterine caruncular endometrium (lanes 3–5), and cotyledonary chorioallantois (lanes 6–8) of an uninfected ewe (171QQ) pregnant (day 145 post mating) with a 171QQ conceptus. Samples in lanes 4, 5, 7, and 8 were prepared with sarkosyl and ultracentrifugation. Tissue homogenates were with (lanes 2, 5, and 8) or without (lanes 1, 3, 4, 6, and 7) PK treatment. (B) Western blot of tissue homogenates of brainstem (lanes 1 and 2), endometrium (lanes 3–5), and myometrium (lanes 6–8) of a non-pregnant scrapie-infected ewe (171QQ). Samples in lanes 4 and 5 and in lanes 7 and 8 were prepared with sarkosyl and ultracentrifugation. Tissue homogenates were with (lanes 2, 5, and 8) or without (lanes 1, 3, 4, 6, and 7) PK treatment. (C) Western blot of tissue homogenates prepared from brainstem, caruncular endometrium, and cotyledonary chorioallantois of a scrapie-infected ewe (177QQ) pregnant with a 171QQ conceptus. Lanes 1 and 2, brainstem prepared without sarkosyl and ultracentrifugation; lanes 3 and 4, uterine caruncular endometrium prepared with sarkosyl and ultracentrifugation; and lanes 5 and 6, placental cotyledonary chorioallantois prepared with sarkosyl and ultracentrifugation. Tissue homogenates were with (lanes 2, 4, and 6) or without (lanes 1, 3, and 5) PK treatment. (D) Western blot of tissue homogenates prepared from brainstem, caruncular endometrium, and cotyledonary chorioallantois of a scrapie-infected ewe (177QQ) pregnant with a 171QR conceptus. Lanes 1 and 2 (brainstem), prepared without sarkosyl and ultracentrifugation; lanes 3 and 4 (uterine caruncular endometrium), prepared with sarkosyl and ultracentrifugation; and lanes 5 and 6, placental cotyledonary chorioallantois prepared with sarkosyl and ultracentrifugation. Tissue homogenates were with (lanes 2, 4, and 6) or without (lanes 1, 3, and 5) PK treatment.

detected in single cell layers of the endometrial epithelium and placental trophoblast cells (Fig. 2 E, F, and G), as well as in the uterine syncytium (data not shown), a structure formed by fusion of endometrial epithelial cells during normal pregnancy (24). Although staining for PrP^{Sc} was present in both uterine epithelium and trophoblast cells throughout the placentome, there was a tendency toward increased accumulation of PrP^{Sc} in trophoblast cells in zone C rather than zones A and B in scrapie-infected ewes. In addition, abundant PrP^{Sc} was detected in chorionic trophoblast cells around the hematoma, which contains maternal blood as a result of degeneration of the distal ends of the maternal endometrial septae in this region (refs. 24 and 25; Fig. 2G).

PrP^{Sc} was not detected by immunohistochemistry in endometrial glands, which were abundant in the interplacentomal region but absent in the placentomal region, nor in intercaruncular endometrium, myometrium, or the interplacentomal endometrial-chorioallantois of scrapie-infected ewes (data not shown). PrP^{Sc} was not found in replicate tissue sections when isotype-matched irrelevant IgG mAbs or mAbs absorbed with peptide-antigen were used as the primary antibody (Fig. 2H).

Discussion

Susceptibility to scrapie in Suffolk and Suffolk cross-bred sheep is determined primarily by codon 171 of the PrP gene. Sheep with a 171QQ genotype are susceptible to scrapie, and those with a 171QR or 171RR genotype are resistant to scrapie (11, 13), with a few exceptions (26, 27). The present study investigated the effects of offspring genotype and pregnancy on PrP^{Sc} accumulation at the fetal–maternal interface of scrapie-infected sheep and possible transmission between tissues of different genotypes by using the ovine placentome as a natural chimeric model.

PrP^{Sc} was present at the fetal–maternal interface, endometrium, and the apposing placental chorioallantois in 171QQ placentomes of most (10/11) infected ewes with PrP^{Sc} in the brain and lymphoid tissue. PrP^{Sc} was not detected in the uterine tissues of non-pregnant infected ewes, nor in the uterine or placental tissues of infected ewes carrying 171QR fetuses. PrP^{Sc} was not detected in 171QQ placentomes from one scrapie-infected ewe. Based on placental morphology, this sheep was considered at an early stage of pregnancy when euthanized, suggesting that detectable levels of PrP^{Sc} had not yet accumulated in placentomes. The results of this study suggest that the accumulation of PrP^{Sc} in the uterus is pregnancy dependent and that PrP^{Sc} accumulation in the placentome requires the presence of a 171QQ conceptus. These observations support earlier reports that transmission is related to lambing (9, 14, 17) and suggest that transmission may be reduced or eliminated if

Table 2. Correlation between stage of pregnancy and spatial distribution of PrP^{Sc} in placentomes of pregnant scrapie-infected ewes

PrP ^{Sc}	Stage of pregnancy		
	Early (n = 7)	Mid (n = 1)	Term (n = 2)
Pattern of PrP ^{Sc} distribution	Numerous small plaques in multinucleate cells	Mostly small plaques and few large plaques	Primarily large plaques spanning several villi
Location of PrP ^{Sc} plaque	Maternal crypts of zone A	Primarily maternal crypt of zone A and some in zone B	Primarily zone C

PrP^{Sc} was defined by immunohistochemistry. Early pregnancy, days 40–60 post mating; mid-pregnancy, days 80–100 post mating; term pregnancy, days 140–150 post mating.

infected ewes are not pregnant or conceive offspring with a scrapie-resistant genotype. The mechanisms by which pregnancy affects PrP^{Sc} deposition in the uterus of genetically susceptible sheep are not known. We showed previously that PrP^C expression is up-regulated in pregnant caruncular endometrium where intimate contact of placental and uterine tissues occur, suggesting that molecules expressed by the cotyledonary trophoblast cells, uterine-placental cell-cell contact in the placentome, and/or the presence of pregnancy-associated hormones are important for PrP^C expression and PrP^{Sc} accumulation in the caruncular endometrium during pregnancy (9). It is intriguing that PrP^{Sc} was absent from the uterine tissue of infected ewes pregnant with 171QR conceptuses. This “blockage” of *in vivo* PrP^{Sc} formation in the presence of placental 171QR and placental/maternal 171QQ PrP may be similar to the interference of PrP^{Sc} formation by heterologous PrP molecules shown *in vitro* (28, 29). One may speculate that placental 171QR PrP expressed by trophoblast cells may interact with uterine PrP^{Sc} through cell-cell contact and inhibit the conversion process. *In vitro* and *in vivo* studies to examine the role of cell-free and cell-associated trophoblast-derived factors in PrP^{Sc} propagation in uterine cells and the systemic effects of pregnancy in the absence of cell-to-cell contact with trophoblast will provide additional information on this critical transmission step. The efficient conversion of PrP in the placentome (detectable during 40–145 days post mating) compared with the lymphoid tissues (10–14 months after exposure during the perinatal period) or the brain (2 to 3 years after exposure) (15) provides a useful model system for examining PrP conversion under natural conditions.

This study demonstrated that PrP^{Sc} was localized to the ovine uterine and placental epithelial cells. Because no antibodies to specific markers of these epithelial cells were available, the maternal or fetal/placental origin of these cells at the interface during early pregnancy was estimated by location and morphol-

ogy under light microscopy. PrP^{Sc} was detected in the uterine cells, mononuclear trophoblast cells, and the multinucleated cells formed by fusion between uterine epithelial and placental trophoblast cells. However, PrP^{Sc} appeared more abundant in the endometrial epithelial cells than in trophoblast cells in zones A and B of the placentome during early and mid pregnancy. The binucleate cells that originate from the trophoblast cells and invade massively into the endometrium were generally not associated with PrP^{Sc} staining (30). PrP^{Sc} was clearly present in both uterine and trophoblast cells in zone C of the prepartum placentome particularly. Abundant PrP^{Sc} was present in trophoblast cells in zone C near the allantoic cavity. The translocation of the PrP^{Sc} from primarily uterine epithelial cells in zone A during early pregnancy to the placental trophoblast cells in zone C at term pregnancy may represent a critical step in the transmission of the scrapie agent from the maternal tissue to placental tissue.

Interestingly, placental PrP^{Sc} was distributed in discrete plaques. The ovine placentome appears to be a homogenous interdigitation between the endometrial septae and chorioallantoic villi, and the PrP^{Sc} plaques in placentomes were not associated with any identifiable landmark structures. However, the spatial distribution pattern of the PrP^{Sc} plaques was highly associated with stage of pregnancy. The transition from smaller PrP^{Sc} plaques in zone A near the endometrial stalk during early pregnancy to relatively larger PrP^{Sc} plaques in zone C near the chorionic plate at term pregnancy may represent a temporal process of PrP^{Sc} accumulation and maternal-placental PrP^{Sc} transmission. The mechanism of such a transient change in plaque size and location are currently unknown. The fact that placental PrP^{Sc} plaques of early pregnancy were formed primarily in the maternal endometrial crypts containing the top of the chorionic villi, but not in the endometrium without the chorionic villous tree, supports the concept that the presence of placental tissue is critical for PrP^{Sc} accumulation. At term pregnancy, PrP^{Sc} plaques were located in the vicinity of the structure known as hematoma in zone C of the placentome. The formation of ovine placentomes is initiated at the time of attachment of uterine caruncles to placental cotyledons and is gradually completed by mid-pregnancy through uterine-placental interdigitation when the distal ends of the endometrial septae and chorionic villi reach the chorionic plate and endometrial stalk, respectively (Fig. 1). During the second half of pregnancy in sheep, the distal ends of the endometrial septae degenerate through an unknown mechanism, and maternal blood leaks through the broken blood vessels into the space surrounded by the intact trophoblast cells of the chorionic villi, leading to the formation of hematoma or the arcade systems in the placentome (24). By term pregnancy, particularly immediately before lambing, massive bleeding is seen in all placentomes, appearing as a dark brown color in the cotyledonary chorioallantois. Thus, hematoma contains maternal blood and degenerated and degenerating endometrial tissue. The trophoblast cells of the hematoma are highly phagocytic, and are thought to play an important role in nonspecific uptake of




Fig. 4. Distribution of PrP^{Sc} plaques in zones A, B, and C of the placentome of ewes infected with scrapie during early (n = 7) and term (n = 2) stages of pregnancy. The total number of PrP^{Sc} plaques in medial sections of two separate placentomes of each scrapie-infected ewe was counted microscopically. The results are expressed as the percentage of total PrP^{Sc} plaques.

maternal endometrial secretions, cellular debris, and maternal blood, especially red blood cells, as a source for nutrients of the fetus (24, 25, 31). Indeed, the phagocytosis of maternal red blood cells by placental trophoblast cells was seen in both normal and scrapie-infected animals. Because phagocytosis at the fetal–maternal interface is a nonspecific process, this may be one way for the placental trophoblast cells to acquire PrP^{Sc} from the degenerating endometrial cells in the hematoma.

This study demonstrated that offspring genotype and pregnancy are determining factors for PrP^{Sc} accumulation at the fetal–maternal interface of scrapie-infected ewes. The characteristic spatial distribution of PrP^{Sc} at the interface during different stages of pregnancy may indicate a temporal course of maternal–placental transmission of scrapie. This study presented evidence that scrapie may not be transmissible between infected genetically susceptible uterine tissue and placental tissue with a resistant genotype using the natural chimeric model, the ovine placentome. The results of this study provide useful information for further understanding of scrapie transmission in species with

extraneural accumulation of PrP^{Sc}. The nature of the PrP^{Sc} converting process in the pregnant uterus, the possible inhibitory effect of placental 171QR PrP molecules on PrP^{Sc} formation in the uterus, the identity of the pregnancy-related cells or molecules that allow the rapid accumulation of PrP^{Sc} in the placentome, and the source of PrP^{Sc} in phagocytic trophoblast cells remain to be determined. Additional *in vivo* studies with sheep at different stages of gestation resulting from genetically controlled mating as well as *in vitro* studies with uterine epithelial and trophoblast cell lines will provide a better understanding of cellular and molecular basis for transmission of scrapie, a prototype transmissible spongiform encephalopathy.

We thank Dr. Charles Leathers for critical review of the manuscript; Anne Anderson, Will Harwood, and Amy Lyda for excellent technical assistance; and Dwayne Chandler, Emma Karel, and Peter Steiner for assistance with animal care and necropsy. This research was supported by grants from the U.S. Department of Agriculture/Agricultural Research Service (CWU 5348-32000-015-00D).

- Prusiner, S. B. (1982) *Science* **216**, 136–144.
- Oesch, B., Westaway, D., Walchli, M., McKinley, M. P., Kent, S. B., Aebersold, R., Barry, R. A., Tempst, P., Teplow, D. B. & Hood, L. E. (1985) *Cell* **40**, 735–746.
- Basler, K., Oesch, B., Scott, M., Westaway, D., Walchli, M., Groth, D. F., McKinley, M. P., Prusiner, S. B. & Weissmann, C. (1986) *Cell* **46**, 417–428.
- Aguzzi, A. & Weissmann, C. (1998) *Haemophilia* **4**, 619–627.
- Prusiner, S. B. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 13363–13383.
- Caughey, B. (2001) *Trends Biochem. Sci.* **26**, 235–242.
- Bendheim, P. E., Brown, H. R., Rudelli, R. D., Scala, L. J., Goller, N. L., Wen, G. Y., Kasczak, R. J., Cashman, N. R. & Bolton, D. C. (1992) *Neurology* **42**, 149–156.
- Horiuchi, M., Yamazaki, N., Ikeda, T., Ishiguro, N. & Shinagawa, M. (1995) *J. Gen. Virol.* **76**, 2583–2587.
- Tuo, W., Zhuang, D., Knowles, D. P., Cheevers, W. P., Sy, M. S. & O'Rourke, K. I. (2001) *J. Biol. Chem.* **276**, 18229–18234.
- Liu, T., Li, R., Wong, B. S., Liu, D., Pan, T., Petersen, R. B., Gambetti, P. & Sy, M. S. (2001) *J. Immunol.* **166**, 3733–3742.
- Westaway, D., Zuliani, V., Cooper, C. M., Da Costa, M., Neuman, S., Jenny, A. L., Detwiler, L. & Prusiner, S. B. (1994) *Genes Dev.* **8**, 959–969.
- Prusiner, S. B. & Scott, M. R. (1997) *Annu. Rev. Genet.* **31**, 139–175.
- O'Rourke, K. I., Holyoak, G. R., Clark, W. W., Mickelson, J. R., Wang, S., Melco, R. P., Besser, T. E. & Foote, W. C. (1997) *J. Gen. Virol.* **78**, 975–978.
- Pattison, I. H., Hoare, M. N., Jebbett, J. N. & Watson, W. A. (1972) *Vet. Rec.* **90**, 465–468.
- Hadlow, W. J., Kennedy, R. C. & Race, R. E. (1982) *J. Infect. Dis.* **146**, 657–664.
- Van Keulen, L. J., Schreuder, B. E., Meloen, R. H., Poelen-van den Berg, M., Mooij-Harkes, G., Vromans, M. E. & Langeveld, J. P. (1995) *Vet. Pathol.* **32**, 299–308.
- Race, R., Jenny, A. & Sutton, D. (1998) *J. Infect. Dis.* **178**, 949–953.
- Van Keulen, L. J., Schreuder, B. E., Vromans, M. E., Langeveld, J. P. & Smits, M. A. (1999) *J. Comp. Pathol.* **121**, 55–63.
- Beekes, M. & McBride, P. A. (2000) *Neurosci. Lett.* **278**, 181–184.
- Heggebo, R., Press, C. M., Gunnes, G., Inge, L. K., Tranulis, M. A., Ulvund, M., Groschup, M. H. & Landsverk, T. (2000) *J. Gen. Virol.* **81**, 2327–2337.
- Andreoletti, O., Berthon, P., Marc, D., Sarradin, P., Grosclaude, J., van Keulen, L., Schelcher, F., Elsen, J. M. & Lantier, F. (2000) *J. Gen. Virol.* **81**, 3115–3126.
- O'Rourke, K. I., Baszler, T. V., Besser, T. E., Miller, J. M., Cutlip, R. C., Wells, G. A., Ryder, S. J., Parish, S. M., Hamir, A. N., Cockett, N. E., *et al.* (2000) *J. Clin. Microbiol.* **38**, 3254–3259.
- O'Rourke, K. I., Baszler, T. V., Miller, J. M., Spraker, T. R., Sadler-Riggleman, I. & Knowles, D. P. (1998) *J. Clin. Microbiol.* **36**, 1750–1755.
- Wimsatt, W. A. (1950) *Am. J. Anat.* **87**, 391–457.
- Mossman, H. W. (1987) *Vertebrate Fetal Membranes* (Rutgers Univ. Press, New Brunswick, NJ).
- Ikeda, T., Horiuchi, M., Ishiguro, N., Muramatsu, Y., Kai-Uwe, G. D. & Shinagawa, M. (1995) *J. Gen. Virol.* **76**, 2577–2581.
- Tranulis, M. A., Osland, A., Bratberg, B. & Ulvund, M. J. (1999) *J. Gen. Virol.* **80**, 1073–1077.
- Priola, S. A., Caughey, B., Race, R. E. & Chesebro, B. (1994) *J. Virol.* **68**, 4873–4878.
- Horiuchi, M., Priola, S. A., Chabry, J. & Caughey, B. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 5836–5841.
- Wooding, F. B., Chambers, S. G., Perry, J. S., George, M. & Heap, R. B. (1980) *Anat. Embryol.* **158**, 361–370.
- Dantzer, V. (1998) in *Encyclopedia of Reproduction*, eds Knobil, E. & Neil, J. (Academic, San Diego), pp. 18–28.