

2009

Nebraska Statewide Groundwater-Level Monitoring Report 2009

Jesse T. Korus Dr.

University of Nebraska-Lincoln, jkorus3@unl.edu

Mark E. Burbach

University of Nebraska-Lincoln, mburbach1@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/conservationsurvey>



Part of the [Geology Commons](#), [Geomorphology Commons](#), [Hydrology Commons](#), [Paleontology Commons](#), [Sedimentology Commons](#), [Soil Science Commons](#), and the [Stratigraphy Commons](#)

Korus, Jesse T. Dr. and Burbach, Mark E., "Nebraska Statewide Groundwater-Level Monitoring Report 2009" (2009). *Conservation and Survey Division*. 161.

<http://digitalcommons.unl.edu/conservationsurvey/161>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Conservation and Survey Division by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Nebraska Statewide Groundwater-Level Monitoring Report

2009

Jesse T. Korus and Mark E. Burbach
Conservation and Survey Division
School of Natural Resources

Nebraska Water Survey Paper Number 76

Institute of Agriculture and Natural Resources
University of Nebraska–Lincoln



Nebraska Statewide Groundwater-Level Monitoring Report 2009

Jesse T. Korus and Mark E. Burbach

*Conservation and Survey Division
School of Natural Resources*

Nebraska Water Survey Paper Number 76

Conservation and Survey Division
School of Natural Resources
Institute of Agriculture and Natural Resources
University of Nebraska—Lincoln

University of Nebraska-Lincoln

Harvey S. Perlman, J.D., Chancellor, University of Nebraska-Lincoln

John C. Owens, Ph.D., Vice Chancellor for Agriculture and Natural Resources

Donald A. Wilhite, Ph.D., Director, School of Natural Resources

Mark S. Kuzila, Ph.D., Director, Conservation and Survey Division

The Conservation and Survey Division of the University of Nebraska-Lincoln is the agency designated by statute to investigate and interpret the geologically related natural resources of the state, to make available to the public the results of these investigations, and to assist in the development and conservation of these resources. It consists of program areas in geology, water, soils, and remote sensing-geographic information systems.

The division is authorized to enter into agreements with federal and state agencies to engage in cooperative surveys and investigations of the state. Publications of the division and the cooperating agencies are available through the Conservation and Survey Division, 101 Hardin Hall, University of Nebraska-Lincoln, Lincoln, NE 68583-0961. Contact the address above, phone : (402) 472-3471, or e-mail snrsales@unl.edu. The Conservation and Survey Division web site is: <http://snr.unl.edu/csd/>.

The University of Nebraska-Lincoln does not discriminate based on gender, age, disability, race, color, religion, marital status, national or ethnic origin or sexual orientation. The University of Nebraska-Lincoln is an equal opportunity educator and employer with a comprehensive plan for diversity.

December 2009

ISBN-13 978-1-56161-012-9

ISBN-10 56161-012-7

ACKNOWLEDGMENTS

The cooperation and assistance of the following agencies and associations in collecting and providing water-level data during 2009 are gratefully acknowledged: U.S. Bureau of Reclamation; Central Nebraska Public Power and Irrigation District; U.S. Geological Survey; and the following Natural Resources Districts; Lower Republican, Middle Republican, Upper Republican, Upper Big Blue, Little Blue, Lower Big Blue, Lower Platte South, Lower Platte North, Central Platte, Twin Platte, North Platte, South Platte, Lower Niobr-

ara, Middle Niobrara, Upper Niobrara-White, Lower Loup, Upper Loup, Lower Elkhorn, Upper Elkhorn, Papio-Missouri River, Lewis and Clark, Nemaha, and Tri-Basin. We also thank the many hundreds of different land-owners who graciously allowed these agencies to collect water-level information from their wells and install observation wells on their land.

Thanks to Dee Ebbeka and Les Howard for their assistance with graphics and maps.

CONTENTS

Introduction	2
Purpose and Methods	4
Factors Causing Groundwater-Level Changes	4
Status of Groundwater-Level Monitoring Program, 2009	6
Changes in Groundwater Levels, Spring 2008 to Spring 2009	8
Changes in Groundwater Levels, Spring 2000 to Spring 2009	10
Changes in Groundwater Levels, Predevelopment to Spring 2009	12
Average Daily Streamflows, 2008	14
Groundwater-Level Change Statistics	16
Groundwater-Level Hydrographs	18

FIGURES

Statewide Subdivisions used to Describe Regional Trends	3
Important Aquifers and Topographic Regions of Nebraska	3
Example of Groundwater-Level Changes at Different Temporal Scales	5
Locations of Spring/Fall and Monthly Observation Wells	7
Graphs showing Numbers of Groundwater-Level Monitoring Wells Statewide since 1930	7
Percent of Normal Precipitation - January 2008 to January 2009	9
Groundwater-Level Changes in Nebraska - Spring 2008 to Spring 2009	9
Percent of Normal Precipitation - January 2000 to January 2009	11
Groundwater-Level Changes in Nebraska - Spring 2000 to Spring 2009	11
Density of Active Registered Irrigation Wells - January 2009	13
Groundwater-Level Changes in Nebraska - Predevelopment to Spring 2009	13
Average Streamflow in 2008, as a Percentage of the Long-term Average	14

TABLES

Average Long-term and 2008 Streamflows at Selected Gaging Stations	15
Groundwater-Level Change Statistics by County	16

HYDROGRAPHS

Groundwater-Level Hydrographs	19
-------------------------------------	----

FOREWORD

Nebraskans are proud keepers of their natural resources. This is especially true of groundwater: it is inextricably linked to our rich heritage, maintains our agricultural economy, and provides steady flows to some of the nation's most admired natural streams. The groundwater resources that lie beneath the State are vast yet vulnerable: even small changes in groundwater levels can have profound impacts.

We are proud to present this report, which is a continuation of the series of water resources reports and maps published by the Conservation and Survey Division of the School of Natural Resources. This information can be used to inform, educate, and guide the citizens of Nebraska as we enter new and challenging times regarding water resources.

INTRODUCTION

Groundwater-level information is valuable to citizens and stakeholders for understanding water resource availability and making informed management decisions.

This report is a statewide synthesis of groundwater-level monitoring programs in Nebraska. It is a continuation of the series of annual reports and maps produced by the Conservation and Survey Division (CSD) of the University of Nebraska in cooperation with the U.S. Geological Survey (USGS) since the 1950's. This information is valuable to citizens and stakeholders for understanding water resource availability and making informed management decisions.

Groundwater-level monitoring began in Nebraska in 1930 in an effort to survey the State's groundwater resources and observe changes in its availability on a continuing basis. The CSD and USGS cooperatively developed, maintained, and operated an observation well network throughout the state. For many years, these two agencies were primarily responsible for collecting, storing, and making this information available to the citizens.

Over the years, other agencies began to assume the responsibilities of building and maintaining observation networks and measuring water levels. The CSD and USGS continue to operate some of the original observation wells, but today the majority of measurements are made by agencies such as the U.S. Bureau of Reclamation, the U.S. Fish and Wildlife Service, Public Power and Irrigation Districts, County Extension offices, municipalities, and particularly, Natural Resources Districts. Since these agencies are located throughout the state, they are able to implement groundwater-level monitoring programs using local field staff, landowner contacts, taxing and regulatory authority, and first-hand knowledge of their particular conditions. Collectively, these agencies have developed an extensive network of observation wells throughout the state.


The CSD plays a vital role in providing technical expertise to these agencies as they develop and implement

groundwater-level monitoring plans. The CSD evaluates the adequacy and accuracy of the water-level data and provides the state-wide assessment of groundwater-level changes across all geographic regions and aquifers. Traditionally, CSD has provided technical services to stakeholders by integrating groundwater-level change data with multiple data sets to:

- 1) Determine the amount of groundwater in storage and its availability for use.
- 2) Assess the water-supply outlook by determining changes in the volume of groundwater in storage.
- 3) Identify areas in which changes in groundwater levels may have an economic impact.
- 4) Assist state and local agencies in the formulation and administration of resource-management programs.
- 5) Determine or estimate the rate and direction of groundwater movement, specific yield of aquifers, base flow of streams, sources and amounts of groundwater recharge, and locations and amounts of groundwater discharge.
- 6) Assess the validity of hydrogeologic interpretations and the assumptions used in developing models of a groundwater system.


The need for this information has increased tremendously, yet the resources available for fulfilling this need have decreased. CSD strives to meet this challenge by focusing on fundamental data, building collaborative relationships with the agencies that depend on the information, and providing scientifically accurate information in a timely manner.

Statewide Subdivisions used to Describe Regional Trends



Divisions are based on Natural Resources District boundaries and are consistent with divisions used in previous reports to categorize and describe regional trends.

Important Aquifers and Topographic Regions of Nebraska



Note: In some areas, the aquifer units shown here may contain little or no saturated thickness.

INTRODUCTION (continued)

Purpose and Methods

This report summarizes annual and long-term changes in Nebraska's groundwater levels. These changes are depicted in maps, tables, and graphs. The maps delineate regional trends on a statewide basis. Although localized conditions may vary considerably, the maps presented in this report provide an overview of the general locations, magnitudes, and extents of rises and declines. The tables summarize statewide statistics regarding observation networks and water-level changes. The water-level hydrographs depict changes at a single observation point. Since there are thousands of observation wells in the state, only a few can be shown in this report. The hydrographs were selected based on the length, accuracy, and completeness of their records and whether or not they are generally representative of a particular area and aquifer.

The annual groundwater-level change map in this report was prepared by comparing groundwater levels measured in spring 2009 to levels collected in the same well in the previous year. Data were used from thousands of sites throughout the state. Contours were computer-generated, then manually edited on maps at a scale of 1:500,000 to conform to hydrogeologic boundaries.

For the drought-period map, water levels were compared in wells for which both Spring 2000 and Spring 2009 levels were measured. In order to be consistent with previous versions of this map, contours were drawn manually with the aid of the previous year's map and with knowledge of major hydrogeologic boundaries.

For the predevelopment to spring 2009 map, comparisons were made between spring 2009 water levels and estimated predevelopment water levels. An estimated predevelopment water level is the approximate average water level at a well site prior to any development that significantly affects water levels. Predevelopment water levels for most of the state are the estimated water levels that generally occurred before the 1930s, 1940s, or early-to-mid-1950s. These dates, which vary throughout Nebraska, generally are dependent on the beginning dates of intensive use of groundwater for irrigation. Typically all available water-level data collected prior to or during the early stages of groundwater development are used to estimate predevelopment water levels. Contours were drawn manually with the aid of the previous year's map and with knowledge of major hydrogeologic boundaries.

Factors Causing Groundwater-Level Changes

Groundwater-level changes are a reflection of the changing balance between recharge to, discharge from, and storage in an aquifer. If recharge and discharge are in balance, groundwater levels generally remain unchanged because storage neither increases nor decreases. If, however, the rate of recharge exceeds the rate of discharge, the amount of water stored in the aquifer increases and groundwater levels rise. Conversely, if the rate of discharge exceeds the rate of recharge, the amount of water in storage is depleted and groundwater levels decline. The physical properties of an aquifer affect this balance and control the magnitude by which groundwater levels change and the rate at which these changes are transmitted through an aquifer.

Groundwater levels respond to a variety of natural and man-made factors affecting recharge and discharge. Recharge occurs from precipitation, but also from irrigation return flow and seepage from canals, reservoirs, and streams. Discharge occurs as baseflow to streams and lakes, but also as a result of groundwater pumping and evapotranspiration.


Groundwater-level changes can be observed at many different temporal scales. Changes may occur over several minutes or hours in response to pumping, floods, or earthquakes. Long-term changes may occur due to the cumulative effects of pumping over many irrigation seasons, prolonged droughts or wet periods, or continual seepage from man-made water bodies.

Groundwater levels can be observed at multiple spatial scales. For example, groundwater levels decline around the immediate vicinity of an individual well during pumping, but also from the cumulative effects of many irrigation wells pumped over many irrigation seasons at the scale of an entire regional aquifer. Similarly, groundwater levels rise along the banks of a stream during a flood, but may also rise significantly over an entire drainage basin during a prolonged wet period.


The maps presented in this report were generally mapped at a scale of 1:500,000. They are intended to identify regional trends at medium and long-term time scales throughout the entire state of Nebraska. As such, these changes chiefly reflect the interplay between precipitation, groundwater pumping, and artificial recharge from reservoirs and canals.

Example of Groundwater-Level Changes at Different Temporal Scales

Long-term groundwater-level changes



Seasonal groundwater-level changes



Daily groundwater-level changes



Based on data from Plymouth Recorder well, Jefferson County

STATUS OF GROUNDWATER-LEVEL MONITORING PROGRAM, 2009

Nebraska has developed a massive network of long-term groundwater-level monitoring sites, but our ability to observe short-term fluctuations has decreased sharply because fewer continuous recorder wells are being measured.

The distribution of observation wells in the state is non-uniform in several ways. First of all, the geographic distribution of wells varies tremendously. The number of observation wells per county ranges from 0 in some counties to more than 250 in others, with the greatest density in areas where substantial changes in water levels have occurred. For this reason, water-level changes in some areas may not be detected or delineated because of insufficient data.

Second, the hydrogeologic distribution of wells varies. Wells are completed to different depths and in different geologic materials depending on the type of well, its purpose, and the availability of aquifers at different depths. Hydrogeology is complex, so hydrogeologists must simplify water-level information collected over large regions in order to analyze results. The water-level change maps in this report show changes over a wide range of aquifer materials and hydrogeologic conditions. Significant hydrogeologic boundaries were adhered to where possible, but only a limited amount of information can be obtained regarding interactions between aquifers.

In 2009, approximately 7,482 water-level measurements, measured in nearly 6,019 observation wells, were reported to CSD. Many, but not all, of these measurements were also reported to the USGS. The reason for this discrepancy is that some observation wells maintained by the various agencies have not been added to the USGS database, a process that requires providing basic information about the well and its construction on a standardized well schedule form. The CSD maintains a database that stores all records from all wells reported from cooperating agencies. Currently there are an additional 13,062 records from 2,096 wells in the CSD database that are not part of the USGS database. Since the local agencies making water-level measurements have the best information regarding their observation well sites, resolving this issue will require cooperation between all entities involved in making water-level measurements in Nebraska.


To compile information regarding the status of water-level monitoring in Nebraska in this report, data from the USGS system were combined with data from the CSD database. Every attempt was made to avoid duplication of records, however, some wells and/or measurements might exist in duplicate due to the large number of records involved, limited information regarding changes at monitoring sites, and other factors unknown to the compiler at the time of the report. For example, a large, but unknown number of

fall measurements are not yet recorded in any centralized database.

The number of wells used to measure groundwater levels has increased dramatically since 1930. Significant increases came in the late 1940's, late 1970's, and from 2000 to 2003. Additionally, the number of wells with long term historical records has increased. The increase in the number of wells with at least 10, 20, or 30 years of previous measurements has increased in a manner similar to the total increase. These trends indicate that many of the same observation wells used in the past are being used today. These wells will provide valuable information regarding long-term historical trends at thousands of individual sites.

In contrast to the increasing number of wells being monitored, the frequency of measurements has decreased markedly over the years. Most observation wells in Nebraska are measured in the spring and late fall. Spring measurements are useful in determining amounts of groundwater in storage prior to the irrigation season under static conditions, whereas fall measurements are useful in evaluating the effects of pumping and canal seepage after the irrigation season. Some wells, however, are measured more frequently, providing a detailed record of seasonal changes. The groundwater-level hydrographs shown later in this report illustrate the utility of long-term, continuous recorder wells. These wells are particularly useful for examining seasonal extremes. The decreasing number of wells being measured in this manner is concerning because key information can be obtained from seasonal fluctuations, including the timing and duration of pumping or recharge events, the degree of confinement of an aquifer, and the adequacy of the well used for monitoring (for example, see the Kimball Recorder Well hydrograph on page 30). Such detailed information can be particularly useful for developing and calibrating groundwater models. It is likely that many wells are actually equipped with automatic water-level sensors, but currently the locations and quantity of such wells or whereabouts of this information are unknown to the authors.


Locations of Spring/Fall and Monthly Observation Wells




Sources: U.S. Geological Survey, Nebraska Water Science Center; Nebraska Natural Resources Districts.

Graphs showing Numbers of Groundwater-Level Monitoring Wells Statewide since 1930

A. Number of wells measured each year since 1930



B. Number of recorder wells measured each year since 1930



Although the number of observation wells and wells with long-term records have increased (A), the number of wells being measured more than twice per year has decreased (B).

CHANGES IN GROUNDWATER LEVELS, SPRING 2008 TO SPRING 2009

Groundwater levels rose throughout much of Nebraska due to above-average precipitation in 2008.

Groundwater levels rose throughout much of central and east-central Nebraska from 2008 to 2009. Areas of greater than one foot rise occurred mainly in a broad area extending from the north-central and northeast toward the south-central and southeast. Rises of greater than five feet occurred in large portions of Clay, Fillmore, and York Counties in south-central Nebraska, Gage County in the southeast, Buffalo County in the central, and Platte County in the northeast. Several smaller areas with greater than five foot rises were scattered throughout this region.

Groundwater levels rose in small, widely scattered areas of western Nebraska. The largest of these areas were in Red Willow, Frontier, Chase, and Perkins Counties in the southwest, and Scottsbluff, Morrill, and Sheridan Counties in the panhandle.

Most of the groundwater level rises can be attributed to above-average precipitation for 2008 across much of the State. Much of central Nebraska experienced greater than 130% of the 30-year average precipitation. The abundant moisture likely resulted in reduced pumping for irrigation and, in areas of shallow water table and permeable soils, increased recharge to the aquifer.


Groundwater-level declines occurred in several, widely scattered areas of the State. The two largest areas were in southwestern Nebraska in portions of Keith, Perkins, Dundy, and Chase Counties, where declines ranged from one to more than eight feet.

Declines in the panhandle occurred in Box Butte, Sheridan, and Cheyenne Counties. These declines were generally between one and five feet. Other significant areas experiencing declines were in Custer and Dawson Counties in the central, and Gosper and Phelps Counties in the south-central. Elsewhere in the State, declines occurred in Holt, Colfax, Dixon, Dakota, Thurston, Burt, and Washington Counties in the north-central and northeast.

Below-average precipitation was mostly limited to the panhandle of Nebraska in 2008. Both rises and declines in groundwater levels were observed in this area, although declines appear to have been more widespread than rises. Other areas experiencing groundwater-level declines, such as the southwest and northeast, received near or slightly above-average precipitation for 2008. Areas of south central Nebraska experiencing groundwater-level declines, such as Custer and Dawson Counties, received abundant precipitation in 2008.


Large areas of little to no change were in the western sandhills and portions of the southwest. These areas received near normal precipitation, and are also areas of relatively low irrigation well density. Areas along major river valleys also were generally unchanged because groundwater-level declines were supplemented by river seepage and rises were limited to the elevation of the discharge points along river beds.

Percent of Normal Precipitation - January 2008 to January 2009



*Sources: National Climate Data Center, Asheville, North Carolina;
High Plains Regional Climate Center, University of Nebraska–Lincoln.*

Groundwater-level Changes in Nebraska - Spring 2008 to Spring 2009



Sources: U.S. Geological Survey, Nebraska Water Science Center; U.S. Bureau of Reclamation, Kansas-Nebraska Area Office; Nebraska Natural Resources Districts; Central Nebraska Public Power and Irrigation District.

CHANGES IN GROUNDWATER LEVELS, SPRING 2000 TO SPRING 2009

Groundwater levels remain below pre-drought levels throughout most of the state, but some groundwater-level rises are occurring in the central and northeast.


Nebraska experienced a widespread multi-year drought during the first part of this decade. By 2006-2007, groundwater levels had declined by at least one foot over almost the entire State, and at least 5 feet to more than 25 feet in areas of major irrigation development (Burbach, 2006; 2007). Although the drought had ended by the beginning of 2007 in the eastern part of the State, and by the beginning of 2009 in the west, groundwater levels remained below pre-drought levels throughout most of the State. As of Spring 2009, widespread areas of 5 or more feet of decline still existed in the southwest, south-central, the panhandle, and the north-central. These areas generally received below-average precipitation from 2000-2009.

Declines in the southwest were most severe in Chase, Dundy, Lincoln, and Perkins Counties. Declines of more than 25 feet occurred in the vicinity of Lake McConaughy in Keith County. Declines in the panhandle were most severe in Cheyenne County near Sidney, in Box Butte County, and in Sheridan County near Gordon. The largest declines in the

east north-central were northwest of O’Niell in Holt County. Declines of up to 15 feet occurred in central Colfax County in the northeast. The east south-central was mostly below pre-drought levels in 2009, with the largest declines occurring in Hamilton, Polk, northern Clay, and northern York Counties. Southern Gage County experienced the largest declines in the southeast, and declines in the central were greatest in Buffalo County and small portions of Custer, Dawson, and Hall Counties.


Spring 2009 groundwater levels rose above Spring 2000 levels in some areas. Areas of Holt, Wheeler, Antelope, Boone, Madison, Pierce, and Wayne Counties in the northeast, and Valley and Sherman Counties in the central had experienced rises of between 1 and 5 feet, with a few areas of greater than 5 feet rise, such as Madison and Valley Counties. These areas received mostly below-average precipitation during the drought (Burbach, 2007), but by 2009 had received enough rainfall that the 9-year period was above the 30-year average.

Percent of Normal Precipitation - January 2000 to January 2009



Sources: National Climate Data Center, Asheville, North Carolina;
High Plains Regional Climate Center, University of Nebraska–Lincoln.

Groundwater-level Changes in Nebraska - Spring 2000 to Spring 2009



Sources: U.S. Geological Survey, Nebraska Water Science Center; U.S. Bureau of Reclamation, Kansas-Nebraska Area Office; Nebraska Natural Resources Districts; Central Nebraska Public Power and Irrigation District.

CHANGES IN GROUNDWATER LEVELS, PREDEVELOPMENT TO SPRING 2009

Long-term groundwater-level changes in Nebraska primarily reflect aquifer depletion in areas of dense irrigation development and increases in storage due to seepage from canals and reservoirs.

Spring 2009 groundwater levels continue to indicate long-term declines and rises in certain areas of Nebraska. The largest groundwater-level declines from predevelopment to spring 2009 occurred in the southwest and in the panhandle. The largest rises occurred in the central and west south-central.

The predevelopment groundwater levels used in the southwest are representative of the approximate average water levels prior to 1953. Available data indicate that, as a result of intensive use of groundwater for irrigation, a general trend of declining water levels began in about 1966 in Chase, Dundy, and Perkins counties. The area of decline extends into portions of Keith, Lincoln, and Hayes Counties. Declines of as much as 70 feet occurred in Chase County since predevelopment.

Predevelopment water levels used to develop the groundwater-level change map in Box Butte County are the approximate average water levels prior to 1946. Intensive groundwater development for irrigation since 1950 has caused water-levels to decline 5 to more than 70 feet from predevelopment levels.

A large portion of east south-central Nebraska has experienced long-term groundwater-level declines since predevelopment. Predevelopment water levels in this area are generally representative of the approximate average water levels prior to 1950. Groundwater levels in large parts of Webster, Adams, Clay, Fillmore, Nuckolls, and Thayer Counties have declined more than 10 feet, and in some areas nearly 30 feet, from predevelopment. Areas of Hamilton, York, Seward, Polk, and Butler have declined at least 5 feet from predevelopment. The areal extent and magnitude of the groundwater-level declines have improved somewhat due to recent above-average precipitation. The most notable improvement was in northeastern Clay County where much of the area that was previously 5 to 10 feet below predevelopment is now less than 5 feet below. The declines in Hamilton, York, Seward, Polk, and Butler are much less severe than in recent years.

Parts of other counties that experienced relatively large areas of decline include Buffalo, Custer, and Dawson in the central; Harlan and Franklin in the south-central; Hitchcock, Frontier, and Red Willow in the southwest; Banner, Kimball,

Morrill, Cheyenne, and Sheridan in the panhandle; and Holt in the north central. A newly developing area of decline is located in central Colfax County in the northeastern part of the State. Many of these areas occur in areas of intense groundwater irrigation, but in other areas, wells are not particularly dense. Other factors such as aquifer characteristics, rates of recharge, and irrigation scheduling could be contributing to the declines.


Groundwater-level rises from predevelopment generally occurred in areas of surface irrigation systems. Storage of water in Lake McConaughy began in 1941, and seepage losses caused water-level rises of as much as 60 feet in nearby observation wells (Ellis and Dreeszen, 1989). Water levels generally stabilized by about 1950 and since then have fluctuated in response to changes in reservoir levels and precipitation (Johnson and Pederson, 1984).

Water released from storage in Lake McConaughy and subsequently diverted from the Platte River near North Platte has been used for irrigation, primarily in Gosper, Keamey, and Phelps counties, since 1941. Deep percolation of water from these irrigation-distribution systems and from excess water applied to crops has raised water levels 10 to 50 feet or greater from predevelopment levels in an area extending from southeastern Keith County in the west to central Kearney County in the east. Rises in Gosper, Phelps, and Kearney Counties exceed 50 feet, and in some areas are as much as 136 feet.

Water-level rises of 5 to 30 feet have occurred south of the South Platte and Platte rivers in Keith, Lincoln, and Dawson counties. Seepage from Sutherland Reservoir, Lake Maloney, and their associated canals caused water levels to start rising south and west of North Platte in about 1935. East of North Platte, water levels began rising in about 1940 as a result of seepage from the Tri-County Supply Canal and Jeffrey Reservoir.


Rises of 10 to more than 60 feet occurred in portions of Howard, Sherman, Valley, and Greeley Counties in the central. The water-level rises in this area are the result of seepage from irrigation canals, seepage from Sherman Reservoir, and deep percolation of irrigation water applied to crops.

Density of Active Registered Irrigation Wells - January 2009



Source: Nebraska Department of Natural Resources.

Groundwater-level Changes in Nebraska - Predevelopment to Spring 2009



Sources: U.S. Geological Survey, Nebraska Water Science Center; U.S. Bureau of Reclamation, Kansas-Nebraska Area Office; Nebraska Natural Resources Districts; Central Nebraska Public Power and Irrigation District.

AVERAGE DAILY STREAMFLOWS, 2008


Streamflows in 2008 varied from 0% to more than 200% of the long-term average, reflecting regional trends in precipitation and long-term effects on baseflow.

The flows in Nebraska streams have several different sources. Snowmelt in the Rocky Mountains west of Nebraska provides springtime flows for the Platte River as it enters Nebraska. Variations in the amount of winter snowpack have a profound impact on discharges, but so also can the timing and amount of releases from dams in Nebraska, Wyoming, and Colorado. Runoff from precipitation is the source of many of the peak flows in Nebraska streams. Runoff is greatest on soils with low infiltration rates and/or high slopes. As such, many streams in eastern Nebraska have 'flashy' discharges characterized by high flows immediately following large precipitation events. Streams with headwaters in the sandhills are characterized by steady flows year-round because high infiltration rates in the sandy soils limit runoff and provide constant groundwater discharge to streams.

Average daily streamflow values varied tremendously across the state in 2008. Flows were well above the long-term average over much of the state due to above-average precipitation. In parts of western Nebraska, where precipitation was near or below average, flows were below the long-term average. The reduced baseflow to these streams is part of a regional trend in the High Plains Aquifer of reduced baseflows due to lowering of the regional water table (Sophocleous, 1998).

The factors affecting streamflows are numerous and complex. Nonetheless, it is commonly known that in areas where streams are well-connected to aquifers, groundwater-level changes can have an effect on baseflows. Continued monitoring of groundwater-level changes throughout Nebraska is necessary in order to evaluate and manage these interconnected resources.

Average Streamflow in 2008, as a Percentage of the Long-term Average



Average Long-term and 2008 Streamflows at Selected Gaging Stations

Stream	long-term average (acre-feet)	2008 (acre-feet)	2008 (% of long-term average)
Sappa Creek nr Stamford	9,720	5,510	57
Red Willow Creek nr Red Willow	7,640	11,900	156
Republican River nr Orleans	109,000	171,000	157
Frenchman River at Culbertson	32,000	33,800	106
Republican River nr Hardy	185,000	207,000	112
Courtland Canal at Kan-Neb line	54,800	25,900	47
Prairie Dog Creek nr Woodruff, KS	5,610	3,270	58
Beaver Creek at Cedar Bluffs, KS	2,060	1,200	58
Haigler (Pioneer) Ditch at Colo-Neb line	5,340	5,410	101
Arikaree River at Haigler (minus spillback)	2,640	240	9
Republican River at Colo-Neb line	25,080	21,070	84
South Platte River at Julesburg, CO	491,000	93,900	19
Lodgepole Creek at Bushnell	2,700	0	0
North Platte River at Wyo-Neb line	1,060,000	859,000	81
Pumpkin Creek near Bridgeport	7,890	730	9
Platte River at Louisville	5,890,000	7,640,000	130
Platte River nr Grand Island	1,310,000	850,000	65
North Platte River at Lewellen	929,000	571,000	61
(Combo) Loup River & Canal nr Genoa	1,900,000	2,000,000	105
South Loup River at St. Michael	160,000	170,000	106
Dismal River nr Thedford	154,000	158,000	103
North Loup River at Taylor	382,000	392,000	103
Niobrara River at Wyo-Neb line	2,190	1,840	84
Niobrara River nr Verdel	1,390,000	1,460,000	105
Keya Paha River at Wewela, SD	68,000	67,900	100
White River nr SD-Neb line	27,700	5,480	20
Little Blue River nr Deweese	100,000	116,000	116
W. Fk. Big Blue River nr Dorchester	137,000	186,000	136
Little Blue River at Hollenberg, KS	369,000	523,000	142
Big Blue River at Barneston	701,000	1,060,000	151
Elkhorn River at Ewing	162,000	185,000	114
Logan Creek at Pender	159,000	288,000	181
Maple Creek nr Nickerson	71,100	97,300	137
No. Fk. Elkhorn River nr Pierce	90,000	188,000	209
Big Nemaha River at Falls City	455,000	575,000	126
Little Nemaha River at Auburn	250,000	493,000	197

GROUNDWATER-LEVEL CHANGE STATISTICS

Groundwater-level change statistics can be used to make generalized comparisons between time periods and counties

The following table summarizes groundwater-level changes by county for Spring 2008 to Spring 2009, Spring 2000 to Spring 2009, and predevelopment to Spring 2009. Since counties are political boundaries, water-level change statistics shown here may reflect conditions averaged over more than one hydrogeologic unit. Counties are readily recognizable boundaries, however, and can be used to generalize changes on a local level for comparison purposes. Ideally, groundwater-level change statistics should be summarized by individual aquifers throughout the state. Such an assessment requires detailed knowledge of the screened

interval of the well and the local geology. This information is not always available, and would require analysis of thousands of additional wells. This process is on-going.

In the table below, 'Count' is the number of observation wells used to compute the statistics, 'Min' is the minimum water-level change during that time period in an individual well, 'Max' is the maximum water-level change during that time period in an individual well, and 'Average' is the average water-level change during that time period computed from all wells in the county.

Groundwater-Level Change Statistics by County

County	Spring 2008 to Spring 2009				Spring 2000 to Spring 2009				Predevelopment to Spring 2009			
	Count	Min	Max	Average	Count	Min	Max	Average	Count	Min	Max	Average
Adams	82	-2.2	3.6	1.1	58	-10.2	3.2	-6.2	59	-27.8	0.5	-10.5
Antelope	68	-2.4	6.9	1.7	36	-8.6	9.0	1.0	34	-5.3	9.7	2.2
Arthur	19	-1.1	0.9	0.0	19	-3.8	-0.3	-1.5	13	-11.7	5.3	-2.2
Banner	93	-2.8	5.7	-0.4	64	-30.1	3.9	-8.0	5	-80.8	-8.6	-24.6
Blaine	12	-0.7	2.9	1.4	12	-2.9	1.0	-0.5	12	-2.2	4.0	1.5
Boone	29	-0.6	4.4	2.3	24	-3.4	10.1	2.2	22	-2.8	15.1	5.3
Box Butte	74	-6.1	16.6	-0.4	43	-21.8	2.9	-10.3	50	-88.4	-1.0	-35.5
Boyd	8	-5.1	1.0	-0.4	8	-7.9	0.4	-2.8	7	-3.5	0.1	-1.6
Brown	40	-4.3	2.7	0.7	34	-7.0	1.5	-2.3	20	-3.6	17.2	3.4
Buffalo	82	-2.2	7.0	1.7	68	-14.2	1.4	-5.4	75	-26.9	4.1	-6.3
Burt	31	-4.7	0.7	-1.5	31	-12.6	3.0	-2.5	30	-19.3	10.3	-2.1
Butler	70	-2.4	7.0	1.7	50	-26.1	2.5	-6.6	48	-22.1	5.8	-4.9
Cass	26	-3.9	3.3	1.4	6	-2.4	8.8	0.7	6	-6.0	-0.6	-3.0
Cedar	29	-1.4	3.0	0.7	25	-21.0	2.7	-1.7	28	-9.5	8.8	0.9
Chase	160	-3.1	2.9	0.0	160	-21.9	0.2	-11.5	122	-73.9	-0.2	-37.0
Cherry	75	-10.0	3.3	0.5	56	-13.1	2.3	-2.2	56	-15.1	30.1	0.2
Cheyenne	84	-5.0	2.5	-0.8	52	-28.9	2.8	-13.8	48	-42.6	6.4	-18.2
Clay	87	-0.8	16.7	2.7	73	-17.4	-1.0	-7.9	69	-20.7	4.2	-9.5
Colfax	32	-7.2	1.7	-0.1	29	-15.4	1.6	-3.6	30	-15.7	3.3	-2.8
Cuming	26	-28.7	1.9	-1.0	21	-57.4	6.6	-3.1	26	-25.3	50.2	2.2
Custer	91	-4.9	3.2	0.6	84	-36.0	15.0	-3.1	85	-44.1	8.9	-4.4
Dakota	2	-1.4	-0.1	-0.7	2	-6.3	-3.2	-4.7	2	-12.5	-3.2	-7.9
Dawes	9	-1.9	9.2	0.9	6	-4.3	-0.3	-2.0	6	-7.0	0.1	-3.4
Dawson	170	-10.2	5.3	0.8	160	-16.1	3.7	-3.3	135	-22.1	33.5	-0.7
Deuel	23	-5.8	3.4	-0.1	17	-10.6	3.1	-3.5	13	-19.0	0.2	-5.8
Dixon	11	-1.9	4.0	0.2	11	-2.4	1.6	-0.4	11	-6.2	8.8	0.4
Dodge	52	-3.4	7.2	0.7	52	-7.3	4.3	-1.7	52	-10.2	10.0	-1.5
Douglas	0	na	na	na	0	na	na	na	0	na	na	na
Dundy	104	-7.2	2.8	-0.7	101	-27.8	3.0	-11.4	83	-63.0	0.4	-34.8
Fillmore	102	-3.4	11.1	2.6	96	-14.3	13.8	-5.0	89	-37.9	18.5	-12.2
Franklin	51	-0.5	2.9	0.5	21	-6.2	0.8	-3.0	37	-12.6	0.5	-4.3
Frontier	36	-8.6	5.4	0.6	35	-8.2	3.6	-0.1	28	-18.8	2.7	-5.4
Furnas	48	-3.3	6.2	0.6	14	-7.5	7.2	-2.8	23	-18.3	5.6	-2.7
Gage	27	-4.3	9.5	2.7	22	-18.7	-0.1	-4.5	25	-12.0	15.6	-1.3
Garden	99	-2.6	1.4	0.1	66	-11.6	1.6	-2.1	30	-14.7	1.9	-3.0
Garfield	29	-0.7	6.5	0.6	13	-5.9	3.0	-0.7	12	0.3	11.4	4.9
Gosper	129	-7.0	4.1	-0.2	102	-65.2	38.7	-5.8	37	-17.4	136.7	36.2

County	Spring 2008 to Spring 2009				Spring 2000 to Spring 2009				Predevelopment to Spring 2009			
	Count	Min	Max	Average	Count	Min	Max	Average	Count	Min	Max	Average
Grant	4	-0.3	1.4	0.8	2	-1.9	-0.6	-1.2	4	-2.4	0.9	-0.8
Greeley	56	-1.3	2.9	0.6	18	-4.7	1.5	-1.4	19	-1.9	15.3	5.6
Hall	97	-1.5	5.6	1.5	79	-13.9	10.6	-3.7	79	-10.8	15.5	-1.7
Hamilton	125	-0.3	6.9	2.3	124	-18.8	0.1	-8.1	121	-14.6	8.0	-5.5
Harlan	95	-8.4	12.0	1.3	68	-10.9	18.7	-3.5	68	-18.0	9.0	-6.7
Hayes	21	-1.8	1.4	0.0	21	-14.5	0.0	-5.9	20	-21.1	-1.6	-9.8
Hitchcock	24	-1.7	1.1	-0.1	25	-10.4	1.0	-1.8	12	-23.5	1.6	-7.0
Holt	138	-14.5	5.6	0.4	78	-17.2	4.3	-3.3	85	-32.3	10.1	-4.8
Hooker	7	-0.4	1.2	0.4	7	-2.7	-0.1	-2.1	7	1.3	7.5	3.7
Howard	109	-5.4	15.6	1.0	34	-9.1	3.0	-1.7	35	-8.5	63.9	5.3
Jefferson	41	-2.6	6.7	1.7	28	-13.0	1.6	-4.3	28	-14.0	1.1	-3.3
Johnson	41	-19.2	3.3	0.2	13	-7.3	3.7	-1.2	20	-10.2	0.9	-2.9
Kearney	83	-3.7	4.7	0.2	59	-13.9	1.9	-5.4	52	-14.9	58.5	5.9
Keith	86	-8.9	5.9	-0.4	89	-46.5	1.8	-6.4	42	-32.2	80.7	4.1
Keya Paha	30	-4.0	6.3	0.3	15	-9.3	-0.4	-3.5	21	-1.0	12.2	3.1
Kimball	29	-4.1	12.8	0.1	20	-25.4	5.2	-6.0	20	-32.4	9.8	-9.3
Knox	25	-3.1	2.6	0.9	21	-11.8	3.3	-2.1	23	-12.2	11.4	2.4
Lancaster	80	-7.4	25.3	1.5	35	-24.8	3.6	-2.0	32	-10.4	20.4	-1.4
Lincoln	86	-3.0	6.4	0.4	85	-16.9	7.1	-4.0	59	-19.2	27.9	0.2
Logan	7	-1.9	1.6	0.4	6	-5.6	-0.1	-3.4	7	-7.9	3.5	-0.6
Loup	36	-0.6	2.6	0.3	10	-4.9	3.5	-1.3	10	-3.3	9.7	1.5
Madison	35	-0.9	3.8	1.5	35	-2.3	8.0	2.7	35	-6.1	19.4	6.8
McPherson	16	-1.6	2.7	0.5	13	-4.8	2.2	-1.8	13	-0.8	5.9	2.0
Merrick	67	-2.3	5.5	0.9	61	-5.2	2.6	0.3	61	-16.1	2.4	-1.2
Morrill	257	-3.2	4.1	0.0	109	-16.3	7.1	-3.4	25	-42.3	17.5	-3.5
Nance	53	-3.0	4.7	0.7	30	-6.3	1.9	-0.6	33	-8.9	26.6	4.4
Nemaha	7	-2.0	1.2	-0.9	1	3.2	3.2	3.2	1	-0.7	-0.7	-0.7
Nuckolls	46	-0.6	5.1	1.4	27	-12.7	3.2	-4.6	36	-19.4	2.7	-8.6
Otoe	27	-21.0	3.9	-0.2	4	-4.4	35.6	11.3	3	-12.2	-4.5	-8.7
Pawnee	0	na	na	na	0	na	na	na	0	na	na	na
Perkins	146	-9.3	4.8	-0.7	146	-28.1	1.6	-11.7	91	-53.6	-2.8	-32.2
Phelps	109	-5.3	2.2	0.0	85	-13.4	5.8	-4.1	70	-34.3	93.7	25.2
Pierce	61	-1.5	3.8	1.0	58	-5.8	6.0	0.2	61	-14.8	12.2	3.8
Platte	67	-1.4	6.4	1.9	38	-13.5	2.6	-2.8	40	-4.3	19.1	2.5
Polk	56	-4.0	11.4	2.0	56	-13.1	1.7	-6.3	56	-11.4	22.9	-2.4
Red Willow	24	-4.1	4.4	0.4	24	-9.3	2.3	-2.3	21	-15.2	-1.6	-7.0
Richardson	7	-12.9	0.0	-3.8	5	-7.6	-0.5	-5.6	4	-14.8	0.0	-8.3
Rock	36	-9.5	3.8	1.2	13	-15.8	2.5	-1.6	18	-13.8	5.4	-0.8
Saline	35	-8.5	8.3	1.4	32	-16.0	0.3	-5.6	28	-14.5	-0.7	-4.8
Sarpy	18	-2.9	1.6	-0.8	15	-1.0	17.1	7.3	1	-2.2	-2.2	-2.2
Saunders	102	-4.8	5.8	1.2	73	-7.4	4.6	-0.4	56	-9.6	5.3	-0.4
Scotts Bluff	225	-9.8	6.9	0.1	204	-15.5	10.3	-2.1	67	-9.4	44.4	4.8
Seward	82	-7.3	5.0	1.8	74	-11.1	6.4	-6.2	73	-10.6	4.4	-5.1
Sheridan	65	-2.5	2.5	-0.5	33	-27.5	0.4	-8.8	34	-26.4	2.1	-14.1
Sherman	18	-0.9	5.0	1.0	18	-7.3	5.0	-0.2	17	-14.5	46.8	15.2
Sioux	82	-2.4	0.7	-0.4	49	-13.8	1.6	-6.7	31	-5.1	61.7	17.5
Stanton	21	-2.3	2.3	0.5	20	-5.5	4.8	-1.1	21	-6.7	6.7	-0.1
Thayer	97	-0.8	4.2	1.3	68	-17.2	0.8	-3.4	83	-26.3	-0.2	-7.8
Thomas	9	-1.2	1.7	0.1	9	-4.3	-1.8	-3.3	9	-0.7	3.0	1.8
Thurston	9	-9.0	0.6	-1.9	8	-4.5	2.9	-0.2	9	-8.4	6.8	-0.8
Valley	103	-9.4	12.9	1.0	22	-3.3	10.6	2.6	17	-5.5	41.2	12.4
Washington	7	-4.5	0.3	-1.3	7	-2.2	3.2	-0.5	4	-3.2	13.0	1.4
Wayne	21	-2.5	3.7	0.3	19	-4.1	3.2	0.8	21	-4.5	10.8	3.2
Webster	53	-0.8	4.1	1.1	17	-15.7	8.4	-2.5	26	-16.5	13.2	-2.5
Wheeler	22	-0.2	4.2	1.7	15	-3.6	4.5	0.7	15	-2.6	10.9	3.4
York	124	-0.7	7.8	2.4	120	-19.4	-0.5	-8.8	118	-20.0	1.6	-6.0

GROUNDWATER-LEVEL HYDROGRAPHS

Groundwater-level hydrographs provide detailed information about seasonal and long-term stresses on aquifers.

Hydrographs are used to illustrate seasonal and long-term water-level fluctuations in wells that represent hydrologic conditions at various locations throughout the State. The observation wells selected include those with long-term records and continuous or semi-continuous measurements. Throughout much of the history of these wells, a float system was used to detect depths to water and record these depths graphically on a chart or digitally on punched tape. Since only a limited amount of data can be stored, only the lowest daily value for about every fifth day was recorded in the archives. More recently, however, submersible pressure transducers have replaced the old float systems for taking automatic measurements. Pressure transducers work by sensing the hydrostatic pressure on the device and automatically calculating the amount of water above the sensor. The depth to water is obtained by subtracting the height of the water column above the sensor from the depth of the sensor. This data can be collected at user-defined intervals and downloaded directly to a computer.

Hydrographs provide a detailed record of water-level

changes in observation wells. It is not practical to record continuous measurements in all observation wells in the state. By selecting a few key wells in certain locations as “index” wells, important short-term changes can be identified. Hydrographs of wells measured only in the spring and fall may not show extremes in water-level fluctuations, but they provide important information on long-term water-level trends.

The time periods shown in the hydrographs in this report vary from well-to-well. So too do the ranges in fluctuations. The top and bottom of the hydrographs usually do not correspond to the top or bottom of the aquifer. Rather, they represent general maximum and minimum water levels measured in each individual well. Water-level changes may appear to fluctuate within similar ranges in all wells, but upon closer examination, it becomes apparent that the water-level range represented in each hydrograph is different.

In the following hydrographs, elevation is in feet above sea level (ft. a.s.l.) and predevelopment water levels are in feet below ground surface (ft. b.g.s.).

Adams County

Roseland Recorder


ID#: 402910098352101

Legal: 6-11W-17 CB

Elevation: 1980 ft. a.s.l.

Estimated predevelopment

water level: 77 ft. b.g.s.



Hastings Recorder


ID#: 403403098244001

Legal: 7-10W-23 AB

Elevation: 1927 ft. a.s.l.

Estimated predevelopment

water level: 99.96 ft. b.g.s.



Antelope County

Elgin Recorder


ID#: 415559098005201

Legal: 23-6W-28 DC

Elevation: 1922 ft. a.s.l.

Estimated predevelopment

water level: 101 ft. b.g.s.



Box Butte County

Alliance Recorder


ID#: 420904102525201

Legal: 25-48W-12 CCA

Elevation: 3950 ft. a.s.l.

Estimated predevelopment


water level: 52 ft. b.g.s.



Buffalo County


Riverdale Recorder

ID#: 405137099085201
Legal: 10-16W-5 DC
Elevation: 2262 ft. a.s.l.
Estimated predevelopment
water level: 106 ft. b.g.s.



Gibbon Recorder


ID#: 404618098504401
Legal: 9-14W-1 DC
Elevation: 2060 ft. a.s.l.
Estimated predevelopment
water level: 17 ft. b.g.s.



Butler County


Dwight Recorder

ID#: 410612096592601
Legal: 13-4E-17 AB
Elevation: 1605 ft. a.s.l.
Estimated predevelopment
water level: 160 ft. b.g.s.



Rising City Recorder


Rising City Recorder
ID#: 411420097173002
Legal: 15-1E-27 DDDD
Elevation: 1618 ft. a.s.l.
Estimated predevelopment
water level: 103 ft. b.g.s.



Chase County


Lamar Recorder

ID#: 403516101560601
Legal: 7-41W-11 DAA
Elevation: 3506 ft. a.s.l.
Estimated predevelopment
water level: 48 ft. b.g.s.




Champion Recorder

Champion Recorder
ID#: 402757101591201
Legal: 6-41W-21 CCC
Elevation: 3513 ft. a.s.l.
Estimated predevelopment
water level: 30 ft. b.g.s.



Imperial Recorder


Imperial Recorder
ID#: 403235101395501
Legal: 7-38W-29 CBB
Elevation: 3290 ft. a.s.l.
Estimated predevelopment
water level: 56 ft. b.g.s.



Cheyenne County


Gurley Recorder

Gurley Recorder
ID#: 412100102592401
Legal: 16-49W-19 BBB
Elevation: 4281 ft. a.s.l.
Estimated predevelopment
water level: 221 ft. b.g.s.




Clay County

Harvard Recorder
ID#: 403910098051401
Legal: 8-7W-23 BB
Elevation: 1812 ft. a.s.l.
Estimated predevelopment
water level: 79 ft. b.g.s.




Glenville Recorder
ID#: 402806098132501
Legal: 6-8W-21 DD
Elevation: 1800 ft. a.s.l.
Estimated predevelopment
water level: 93 ft. b.g.s.




Dawson County

Overton Recorder
ID#: 404553099341301
Legal: 9-20W-10 ADDD
Elevation: 2327 ft. a.s.l.
Estimated predevelopment
water level: n.a.




Lexington Recorder
ID#: 404949099445701
Legal: 10-21W-18 DDD
Elevation: 2421 ft. a.s.l.
Estimated predevelopment
water level: 12 ft. b.g.s.




Dawson County

Gothenburg Recorder
ID#: 405445100074001
Legal: 11-25W-24 BCB
Elevation: 2545 ft. a.s.l.
Estimated predevelopment
water level: 5 ft. b.g.s.




Dundy County


Enders Recorder
ID#: 401703101394801
Legal: 4-38W-30 BCC
Elevation: 3317 ft. a.s.l.
Estimated predevelopment
water level: 46 ft. b.g.s.



Haigler (New) Recorder
ID#: 400155101521302
Legal: 1-40W-29 BB
Elevation: 3025 ft. a.s.l.
Estimated predevelopment
water level: 13 ft. b.g.s.




Lamont Recorder
ID#: 401401101510701
Legal: 3-40W-16 BBB
Elevation: 3458 ft. a.s.l.
Estimated predevelopment
water level: n.a.




Dundy County

Benkelman Recorder
ID#: 400852101352701
Legal: 2-38W-10 DD
Elevation: 3265 ft. a.s.l.
Estimated predevelopment
water level: 84 ft. b.g.s.




Fillmore County


Exeter Recorder
ID#: 403800097300701
Legal: 8-2W-26 AD
Elevation: 1610 ft. a.s.l.
Estimated predevelopment
water level: 24 ft. b.g.s.



Shickley Recorder
ID#: 402504097432201
Legal: 5-4W-12 BDC
Elevation: 1651 ft. a.s.l.
Estimated predevelopment
water level: 82 ft. b.g.s.



Burress Recorder
ID#: 403356097275602
Legal: 7-1W-19 AA
Elevation: 1615 ft. a.s.l.
Estimated predevelopment
water level: 57 ft. b.g.s.



Franklin County


Upland Recorder

ID#: 401735098522701

Legal: 4-14W-23 CCDA

Elevation: 2120 ft. a.s.l.

Estimated predevelopment
water level: 170 ft. b.g.s.



Frontier County


Orafino Recorder

ID#: 403042100093201

Legal: 6-25W-4 DD

Elevation: 2500 ft. a.s.l.

Estimated predevelopment
water level: 65 ft. b.g.s.



Gage County


Ellis Recorder

ID#: 400917096525101

Legal: 2-5E-8 AD

Elevation: 1360 ft. a.s.l.

Estimated predevelopment
water level: 70 ft. b.g.s.




DeWitt Recorder

ID#: 402155096523101

Legal: 5-5E-32 AAAA

Elevation: 1278 ft. a.s.l.


Estimated predevelopment
water level: 12 ft. b.g.s.



Hall County


Alda (New) Recorder
ID#: 405315098304302
Legal: 11-11W-25 CC
Elevation: 1924 ft. a.s.l.
Estimated predevelopment
water level: 15 ft. b.g.s.

Alda (New) Recorder




Doniphan Recorder
ID#: 404513098181201
Legal: 9-9W-14 AAB
Elevation: 1915 ft. a.s.l.
Estimated predevelopment
water level: 53 ft. b.g.s.

Doniphan Recorder



Cameron Recorder
ID#: 405553098363001
Legal: 11-12W-12 DDD
Elevation: 1942 ft. a.s.l.
Estimated predevelopment
water level: 21 ft. b.g.s.


Cameron Recorder



Hamilton County

Aurora Recorder
ID#: 404836097584101
Legal: 10-6W-27 ACAA
Elevation: 1791 ft. a.s.l.
Estimated predevelopment
water level: 91 ft. b.g.s.


Aurora Recorder



Hamilton County


Kronberg Recorder

ID#: 405921097514701
Legal: 12-5W-23 CCC
Elevation: 1774 ft. a.s.l.
Estimated predevelopment
water level: 81 ft. b.g.s.



Murphy Recorder


Murphy Recorder
ID#: 405648098064201
Legal: 11-7W-3 CCB
Elevation: 1846 ft. a.s.l.
Estimated predevelopment
water level: 85 ft. b.g.s.



Harlan County


Ragan Recorder

ID#: 401857099195201
Legal: 4-18W-15 AD
County: Harlan
Elevation: 2290 ft. a.s.l.
Estimated predevelopment
water level: 173 ft. b.g.s.



Alma Recorder


Alma Recorder
ID#: 400920099215501
Legal: 2-18W-9 BCC
Elevation: 2120 ft. a.s.l.
Estimated predevelopment
water level: n.a.



Hitchcock County

Palisade Recorder


ID#: 401548101080501
Legal: 4-34W-34 DDD
Elevation: 2975 ft. a.s.l.
Estimated predevelopment
water level: 187 ft. b.g.s.



Holt County


Atkinson Recorder

ID#: 423730098560001
Legal: 31-14W-27 DDD
Elevation: 2080 ft. a.s.l.
Estimated predevelopment
water level: 30 ft. b.g.s.




O'Neill Recorder

O'Neill Recorder
ID#: 423148098300601
Legal: 30-10W-32 DAA
Elevation: 1952 ft. a.s.l.
Estimated predevelopment
water level: 31 ft. b.g.s.



Chambers Recorder


Chambers Recorder
ID#: 421210098402001
Legal: 26-12W-26 AAA
Elevation: 2060 ft. a.s.l.
Estimated predevelopment
water level: 6 ft. b.g.s.



Jefferson County

Fairbury Recorder

ID#: 400813097112401
Legal: 2-2E-15 CAD
Elevation: 1314 ft. a.s.l.
Estimated predevelopment
water level: 9 ft. b.g.s.




Daykin Recorder

ID#: 401626097210701
Legal: 4-1E-31 AA
Elevation: 1480 ft. a.s.l.
Estimated predevelopment
water level: 73.5 ft. b.g.s.



Plymouth Recorder


ID#: 401837097015301
Legal: 4-3E-13 DA
Elevation: 1440 ft. a.s.l.
Estimated predevelopment
water level: 107 ft. b.g.s.



Johnson County

Cook Recorder


ID#: 403032096104801
Legal: 6-11E-4 DDDD
County: Johnson
Elevation: 1070 ft. a.s.l.
Estimated predevelopment
water level: n.a.



Keya Paha County

Springview Recorder
ID#: 424837099425201
Legal: 33-20W-19 DCDD
Elevation: 2446 ft. a.s.l.
Estimated predevelopment
water level: n.a.


Springview Recorder



Kimball County

Kimball Recorder
ID#: 411739103401501
Legal: 15-55W-7 ABB
Elevation: 4874 ft. a.s.l.
Estimated predevelopment
water level: 210 ft. b.g.s.


Kimball Recorder



Lancaster County


Princeton Recorder
ID#: 403400096435501
Legal: 7-6E-15 DCCC
Elevation: 1450 ft. a.s.l.
Estimated predevelopment
water level: n.a.

Princeton Recorder



Van Dorn Recorder

Van Dorn Recorder
ID#: 404706096413001
Legal: 10-6E-36 CDD
Elevation: 1200 ft. a.s.l.
Estimated predevelopment
water level: 68 ft. b.g.s.



Lincoln County


Dickens Recorder

ID#: 404742101010801

Legal: 10-33W-36 ADD

Elevation: 3089 ft. a.s.l.

Estimated predevelopment
water level: 105 ft. b.g.s.



Farnam Recorder


Farnam Recorder

ID#: 404233100132301

Legal: 9-26W-36 ADD

Elevation: 2785 ft. a.s.l.

Estimated predevelopment
water level: 228 ft. b.g.s.



Hershey Recorder


Hershey Recorder

ID#: 410250101004201

Legal: 12-32W-6 BAB

Elevation: 3088 ft. a.s.l.

Estimated predevelopment
water level: n.a.



Lake Maloney Recorder


Lake Maloney Recorder

ID#: 405732100531201

Legal: 11-31W-5 BBB

Elevation: 3081 ft. a.s.l.


Estimated predevelopment
water level: n.a.



Merrick County


Archer Recorder

ID#: 410618098113401
Legal: 13-8W-11 DDDD
Elevation: 1772 ft. a.s.l.
Estimated predevelopment
water level: 3.5 ft. b.g.s.



Central City Recorder


Central City Recorder
ID#: 410943097575001
Legal: 14-6W-26 AAAA
Elevation: 1673 ft. a.s.l.
Estimated predevelopment
water level: n.a.



Perkins County


Grainton Recorder

Grainton Recorder
ID#: 404519101170301
Legal: 9-35W-23 BBB
Elevation: 3245 ft. a.s.l.
Estimated predevelopment
water level: 165 ft. b.g.s.



Grant South Recorder


Grant South Recorder
ID#: 404620101433401
Legal: 9-39W-2 DDDD
Elevation: 3413 ft. a.s.l.
Estimated predevelopment
water level: 142 ft. b.g.s.



Perkins County

Grant North Recorder
ID#: 405738101423202
Legal: 12-38W-31 CCCC
Elevation: 3423 ft. a.s.l.
Estimated predevelopment
water level: 183 ft. b.g.s.


Grant North Recorder



Pierce County

Osmond Recorder
ID#: 422150097402401
Legal: 28-3W-33 BA
Elevation: 1673 ft. a.s.l.
Estimated predevelopment
water level: 26 ft. b.g.s.


Osmond Recorder



Polk County


Shelby Recorder
ID#: 411738097264301
Legal: 15-1W-9 BBBB
Elevation: 1525 ft. a.s.l.
Estimated predevelopment
water level: n.a.

Shelby Recorder



Osceola Recorder


Osceola Recorder
ID#: 411012097325201
Legal: 14-2W-21 DB
Elevation: 1662 ft. a.s.l.
Estimated predevelopment
water level: 80 ft. b.g.s.



Red Willow County

Indianola Recorder
ID#: 401919100223001
Legal: 4-27W-17 ABB
Elevation: 2575 ft. a.s.l.
Estimated predevelopment
water level: 115 ft. b.g.s.


Indianola Recorder



Saline County

Dorchester Recorder
ID#: 403855097072501
Legal: 8-3E-19 ADA
Elevation: 1496 ft. a.s.l.
Estimated predevelopment
water level: 96 ft. b.g.s.


Dorchester Recorder



Sarpy County

Lincoln Well Field M90-28R
ID#: 410308096190701
Legal: 13-10E-32 DBBA
Elevation: 1055 ft. a.s.l.
Estimated predevelopment
water level: n.a.


Lincoln Well Field M90-28R



Saunders County

Mead Recorder
ID#: 411005096281502
Legal: 14-8E-24 ACD
Elevation: 1171 ft. a.s.l.
Estimated predevelopment
water level: 40.5 ft. b.g.s.


Mead Recorder



Scottsbluff County

Harvard Recorder
Scottsbluff Recorder
ID#: 415325103392801
Legal: 22-55W-11
Elevation: 3950 ft. a.s.l.
Estimated predevelopment
water level: n.a.


Scottsbluff Recorder



Seward County

Seward Recorder
ID#: 405406097115001
Legal: 11-2E-21 DD
Elevation: 1550 ft. a.s.l.
Estimated predevelopment
water level: 74 ft. b.g.s.


Seward Recorder



Sheridan County

Mirage Flats Recorder
ID#: 423034102415001
Legal: 29-46W-10 AA
Elevation: 3794 ft. a.s.l.
Estimated predevelopment
water level: 38.5 ft. b.g.s.


Mirage Flats Recorder



Thayer County

Carleton Recorder
ID#: 401537097434101
Legal: 3-4W-2 AA
Elevation: 1605 ft. a.s.l.
Estimated predevelopment
water level: 96 ft. b.g.s.


Carleton Recorder



Valley County

Ord Recorder


ID#: 413156098591201
Legal: 18-15W-13 ACD
Elevation: 2160 ft. a.s.l.
Estimated predevelopment
water level: 69 ft. b.g.s.



Wheeler County

Bartlett Recorder


ID#: 415445098252501
Legal: 22-10W-1
Elevation: 2070 ft. a.s.l.
Estimated predevelopment
water level: n.a.



York County


York Recorder

ID#: 405305097351503
Legal: 11-2W-31 BA
Elevation: 1659 ft. a.s.l.
Estimated predevelopment
water level: 80 ft. b.g.s.



York Pederson Recorder

York Pederson Recorder
ID#: 405305097351504
Legal: 11-2W-31 BAD
Elevation: 1659 ft. a.s.l.
Estimated predevelopment
water level: 85 ft. b.g.s.



REFERENCES

- Burbach, M., 2006, Groundwater-Level Changes in Nebraska, Spring 2000 to Spring 2006: University of Nebraska–Lincoln Conservation and Survey Division, Water Survey Map 63.
- Burbach, M., 2007, Groundwater-Level Changes in Nebraska, Spring 2000 to Spring 2007: University of Nebraska–Lincoln Conservation and Survey Division, Water Survey Map 74.
- Ellis, M.J., and Dreeszen, V.H., 1987, Groundwater levels in Nebraska, 1986: University of Nebraska–Lincoln, Conservation and Survey Division, Nebraska Water Survey Paper Number 62, 68 p.
- Johnson, M.S., and Pederson, D.T., 1984, Groundwater levels in Nebraska, 1983: University of Nebraska–Lincoln, Conservation and Survey Division, Nebraska Water Survey Paper Number 57, 67 p.
- Sophocleous, M.A., 1998, On the elusive concept of safe yield and the response of interconnected stream-aquifer systems to development, In Perspectives on Sustainable Development of Water Resources in Kansas, ed. M.A. Sophocleous, p. 6-85. Kansas Geological Survey Bulletin 239. KGS, Lawrence, KS.

Groundwater-Level Changes in Nebraska Map Series

Available on-line at <http://snr.unl.edu/data/water/groundwatermaps.asp>

Reports Containing Water-level Information

Year	Publication and number	Author(s) and year published
pre 1954	U.S.G.S. Open-File Rpt. 54-138	Keech, C.F.; Case, R.L., 1954
1954	U.S.G.S. Open-File Rpt. 55-80	Keech, C.F.; Case, R.L., 1955
1955	U.S.G.S. Open-File Rpt. 56-70	Keech, C.F., 1956
1956	U.S.G.S. Open-File Rpt. 57-61	Keech, C.F., 1957
1957	Nebraska Water Survey Paper 4*	Keech, C.F., 1958
1958	Nebraska Water Survey Paper 5*	Keech, C.F., 1959
1959	Nebraska Water Survey Paper 6	Keech, C.F., 1960
1960	Nebraska Water Survey Paper 9	Keech, C.F., 1961
1961	Nebraska Water Survey Paper 12	Keech, C.F.; Hyland, J.B., 1962
1962	Nebraska Water Survey Paper 13	Emery, P.A.; Malhoit, M.M., 1963
1963	Nebraska Water Survey Paper 14	Emery, P.A.; Malhoit, M.M., 1964
1964	Nebraska Water Survey Paper 17	Emery, P.A.; Malhoit, M.M., 1965
1965	Nebraska Water Survey Paper 18	Emery, P.A.; Malhoit, M.M., 1966
1966	Nebraska Water Survey Paper 20*	Keech, C.F., 1967
1967	Nebraska Water Survey Paper 23	Keech, C.F., 1968
1968	Nebraska Water Survey Paper 24*	Keech, C.F.; Svoboda, G.R., 1969
1969	Nebraska Water Survey Paper 26*	Keech, C.F., 1970
1970	Nebraska Water Survey Paper 28*	Keech, C.F., 1971
1971	Nebraska Water Survey Paper 33	Keech, C.F., 1972
1972	Nebraska Water Survey Paper 34	Ellis, M.J., 1973
1973	Nebraska Water Survey Paper 36	Ellis, M.J., 1974
1974	Nebraska Water Survey Paper 40*	Ellis, M.J., 1975
1975	Nebraska Water Survey Paper 43*	Ellis, M.J.; Pederson, D.T., 1976
1976	Nebraska Water Survey Paper 44	Ellis, M.J.; Pederson, D.T. 1977
1977	Nebraska Water Survey Paper 45	Ellis, M.J.; Pederson, D.T., 1978
1978	Nebraska Water Survey Paper 49	Pederson, D.T.; Johnson, M.S., 1979
1979	Nebraska Water Survey Paper 50*	Johnson, M.S.; Pederson, D.T., 1980
1980	Nebraska Water Survey Paper 51	Johnson, M.S.; Pederson, D.T., 1981
1981	Nebraska Water Survey Paper 52	Johnson, M.S.; Pederson, D.T., 1982
1982	Nebraska Water Survey Paper 56	Johnson, M.S.; Pederson, D.T., 1983
1983	Nebraska Water Survey Paper 57	Johnson, M.S.; Pederson, D.T., 1984
1984	Nebraska Water Survey Paper 59	Ellis, M.J.; Pederson, D.T., 1985
1985	Nebraska Water Survey Paper 61	Ellis, M.J.; Pederson, D.T., 1986
1986	Nebraska Water Survey Paper 62	Ellis, M.J.; Dreeszen, V.H., 1987
1987	Nebraska Water Survey Paper 65	Ellis, M.J.; Wigley, P.B., 1988
1988	Nebraska Water Survey Paper 66	Ellis, M.J.; Steele, G.V.; Wigley, P.B., 1989
1989	Nebraska Water Survey Paper 67	Ellis, M.J.; Steele, G.V.; Wigley, P.B., 1990
1990	Nebraska Water Survey Paper 69	Steele, G.V.; Wigley, P.B., 1991
1991	Nebraska Water Survey Paper 71	Steele, G.V.; Wigley, P.B., 1992
1992	Nebraska Water Survey Paper 72	Steele, G.V.; Wigley, P.B., 1994
1994	Nebraska Water Survey Paper 74	Chen, A.H.; Wigley, P.B., 1996
1996	Nebraska Water Survey Paper 75	Chen, A.H.; Khisty, M.J., 1999

*Out of print; but available for study at the address below:

Nebraska Maps and More
 101 Hardin Hall
 3310 Holdrege Street
 University of Nebraska-Lincoln
 Lincoln, Nebraska 68583-0961

Voice 402-472-3471
 Fax 402-472-2946
 Email snrsales@unl.edu
 Website snr.unl.edu



Conservation and Survey Division
School of Natural Resources
Institute of Agriculture and Natural Resources
University of Nebraska–Lincoln

ISBN-13 978-1-56161-012-9
ISBN-10 56161-012-7