

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Nebraska Swine Reports

Animal Science Department

1995

Effect of Cooking Method on Nutrient Content of Boneless Pork Loin Roasts

Judy A. Driskell

University of Nebraska-Lincoln, jdriskell1@unl.edu

Judith H. Batenhorst

University of Nebraska-Lincoln

Fayrene Hamouz

University of Nebraska-Lincoln, fhamouz1@unl.edu

Follow this and additional works at: https://digitalcommons.unl.edu/coopext_swine



Part of the [Animal Sciences Commons](#)

Driskell, Judy A.; Batenhorst, Judith H.; and Hamouz, Fayrene, "Effect of Cooking Method on Nutrient Content of Boneless Pork Loin Roasts" (1995). *Nebraska Swine Reports*. 162.

https://digitalcommons.unl.edu/coopext_swine/162

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Swine Reports by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



composition (Table 3) for the 70-pound samples. Conversely, neither grinding nor temperature nor weight improved prediction for the 40-pound samples.

Electromagnetic scanning is effective and accurate (within 2 pounds or < 3%) for prediction of fat-free lean in pork trim, presuming sample size is sufficient.

¹N. L. Meseck and B. L. Gwartney were graduate students, and C. R. Calkins is a Professor in the Animal Science Department at the University of Nebraska - Lincoln.

Table 3. Prediction of fat-free lean in pork trim.

Tub weight, lb.	Particle size	Fat-free lean weight			Fat-free lean, %		
		Model	R ²	RMSE, lb. ^a	Model	R ²	RMSE, %
70	Lean trim	Peak, temp.	.714	2.04	Peak, temp., wt.	.723	2.92
	Ground lean	Peak, temp.	.821	1.62	Peak, temp., wt.	.829	2.29
40	Lean trim	Peak	.451	1.28	Peak	.455	3.22
	Ground lean	Peak	.358	1.38	Peak	.362	3.48

^aRMSE = root mean square error.

Effect of Cooking Method on Nutrient Content of Boneless Pork Loin Roasts

Judy A. Driskell
Judith H. Batenhorst
Fayrene L. Hamouz¹

The nutritive qualities of boneless Chef's Prime™ pork loin roasts cooked by three household cooking methods to two internal temperatures were evaluated. Fresh pork loins were obtained from a vendor and prepared by UNL's Meat Laboratory according to National Pork Producers Council's specifications for the Chef's Prime™ trademarked cut with 1/8-inch fat trim. The roasts were frozen for less than two months before defrosting in the refrigerator and cooking.


The National Pork Producers Council now recommends that pork be cooked to an endpoint internal temperature of 160° F rather than the previously recommended temperature of 180° F. This is because new swine production practices have reduced concerns about trichinosis. Roasts were cooked in a household oven at 325° F to internal temperatures of 160° F (the new recommendation) and 180° F (the former recommendation). The loins were cooked by roasting, braising, and cooking in a large (Reynolds) oven bag. The loin roasts were between 2.4 and 4 pounds. The average cooking times for the roasts are given in Table 1. Pork that was braised

Table 1. Average Cooking Time

Cooking method	Internal temperature	
	160° F	180° F
	(minutes)	
Roast	131	164
Braise	107	121
Bag	109	122

or cooked in a bag reached 160° F or 180° F internal temperature much more quickly than pork that was roasted. The cooked pork contained a mean of 68% moisture and 8% crude fat.

Pork cuts are "good" to "major" sources of many nutrients that Americans frequently consume in less than adequate quantities. These include



Values represent least squares means and standard errors. Values for each nutrient not sharing a common superscript are significantly different at P < .01.

Figure 1. True Retention Values for Three Vitamins in Pork Roasts Prepared by Three Cooking Methods to Two Internal Temperatures



vitamin B-6, vitamin E, iron, magnesium, zinc, and selenium. The amounts of these nutrients in the cooked pork roasts and their retention values were determined. Thiamin was used as the index nutrient. Cooked pork roasts (3.5 ounces) were found to contain approximately 20% of the vitamin B-6, 49% of the thiamin, 2% of the vitamin E, 10% of the iron, 6% of the magnesium, 20% of the zinc, and 89% of the selenium needed to meet the Recommended Dietary Allowances of adults for a day.

True retention is a term that relates the percentage of nutrient content of the food as cooked to the content before cooking. The true retention of the vitamins in the pork roasts prepared by the three cooking methods to the two internal temperatures are given in Figure 1. Retention values for vitamin B-6 and thiamin were significantly higher ($P < .01$) in pork cooked to 160° F than to 180° F. Vitamin B-6 retention values for pork cooked in a bag were significantly higher ($P < .01$) than for pork that was

roasted, whereas pork that was roasted had significantly higher ($P < .01$) values than pork that was braised. Thiamin retention values were significantly higher ($P < .01$) in pork that was cooked in a bag or roasted than in pork that was braised. Mean true retention values were 58% for vitamin B-6 and 51% for thiamin. Hence, almost half of the vitamin B-6 and thiamin were destroyed during cooking. The highest true retention values for these two vitamins were for pork cooked in a bag.

The vitamin E retention in pork prepared by the different cooking: temperature methods was similar. The pork roasts contained a small amount of vitamin E, only enough to meet about 2% of recommended intakes per serving. However, the mean true retention value for vitamin E was only 44%, indicating that over half of the vitamin E was destroyed during cooking. This was independent of the cooking: temperature method used. The lower fat trim of today's pork cuts may result in the lower vitamin E content.

True retention values for iron, magnesium, zinc, and selenium were similar for the different cooking: temperature methods and were close to 100%. Hence, no loss of minerals occurred while the pork was being cooked.

True retention values for vitamin B-6, thiamin, and vitamin E were highest for pork roasts cooked in the bag to an internal temperature of 160° F. However, true retention values for iron, magnesium, zinc, and selenium were similar in pork cooked in the bag, braised, or roasted to either 160° or 180° F internal temperature. Chef's Prime™ loin roasts were found to be "major" sources of vitamin B-6, thiamin, zinc, and selenium and a "good" source of iron.

¹Judy A. Driskell is a Professor, Judith H. Batenhorst, a graduate student, and Fayrene L. Hamouz, an Assistant Professor, in the Department of Nutritional Science and Dietetics., University of Nebraska, Lincoln.

Utilization of Twin Screw Cold Extrusion to Manufacture Restructured Chops from Lower-Valued Pork

Wesley N. Osburn
Roger W. Mandigo
Paul S. Kuber¹

Restructured meat products are commonly manufactured by using lower-valued meat trimmings reduced in size by comminution (flaking, chunking, grinding, chopping or slicing). The comminuted meat mixture is mixed with salt and water to extract salt-soluble proteins. These extracted proteins are critical to produce a "glue" which binds muscle pieces together. These muscle pieces may then be reformed to produce a "meat log" of specific form or shape. The log is then cut into steaks or chops which, when

cooked, are similar in appearance and texture to their intact muscle counterparts.

Two concerns must be addressed in the manufacture of restructured meat products: texture, and the removal and degradation of connective tissue. Lower-valued meat trimmings used in restructuring tend to contain more connective tissue which may affect product texture.

Mechanical desinewing is used to remove connective tissue from boneless meat trimmings. Reducing the connective tissue in trimmings increases their value for use in various restructured meat products. The method of comminution also affects the final prod-

uct texture, which usually is somewhere between that of ground (hamburger) and an intact muscle (steak or chop) meat product.

Recently, twin screw cold extrusion has been used as a processing technology to produce restructured meat products. In this process, a comminuted meat mixture is forced to flow through an enclosed twin-screw extruding horn to form "extruded ropes" of a specific shape and size. These ropes can be pressed together to form meat logs which then can be cleaved into restructured steaks or chops. This process is believed to partially realign muscle fibers and modify the texture of meat products.

(Continued on next page)