

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Bird Control Seminars Proceedings

Wildlife Damage Management, Internet Center
for

September 1968

DRC-1339 IN FEEDLOTS

John W. DeGrazio

U.S. Bureau of Sport Fisheries and Wildlife, Denver, CO

Follow this and additional works at: <https://digitalcommons.unl.edu/icwdmbirdcontrol>



Part of the [Environmental Sciences Commons](#)

DeGrazio, John W., "DRC-1339 IN FEEDLOTS" (1968). *Bird Control Seminars Proceedings*. 167.
<https://digitalcommons.unl.edu/icwdmbirdcontrol/167>

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Bird Control Seminars Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

DRC-1339 IN FEEDLOTS

John W. De Grazio
U.S. Bureau of Sport Fisheries and Wildlife
Federal Center, Denver, Colorado

Since 1960, the Denver Wildlife Research Center has been investigating methods of controlling starlings (*Sturnus vulgaris*) at livestock feedlots. One chemical developed at the Denver Center is a slow-acting avicide, DRC-1339 (3-chloro-*p*-toluidine); it is well accepted and highly toxic to starlings, generally less toxic to other birds, and relatively nontoxic to mammals. It possesses no secondary hazard to avian and mammalian predators. A Cooper's hawk (*Accipiter cooperii*), a marsh hawk (*Circus cyaneus*), and a sparrow hawk (*Falco sparverius*) subsisted on field-killed DRC-1339 starlings for 3- and 4-month periods with no ill effects.

In our tests, poultry pellets (Purina Layena Checkers') have been the most effective and selective starling bait. Field tests in many states have shown the utility of this compound when applied to pellets. Pellets were treated to contain approximately 1 percent DRC-1339, diluted with untreated pellets and grains, and broadcast in pens and alleys at cattle feedlots and on the perimeters of pens at turkey feedlots. Baiting rates varied from 2 to 10 lbs. of treated pellets per acre, depending on the size of the lots; dilution rates were generally 1:10.

DRC-1339 has proved to be an extremely effective and safe toxicant for starlings at animal feedlots. It is now marketed commercially under the trade name of Starlicide.

¹ Reference to trade names does not imply endorsement of commercial products by the Federal Government.