

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

March 1995

Two species of Cerylonidae new to Florida and the U.S. (Coleoptera)

Michael C. Thomas

*Florida State Collection of Arthropods, Florida Department of Agriculture and Consumer Services,
Gainesville, FL*

Paul E. Skelley

*Florida State Collection of Arthropods, Florida Department of Agriculture and Consumer Services,
Gainesville, FL*

Randall W. Lundgren

*Florida State Collection of Arthropods, Florida Department of Agriculture and Consumer Services,
Gainesville, FL*

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Entomology Commons](#)

Thomas, Michael C.; Skelley, Paul E.; and Lundgren, Randall W., "Two species of Cerylonidae new to Florida and the U.S. (Coleoptera)" (1995). *Insecta Mundi*. 171.

<https://digitalcommons.unl.edu/insectamundi/171>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Two species of Cerylonidae new to Florida and the U.S. (Coleoptera)

Michael C. Thomas, Paul E. Skelley, and Randall W. Lundgren

Florida State Collection of Arthropods, Division of Plant Industry,
Florida Department of Agriculture and Consumer Services,
P.O. Box 147100, Gainesville, FL 32614-7100, U.S.A.

In their paper on the Cerylonidae of America north of Mexico, Lawrence and Stephan (1975) noted that *Euxestus erithacus* Chevrolat is widespread in the Antilles and, although not recorded from Florida, might be found in the southern part of the state. We here report the accuracy of their prediction.

While recording specimen data for a checklist of Florida Coleoptera, we noticed that some specimens of what we assumed to be *Hypodacne punctata* LeConte, a widespread eastern North American species, appeared slightly different than others. Closer examination revealed that these specimens possessed the distinguishing features of *Euxestus* Wollaston as mentioned by Lawrence and Stephan (1975).

Euxestus keys to *Hypodacne* LeConte in Lawrence and Stephan (1975). Their key is modified as follows to include *Euxestus*:

2. Procoxal cavities closed behind; length more than 1.5 mm.; body oval and highly convex, with smooth, even, lateral edges and no antennal cavities 2a
 - Procoxal cavities open behind; length less than 1.5 mm.; without other characters in combination 3
- 2a. Tarsi not lobed; penultimate antennomere asymmetrical, almost as wide as ultimate; length 2.2 - 2.6 mm.; color usually castaneous *Hypodacne* LeConte
 - Tarsomere I lobed; penultimate antennomere symmetrical, much narrower than ultimate; length 1.8 - 2.0 mm.; color usually piceous *Euxestus* Wollaston

We have seen seven Florida specimens of *Euxestus erithacus* from the following localities (specimens are deposited in the Florida State Collection of Arthropods (FSCA) and in the collections of the second and third authors (PESC and RWLC, respectively)): **Alachua Co.:** 29°34.5' N 82°29' W 29-XI-1994 flight barrier trap in hardwood hammock (RWLC); **Dade Co.:** Everglades National Park, Compartment M, 23-III-1994, ethanol/turpentine trap (FSCA); same, except Compart-

ment J (FSCA); Homestead, 14-IX-1993 (FSCA); Miami, Deering Estate, 27-XI-1993, ethanol/turpentine trap (FSCA); Camp Mahachie, nr. Matheson Hammock, V-1991, blacklight trap (FSCA).

We have also examined two Florida specimens of *Hypodacne punctata* from: **Liberty Co.:** Torreya State Park, 17-VII-1987, berlese basal tree-hole litter; same data except 26-27-III-1988, under bark (PESC).

Lawrence and Stephan (1975) described *Ostomopsis neotropicalis* from specimens collected in Mexico and Panama. They reported that they could not locate the Florida specimen of this genus that was mentioned by Sen Gupta and Crowson (1973). We can confirm the presence of this genus in Florida based on the collection of a single specimen of *O. neotropicalis* with the following data: **Dade Co.** Camp Mahachie, nr. Matheson Hammock, 18-I-1991, J. Gleason, blacklight trap (FSCA).

Acknowledgements

We thank W. N. Dixon, G. J. Steck, and W. C. Welbourn, Division of Plant Industry, for critically reading this manuscript; J.R. Meeker, Division of Forestry, for supplying samples from the post-Hurricane Andrew scolytid survey; and John Gleason and the Girl Scout Council of Tropical Florida for facilitating the operation of a blacklight trap at Camp Mahachie. This is Entomology Contribution No. 819, Bureau of Entomology, Nematology, and Plant Pathology, Division of Plant Industry, Florida Department of Agriculture and Consumer Services.

References

- Lawrence, J. F., and K. Stephan. 1975. The North American Cerylonidae (Coleoptera: Clavicornia). *Psyche* 82(2): 131-166.
- Sen Gupta, T., and R. A. Crowson. 1973. A review of the classification of Cerylonidae (Coleoptera, Clavicornia). *Trans. Roy. Ent. Soc. London* 124: 365-446.