

2009

ANHYDROUS FLUORIDE SALTS AND REAGENTS AND METHODS FOR THEIR PRODUCTION

Stephen G. DiMagno
Lincoln, NE, sdimagno@uic.edu

Haoran Sun
Lincoln, NE

Follow this and additional works at: <https://digitalcommons.unl.edu/chemfacpub>

 Part of the [Analytical Chemistry Commons](#), [Medicinal-Pharmaceutical Chemistry Commons](#), and the [Other Chemistry Commons](#)

DiMagno, Stephen G. and Sun, Haoran, "ANHYDROUS FLUORIDE SALTS AND REAGENTS AND METHODS FOR THEIR PRODUCTION" (2009). *Faculty Publications -- Chemistry Department*. 185.
<https://digitalcommons.unl.edu/chemfacpub/185>

This Article is brought to you for free and open access by the Published Research - Department of Chemistry at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications -- Chemistry Department by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



US007592486B2

(12) **United States Patent**
DiMagno et al.

(10) **Patent No.:** **US 7,592,486 B2**
(45) **Date of Patent:** **Sep. 22, 2009**

(54) **ANHYDROUS FLUORIDE SALTS AND REAGENTS AND METHODS FOR THEIR PRODUCTION**

6,794,401 B2 9/2004 Nag

FOREIGN PATENT DOCUMENTS

CA 2035561 11/1991

(75) Inventors: **Stephen G. DiMagno**, Lincoln, NE (US); **Haoran Sun**, Lincoln, NE (US)

OTHER PUBLICATIONS

(73) Assignee: **Board of Regents of the University of Nebraska**, Lincoln, NE (US)

ChemBioChem Special Issue: Fluorine in the Life Sciences 2004, 5, 557726.

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 312 days.

Christe, K.O., et al., J. Am. Chem. Soc. 1990, 112, 7619-25.

Gnann, R.Z., et al., J. Am. Chem. Soc. 1997, 119, 112-115.

Kornath, A., et al. Inorg. Chem. 2003, 42, 2894-2901.

JACS, 1990, 112, 7619-7625.

J. Am. Chem. Soc., Bennett, B.K. et al., 1994, 116, 11165-11166.

JACS, 1997, 119, 112-115.

Inorg. Chem., 2003, 42, 2894-2901.

Cox, D.P., et al., J. Org. Chem. 1984, 49, 3216-19.

Shann, R.K., et al., J. Org. Chem. 1983, 48, 2112-14.

Pilcher, et al. J. Am. Chem. Soc. 1995, 117, 5166-5167.

Akiyama, Y, et al., S. Synlett 2003, 1530-1532.

(21) Appl. No.: **11/226,277**

(22) Filed: **Sep. 15, 2005**

(65) **Prior Publication Data**

US 2006/0089514 A1 Apr. 27, 2006

Related U.S. Application Data

(60) Provisional application No. 60/610,412, filed on Sep. 16, 2004.

(51) **Int. Cl.**
C07C 211/63 (2006.01)

(52) **U.S. Cl.** **564/281**; 568/9

(58) **Field of Classification Search** 564/281;
568/9

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,369,212 A 11/1994 Christe

5,854,084 A 12/1998 Drukier

6,156,812 A 12/2000 Lau

6,451,921 B2 9/2002 Weisse

(57) **ABSTRACT**

Anhydrous organic fluoride salts and reagents prepared by a method comprising the nucleophilic substitution of a fluorinated aromatic or fluorinated unsaturated organic compound with a salt having the formula:



in an inert polar, aprotic solvent; wherein M is an atom capable of supporting a formal positive charge, the n groups Q are independently varied organic moieties, n is an integer such that the $[Q_nM]$ carries at least one formal positive charge, x is an integer defining the number of formal positive charge(s), +, carried by the $[Q_nM]$, A^- is an anionic nucleophile capable of substituting for F in the fluorinated compound and F represents fluorine or a radioisotope thereof.

16 Claims, 10 Drawing Sheets




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10

ANHYDROUS FLUORIDE SALTS AND REAGENTS AND METHODS FOR THEIR PRODUCTION

This application claims benefit of 60/610,412, filed Sep. 16, 2004.

FIELD OF THE INVENTION

The invention relates to novel organic fluorides and methods for their production.

BACKGROUND OF THE INVENTION

Fluorine substitution is a powerful tool to improve the bioavailability of pharmaceuticals and agrochemicals. Thus, an expansive set of nucleophilic and electrophilic reagents have been developed to replace various C—X functional groups with C—F.

[*CheMBioChem Special Issue: Fluorine In the Life Sciences* 2004, 5, 557726]. Simplest among the nucleophilic fluorinating reagents are “anhydrous” or “naked” organic fluoride salts, represented by tetramethylammonium fluoride (TMAF) [Christe, K. O, et al, *J. Am. Chem. Soc.* 1990, 112, 7619-25, 1-methylhexamethylenetetramine fluoride (MHAF) [Gnann, R. Z., et al, *J. Am. Chem. Soc.* 1997, 119, 112-115] and tetramethylphosphonium fluoride (TMPF) [Kornath, A, et al, *Inorg. Chem.* 2003, 42, 2894-2901]. Highly soluble anhydrous fluoride salts possessing a wide variety of alkyl groups are desirable for synthetic purposes, but these compounds cannot be prepared according to current methodologies.

Typical of prior art methods for preparing such salts are those described in U.S. Pat. No. 5,369,212 and Canadian patent no. 2035561.

The preparation of absolutely anhydrous fluoride salts whose cations are substituted with alkyl groups possessing beta-hydrogen atoms has proved to be a significant challenge. Approximately 20 years ago, the first claims for “anhydrous” tetrabutylammonium fluoride appeared. The compounds were prepared by physical drying of the hydrated salt, i.e., dynamic high vacuum (<0.1 mmHg) to remove water for at least 48 hours from TBAF·3H₂O at 40~45° C. (JOC, 1984, 49, 3216-3219). However, there was still 0.1 to 0.3 equiv of water in this “anhydrous” TBAF and copious quantities of the elimination products (tributylamine, bifluoride ion, and butane) as a result of this process. The side reactions and the presence of water and tributylamine significantly decrease the reactivity of the fluoride ion and lead to significant side reactions, such as hydrolysis of the starting substrates. An example of water’s deleterious effects upon the reactivity of TBAF can be seen in simple model reactions. For example, if TBAF that is dried using physical methods is combined with benzyl chloride or benzyl bromide at room temperature to 40° C., formation of benzyl fluoride required 8 to 12 hours. In comparison, if truly anhydrous TBAF were employed, the same reaction would only take a few minutes or less at low temperatures and give quantitative yields.

Later, individual syntheses of tetramethylammonium fluoride (TMAF) (JACS, 1990, 112, 7619-7625), cobaltocenium fluoride (Cp₂CoF), (JACS, 1994, 116, 11165-11166), 1-Methylhexamethylenetetramine fluoride (MHAF) (JACS, 1997, 119, 112-115), tetramethylphosphonium fluoride (TMPF) (Inorg. Chem., 2003, 42, 2894-2901) as well as several others were synthesized and characterized as “naked” or “anhydrous” fluoride salts. However, each of these salts has specific drawbacks in terms of solubility or reactivity, and the pre-

parative methods for synthesizing these individual salts are not applicable for the preparation of a wide variety of fluoride salts. Anhydrous fluoride salts with alkyl groups capable of beta-elimination (ethyl, propyl, butyl, isopropyl, pentyl, isobutyl, etc.) in particular are not accessible by current methods.


Generally, then, these compounds are commonly prepared in a hydrated state and are subsequently dried by heating under dynamic vacuum or by azeotropic distillation. However, the conditions used to dry these salts are often incompatible with a variety of desirable cations. For example, dried tetrabutylammonium fluoride, (TBAF) [Cox, D. P., et al, *J. Org. Chem.* 1984, 49, 3216-19] is reported to decompose by Hofmann elimination at room temperature. The salt isolated after dehydration is contaminated with copious amounts of bifluoride ion (HF₂) and tributylamine [Shannn, R. K., et al, *J. Org. Chem.* 1983, 48, 2112-14]. These considerations and findings have led to the belief among those skilled in the art that “it is very unlikely that pure, anhydrous tetraalkylammonium fluoride salts have ever, in fact, been produced in the case of ammonium ions susceptible to E2 eliminations” [Sharma et al, supra].

It is an object of the present invention to provide a novel method of producing truly anhydrous organic fluoride salts and reagents.


It is a further object of the invention to provide novel anhydrous organic fluoride salts and reagents.

SUMMARY OF THE INVENTION

The above and other objects are realized by the present invention one embodiment of which relates to a method of synthesizing an anhydrous fluoride salt having the formula:



comprising the nucleophilic substitution of a fluorinated aromatic or fluorinated unsaturated organic compound with a salt having the formula:



in an inert polar, aprotic solvent; wherein M is an atom capable of supporting a formal positive charge, the n groups Q are independently varied organic moieties, n is an integer such that the [Q_nM] carries at least one formal positive charge, x is an integer defining the number of formal positive charge(s), +, carried by the [Q_nM], A⁻ is an anionic nucleophile capable of substituting for F in the fluorinated compound and F represents fluorine or a radioisotope thereof.

Another embodiment of the invention concerns anhydrous organic fluoride salts and reagents of the above formula, preferably produced by the above-described invention.

Still other embodiments of the invention relate to the use of the anhydrous organic fluoride salts and reagents of the invention in methods, processes and syntheses wherein the non-anhydrous salts and reagents are employed.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1-10 depict ¹⁹F NMR and ¹H NMR spectra for various reaction products produced by the reactions described herein (vide infra).

DETAILED DESCRIPTION OF THE INVENTION

The present invention is predicated on the discovery that a wide variety of truly anhydrous organic fluoride salts and reagents may be synthesized in one step by the nucleophilic substitution of various fluorinated organic compounds with organic salts of diffusely charged anionic nucleophiles capable of forming strong bonds to carbon in a nucleophilic substitution reaction. Thus, employing the methods of the invention a wide range of novel anhydrous salts can be prepared using one simple procedure. Moreover, as the examples set forth below demonstrate, the method of the invention allows many sensitive or unstable fluoride salts to be prepared easily. Such compounds would decompose rapidly under the conditions employed in typical literature preparations of similar compounds.

Although the invention is principally exemplified and illustrated herein for preparing tetrabutylammonium fluoride, it will be understood by those skilled in the art that the inventive method may be utilized to prepare many and varied anhydrous fluoride salts. It will also be understood by those skilled in the art that the method of the invention may also be utilized to prepare radioisotopic fluoride salts (e.g., ^{18}F).


The reaction may be carried out at low temperatures [-35°C . to RT] in polar aprotic solvents such as tetrahydrofuran, dimethyl sulfoxide, diethyl ether, dioxane, dimethoxyethane, methyl tert-butyl ether, acetonitrile, acetone, methylethylketone, tetrahydrofuran, dimethylformamide, dimethylacetamide, N-methylpyrrolidinone, butyronitrile, or in aromatic solvents such as toluene, pyridine, benzonitrile, or diphenyl ether. Other suitable solvents include carbonates such as diethyl carbonate and hexamethylphosphoric triamide. In preferred embodiments tetrahydrofuran, dimethylsulfoxide, and acetonitrile are the solvents employed. Halogenated solvents such as methylene chloride or dichloroethane are decomposed rapidly by anhydrous fluoride salts, and are thus generally not useful for this synthetic procedure.

In the above-described formulae, Q is an organic moiety capable of undergoing E2 elimination and may be, e.g., alkyl, alkenyl, alkynyl, or form the backbone or sidechain of a polymer. M may be N, P or any element capable of supporting a formal positive charge. The anion A may any diffusely charged anionic nucleophile capable of forming strong bonds to carbon in a nucleophilic substitution reaction, such as, e.g., cyanide, isothiocyanate, thiocyanate, alkyl- and arylthiolates, or azide. In preferred embodiments cyanide is the nucleophile employed.

$[\text{Q}_n\text{M}]^{x+}\text{A}_x^-$ is preferably a tetraalkylammonium cyanide, a trialkylarylammonium cyanide, a dialkyl diarylammonium cyanide, an alkyl triarylammonium cyanide, or a tetraarylammonium cyanide; Q being an organic moiety capable of undergoing E2 elimination.

The fluorinated compound nucleophilically substituted in the method of the invention is preferably a fluorinated benzene, alkene or alkyne with a large number of fluorine atoms per unit weight e.g., hexafluorobenzene, octafluoronaphthalene, octafluorotoluene, pentafluorobenzonitrile, pentafluoropyridine, decafluorobiphenyl, etc. For the generation of isotopically labeled anhydrous fluoride salts (i.e., TBA ^{18}F) a singly fluorinated arene is sufficient, e.g., 4-fluorobenzonitrile.

The reaction scheme for the method of the invention is:



Nucleophile substitution reactions are generally well known in the art as exemplified in U.S. Pat. Nos. 6,794,401; 6,451,921; 6,156,812 and 5,854,084, inter alia.






Thus tetrabutylammonium fluoride (TBAF) is easily prepared in one step at low temperatures by the nucleophilic substitution of the hexafluorobenzene with tetrabutylammonium cyanide. Adventitious water is readily scavenged by the hexacyanobenzene by-product of the reaction.

The constraints on a fluoride-generating synthesis grounded in nucleophile substitution reactions are quite severe and dictate a careful choice of the nucleophile. Because the enthalpic driving force for fluoride liberating reaction derives almost exclusively from ion-pairing and ΔBDE terms, and because the $\text{C}_{sp^2}\text{-F}$ bond in aromatics (as well as unsaturated compounds) is exceptionally strong (126 kcal/mol), only diffusely charged anionic nucleophiles capable of forming strong bonds to carbon are capable of acting in nucleophile substitution reactions at low temperature in polar aprotic solvents. Cyanide ion, a potent, weakly basic nucleophile that forms strong bonds to sp^2 -hybridized carbon (BDE=133 kcal/mol) is an excellent candidate. It will be understood by those skilled in the art, however, that any similar diffusely charged anionic nucleophile may also be employed in the method of the invention, such as, e.g., isothiocyanate, isocyanate, cyanate, thiocyanate, alkyl- and arylthiolates, or azide.

As illustrated in the examples below, treatment of hexafluorobenzene with tetrabutylammonium cyanide (TBACN) (in 1:1 to 1:6 molar ratios) in the polar aprotic solvents THF, acetonitrile, or DMSO at or below room temperature gives excellent yields of anhydrous TBAF. ^{19}F NMR spectroscopy indicates that the overall yield of TBAF in solution in all cases is $>95\%$. Cyano substitution dramatically increases the fluorinated benzene ring's susceptibility to further nucleophilic attack, as is evidenced by the observation of pentacyanofluorobenzene and hexafluorobenzene as the principal fluorinated aromatic species in the reaction solution, even if 1:1 TBACN: C_6F_6 stoichiometry is employed.

In THF, colorless to light yellow anhydrous TBAF precipitated from cooled (-35°C .) solutions and yields of the isolated salt ranged from 40% to 70%. Freshly isolated TBAF

TABLE 2

Fluorination of various substrates using anhydrous TBAF								
Run	Substrate	Reagent	Solvent	Conditions	Product	Yield* (%)	Comments	Ref.
1	PhCH ₂ Br	1.3~1.5 eq. TBAF	CD ₃ CN	-35° C., <5 min	PhCH ₂ F	100		This work ³
2	PhCH ₂ Br	2 eq. TBAF "anhydrous"	THF	RT, 8 h	PhCH ₂ F	>90	PhCH ₂ OH (5%)	
3	CH ₃ I	1.5 eq. TBAF	CD ₃ CN	-40° C., <5 min	CH ₃ F	100		This work
4	CH ₃ I	CoCp ₂ F	THF	RT, 6 h	CH ₃ F	100		This work
5	CH ₃ (CH ₂) ₂ Br	TBAF	THF	RT, <5 min	CH ₃ (CH ₂) ₂ F	40-50	(remainder alkene)	This work
6	CH ₃ (CH ₂) ₂ Br	6 eq. TBAT	CD ₃ CN	Reflux, 24 h	CH ₃ (CH ₂) ₂ F	85		This work
7	CH ₃ (CH ₂) ₂ Br	2 eq. TBAF "anhydrous"	THF	RT, 1 h	CH ₃ (CH ₂) ₂ F	48	40% octanol	This work ³
8		TBAF	THF	RT, <5 min	CH ₃ (CH ₂) ₁₃ F	100		This work
9		4 eq. TBAF	THF, or CD ₃ CN	RT, <5 min		>90		This work
10		1.3 eq. TBAF	CD ₃ CN	RT, <2 min		>95		This work
11	PhCOCl	1 eq. TBAF	THF	RT, <2 min	PhCOF	100		This work
12	Tosyl-Cl	1 eq. TBAF	THF	RT, <2 min	Tosyl-F	100		This work

*yields were calculated by integration of starting material and product signals in the ¹H and/or ¹³F NMR spectra.

Taken together, the results presented here show that exceptionally nucleophilic, highly soluble fluoride ion sources featuring ammonium cations can be prepared readily even if the cations are thought susceptible to E2 elimination. The self dehydrating nature of the nucleophilic aromatic substitution method makes it an exceptionally forgiving synthetic route to anhydrous fluoride salts.

Generally, the method of the invention produces anhydrous organic fluoride salts and reagents containing less than 0.01% H₂O in one efficient step, in high yields and low temperatures without deleterious effects on the product or reaction mechanism. Moreover, the reactivity of the anhydrous fluoride salts of the invention in solution or liquid approaches that of conventional fluoride salts in the gas phase, e.g., a reaction between nitrobenzene and conventionally produced organic fluoride salt in solution will not proceed but will in gas phase. The anhydrous fluoride salts of the invention will react with nitrobenzene in solution.

EXAMPLES

All reagents were handled under N₂. Hexafluorobenzene (C₆F₆) (99%, SynQuest) was passed through a column of activated (130° C. for 5 h) silica gel and distilled from CaH₂.

Acetonitrile (HPLC grade, Aldrich) was distilled from P₂O₅ and redistilled under reduced pressure from CaH₂. THF (anhydrous, Aldrich) was distilled from LiAlH₄. Purified solvents were stored under N₂ in Schlenk-style flasks under N₂. Tetra-n-butylammonium cyanide (TBACN) (97%) was obtained from Fluka Chemical Co. TBACN was dried under vacuum at 40° C. overnight prior to use. For initial work, TBACN was recrystallized from THF/Hexane by layering, subsequent studies showed that this purification step was unnecessary. Tetramethylammonium hexafluorophosphate (TMAPP₆) was obtained from Fluka and dried under vacuum. All other reagents were of analytical grade, from Aldrich. All chemical handling was performed under N₂ in a glove box.

¹H, ¹³C and ¹⁹F NMR spectra were determined in the Instrumentation Center at the University of Nebraska-Lincoln. 400 MHz (QNP probe for ¹H, ¹³C and ¹⁹F NMR spectra), 500 MHz (QNP probe for ¹H, ¹³C and ¹⁹F NMR spectra) and 600 MHz (HF probe for ¹H and ¹⁹F NMR spectra) NMR spectrometers were used in this study. ¹⁹F NMR chemical shifts were referenced to an internal standard, hexafluorobenzene

Anhydrous tetrabutylammonium fluoride (TBAF):

0.67 g TBACN was dissolved in 2.5 ml THF and the resulting solution was cooled to -65°C . A chilled solution (-65°C .) of 0.3 ml hexafluorobenzene (C_6F_6) in 0.5 ml THF was added, and the mixture was allowed to warm gradually (over 4 hours) to -15°C . During this time the solution changed from colorless to yellow-green, and a white solid precipitated. The mixture was again cooled to -65°C ., the solid was filtered and washed two times with cold THF. All isolation procedures were kept below -36°C . The white or light yellow TBAF solid was collected and put into a -36°C . freezer for short term storage. Total TBAF yield was over 95% (based upon TBACN, confirmed by quenching experiments with benzyl chloride) if the mixture was used directly. Isolated yields of the solid material varied from 40% to 70% depending on the rapidity of the wash and filtration steps. ^1H NMR ($(\text{CD}_3)_2\text{SO}$) 3.23 (8H, m), 1.56 (8H, m), 1.28 (8H, sext, $J=7.31$ Hz), 0.86 (12H, t, $J=7.31$ Hz); ^{19}F NMR ($(\text{CD}_3)_2\text{SO}$) -72.6 ppm (s); ^{13}C NMR ($(\text{CD}_3)_2\text{SO}$): 57.5, 23.1, 19.2, 13.5 ppm.

Generation of TBAF in CH_3CN :

TBACN (0.134 g, 0.5 mmol) was dissolved in anhydrous acetonitrile (0.5 ml). At 25°C ., 9.6 μl (0.083 mmol) C_6F_6 was added, and the initially colorless solution changed to dark-red immediately. The reaction was monitored by ^{19}F NMR spectroscopy. Fluoride generation was complete within 1 h. A representative ^{19}F NMR spectrum is shown in FIG. 1.

Generation of TBAF in DMSO:

A very similar procedure was used to generate TBAF in DMSO. TBACN (0.134 g, 0.5 mmol) was dissolved in anhydrous acetonitrile (0.5 ml). At 25°C ., 9.6 μl (0.083 mmol) C_6F_6 was added, and the mixture was allowed to stand for one

h. The solubility of TBAF in both CH_3CN and DMSO was excellent (up to 2 M). The solution was directly used in the fluorination reaction.

TMACN—TMACN was prepared by metathetical ion-exchange of TBACN with TMAPF_6 in acetonitrile/THF. 110 mg (0.5 mmol) TMAPF_6 was dissolved in a minimum amount of acetonitrile, and a saturated acetonitrile solution of TBACN (134 mg, 0.5 mmol) was added. The precipitated TMACN was filtered, washed with a small amount of acetonitrile, and the residual solvents were evaporated. ^1H NMR (CD_3CN) 3.11 ppm (s), ^{13}C NMR (CD_3CN) 54.27, 167.20.

TMAF—TMAF was synthesized from TMACN and C_6F_6 in acetonitrile by a method similar to that described for TBAF. 4.6 mg TMACN dissolve in 0.6 ml of $(\text{CD}_3)_2\text{SO}$ at room temperature. 1.0 μl hexafluorobenzene (C_6F_6) was and the mixture was allowed to stand at room temperature for 12 h.

General Procedure for Fluorination Reactions

The general procedure given below was used for all fluorination reactions employing in situ generated TBAF. Yields were calculated by integration of the relevant peaks in the ^1H and ^{19}F NMR spectra.









In an NMR tube equipped with a PTFE resealable closure, TBACN (0.134 g, 0.5 mmol) was dissolved in anhydrous CD_3CN (or $(\text{CD}_3)_2\text{SO}$) (0.5 ml). At 25°C ., 9.6 μl (0.083 mmol) C_6F_6 was added, and the mixture was held at room temperature for 1 h. The mixture was cooled to -40°C . and the substrate (0.25-0.5 mmol) was added. The solution was mixed vigorously and the tube was transferred to a precooled (-35°C .) NMR probe and spectra were gathered. The time elapsed from the sample mixing until completion of the first NMR spectrum was approximately 3 min. The reaction was monitored by ^{19}F NMR spectra every 2 minutes until no further change was observed.

Table 3 shows results of fluorination of various substrates under different conditions. For comparison, the literature results by other fluorination reagents are listed in table 3.

TABLE 3

Run	Substrate	Reagent	Solvent	Temp and Time	Product	Yield (%)	Comments	Ref.
1	PhCH_2Br	1.3~1.5 eq. TBAF	acetonitrile	-35°C ., <5 min	PhCH_2F	100	No PhCH_2OH	This work
2	PhCH_2Br		DMSO	RT, <2 min	PhCH_2F	100		This work
3	PhCH_2Br		THF	RT, <2 min	PhCH_2F	100		This work
4	PhCH_2Br	2 eq. TBAF "anhydrous"	THF	RT, 8 hours	PhCH_2F	>90	PhCH_2OH (5%)	1
5	PhCH_2Cl	1.5 eq. TBAF	THF	RT, <2 min	PhCH_2F	100		This work
6	PhCH_2Cl	2 eq. TBAF "anhydrous"	THF	40°C ., 12 hours	PhCH_2F			1
7	PhCH_2Cl	CoCp_2F	THF	RT, 90 min	PhCH_2F	95		2
8	CH_3I	1.5 eq. TBAF	acetonitrile	-40°C ., <5 min	CH_3F	100		This work
9	CH_3I	CoCp_2F	THF	RT, 6 hours	CH_3F	100		2
10	$\text{CH}_3(\text{CH}_2)_7\text{Br}$	TBAF	THF	RT, <5 min	$\text{CH}_3(\text{CH}_2)_7\text{F}$	40~50	No octanol	This work
11	$\text{CH}_3(\text{CH}_2)_7\text{Br}$	6 eq. TBAT	acetonitrile	Reflux, 24 h	$\text{CH}_3(\text{CH}_2)_7\text{F}$	85		3
12	$\text{CH}_3(\text{CH}_2)_7\text{Br}$	2 eq. TBAF "anhydrous"	THF	RT, 1 hour	$\text{CH}_3(\text{CH}_2)_7\text{F}$	48	40% octanol	1
13	$\text{CH}_3(\text{CH}_2)_{17}(\text{p-Cl-benzenesulfonate})$	TBAF	THF	RT, <5 min	$\text{CH}_3(\text{CH}_2)_{17}\text{F}$	100		This work
14	$\text{CH}_3(\text{CH}_2)_7\text{OTs}$	2 eq. TBAF "anhydrous"	none	RT, 1 hour	$\text{CH}_3(\text{CH}_2)_7\text{F}$	98	2% alkene	1

TABLE 3-continued

Run	Substrate	Reagent	Solvent	Temp and Time	Product	Yield (%)	Comments	Ref.
15	CH ₃ (CH ₂) ₇ OTs	4 eq. TBAF	acetonitrile	Reflux, 24 h	CH ₃ (CH ₂) ₇ F	99	Trace alkene	³
16		2 eq. TBAF	THF, or acetonitrile	RT, <5 min		100		This work
17		1.3 eq TBAF	DMSO, or acetonitrile	RT, <8 hours		>90		This work
18		1.3 eq TBAF	acetonitrile	RT, <2 min		~95		This work
19	PhCOCl	1 eq. TBAF	THF	RT or below RT, <2 min	PhCOF	100		This work
20	PhCOCl	2 eq. TBAF "anhydrous"		RT, 1 hour	PhCOF	81*		¹
21	Tosyl-Cl	1 eq. TBAF	THF	RT, <2 min	Tosyl-F	100		This work
22		2.5 eq. TBAF, 3 h; followed by add H ₂ O	DMSO or acetonitrile	RT, ~3 hours				This work
23		TBAF-KHF ₂	none	120° C., 2 hours		86	Contains 10% PhCHFCH ₂ OH	⁴

¹Cox, D. P.; Terpinski, J.; Lawrynowicz, W. J. *Org. Chem.* 1984, 49, 3216-3219.

²Bennett, B. K.; Harrison, R. G.; Richmond, T. G. *J. Am. Chem. Soc.* 1994, 116, 11165-11166.

³Pilcher, A. S.; Ammon, H. L.; DeShong, P. J. *Am. Chem. Soc.* 1995, 117, 5166-5167.

⁴Akiyama, Y.; Fukuhara, T.; Hara, S. *Synlett* 2003, 1530-1532.

NMR Spectra:

Generation of TBAF—FIG. 1: ¹⁹F NMR spectra recorded over the course of 40 minutes following the mixing of 134 mg TBACN and 9.6 μl C₆F₆ in CD₃CN. The peak at -72 ppm is due to fluoride ion; the peak at -164 ppm is the C₆F₆; the small peak at -147 ppm (d, J_{HF}=148 Hz) is due to HF₂⁻ (The signal marked with * at -151 ppm is an artifact).

Debromofluorination of an aromatic compound—FIG. 2: Conversion of 3,5-bis(trifluoromethyl)bromobenzene to 3,5-bis(trifluoromethyl)fluorobenzene by TBAF in (CD₃)₂SO. a: ¹⁹F NMR spectrum before the addition of 3,5-bis(trifluoromethyl)bromobenzene; b-e: ¹⁹F NMR spectrum after the addition of 3,5-bis(trifluoromethyl)bromobenzene. The total elapsed time was 8 h. Chemical shift assignments: -74 ppm (F⁻), -64 ppm (CF₃), -108 ppm (Ar-F).

Removal of protic solvent by hexacyanobenzene—FIG. 3: ¹⁹F NMR spectra showing the effect of adding 0.08 eq. benzyl alcohol to a solution of in-situ generated TBAF (CD₃)₂SO. a: Spectrum recorded before the addition of benzyl alcohol; b: 5 min after addition of benzyl alcohol; c: 1 h after addition; d:

4 h after addition; e: 20 h after addition. For spectra b and c the bottom spectrum is presented with the normal Y-scale, the top spectrum has the Y-scale multiplied by 8.

Impact of protic solvent in the absence of hexacyanobenzene—FIG. 4: ¹⁹F NMR spectra showing the effect of adding 0.08 eq. benzyl alcohol to a solution of purified TBAF (CD₃)₂SO. a: Spectrum recorded before the addition of benzyl alcohol; b: 10 min after addition of benzyl alcohol; c: 1 h after addition; d: 7 h after addition; e: 20 h after addition. For spectra b and c the bottom spectrum is presented with the normal Y-scale, the top spectrum has the Y-scale multiplied by 64.

Reaction of in-situ generated TBAF with water—FIG. 5: ¹⁹F NMR spectra of the reaction of in situ generated TBAF with 0.083 eq. water in (CD₃)₂SO. a, before addition of water; b-h, after addition of water.

Detail of FIG. 5—FIG. 6: ¹⁹F NMR spectra (expanded area from Figure S-5) of the reaction of in-situ generated TBAF with 0.083 eq water in DMSO-d₆. a, before addition of water; b-h, after addition of water.

13

Reaction of in-situ generated TBAF with water—FIG. 7: ¹H NMR spectra of the reaction of in-situ generated TBAF with 0.083 eq water in (CD₃)₂SO. a, before addition of water; b~f, after addition of water. The signal at 5.6 ppm is assigned to H₂O.

Reaction of isolated TBAF with water—FIG. 8: ¹⁹F NMR spectra of the reaction of isolated TBAF with 0.083 eq water in (CD₃)₂SO. a, before addition of water; b~g, after addition of water.

Detail of FIG. 8—FIG. 9: ¹⁹F NMR spectra of the reaction of isolated TBAF with 0.083 eq water in (CD₃)₂SO. (Detail from Figure S-8.)

Reaction of isolated TBAF with water—FIG. 10: ¹⁹F NMR spectra of the reaction of isolated TBAF with 0.083 eq water in (CD₃)₂SO. a, before addition of water; b~e, after addition of water. The signal at 5.6 ppm is assigned to H₂O; the signal at 5.8 ppm is assigned to HOD.

The invention claimed is:

1. A method of synthesizing an anhydrous fluoride salt having the formula [Q_nM]ⁿ⁺F_x⁻, the method comprising a nucleophilic substitution of a fluorinated aromatic or fluorinated unsaturated organic compound with a salt having the formula [Q_nM]ⁿ⁺A_x⁻ in an inert polar, aprotic solvent wherein:

M is nitrogen or phosphorus;

Q groups are independently selected from the group consisting of an alkyl, an alkenyl, an alkynyl, an aryl, a heteroaryl, and a polymer;

n is an integer such that said [Q_nM] carries at least one formal positive charge;

x is an integer defining the number of formal positive charge(s), +, carried by said [Q_nM];

A⁻ is an anionic nucleophile selected from the group consisting of cyanide, isothiocyanate, isocyanate, cyanate, thiocyanate, alkylthiolates, arylthiolates, or azide; and

F represents fluorine or a radioisotope thereof wherein the anhydrous fluoride salt contains less than 0.01% water of hydration.

2. The method of claim 1 wherein the nucleophilic substitution is conducted at a low temperature.

3. The method of claim 1 conducted at a temperature that is at least -35° C. and no greater than room temperature.

4. The method of claim 1 wherein said inert polar, aprotic solvent is tetrahydrofuran, dimethyl sulfoxide, diethyl ether, dioxane, dimethoxyethane, methyl tert-butyl ether, acetonitrile, acetone, methylethylketone, tetrahydrofuran, dimethyl-

14

formamide, dimethylacetamide, N-methylpyrrolidinone, butyronitrile, toluene, pyridine, benzonitrile, diphenyl ether, diethyl carbonate, or hexamethylphosphoric triamide.

5. The method of claim 1 wherein said fluorinated aromatic compound is a fluorinated benzene and said a fluorinated unsaturated organic compound is a fluorinated alkene or a fluorinated alkyne.

6. The method of claim 1 wherein said [Q_nM]ⁿ⁺A_x⁻ is a tetraalkylammonium cyanide, a trialkylarylammonium cyanide, a dialkyldiarylammonium cyanide, an alkyltriarylammonium cyanide, or a tetraarylammonium cyanide.

7. The method of claim 6 wherein said tetraalkylammonium cyanide is tetrabutylammonium cyanide.

8. The method of claim 1 wherein said fluorinated aromatic compound is hexafluorobenzene, octafluoronaphthalene, octafluorotoluene, pentafluorobenzonitrile, pentafluoropyridine or decafluorobiphenyl.

9. A method for preparation of an anhydrous tetraalkylammonium fluoride salt, the method comprising reacting a fluorinated aromatic compound, in a nucleophilic aromatic substitution reaction, with a tetraalkylammonium cyanide in an inert polar, aprotic solvent, to produce the anhydrous tetraalkylammonium fluoride salt, that contains less than 0.01% water of hydration.

10. The method of claim 9 wherein said tetraalkyl ammonium fluoride salt is tetrabutylammonium fluoride.

11. The method of claim 9 wherein said tetraalkylammonium cyanide is tetrabutylammonium cyanide.

12. The method of claim 9 wherein said fluorinated aromatic compound is hexafluorobenzene.

13. A method for preparation of a fluorinated compound, the method comprising reacting an anhydrous organic fluoride salt, in a nucleophilic substitution reaction, with an alkyl halide, a tosylate, a triflate, a nitroaromatic, a chloroaromatic, or an aromatic triflate compound to produce a fluorinated compound that contains less than 0.01% water of hydration.

14. A method for deprotection of a silylated species, comprising reacting the silylated species with an anhydrous organic fluoride salt that contains less than 0.01% water of hydration to deprotect the silylated species.

15. A method as set forth in claim 14 wherein the silylated species is a silylated aromatic, a silylated heteroaromatic, a silylated alcohol, or a silylated amine.

16. The method of claim 1 wherein [Q_nM]ⁿ⁺ is chiral.

* * * * *