

2008

Factors influencing greater sandhill crane nest success at Malheur National Wildlife Refuge, Oregon

Gary L. Ivey
Department of Fisheries and Wildlife

Bruce D. Dugger
Department of Fisheries and Wildlife

Follow this and additional works at: <http://digitalcommons.unl.edu/nacwgproc>

 Part of the [Behavior and Ethology Commons](#), [Biodiversity Commons](#), [Ornithology Commons](#), [Population Biology Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Ivey, Gary L. and Dugger, Bruce D., "Factors influencing greater sandhill crane nest success at Malheur National Wildlife Refuge, Oregon" (2008). *North American Crane Workshop Proceedings*. 173.
<http://digitalcommons.unl.edu/nacwgproc/173>

This Article is brought to you for free and open access by the North American Crane Working Group at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in North American Crane Workshop Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

FACTORS INFLUENCING GREATER SANDHILL CRANE NEST SUCCESS AT MALHEUR NATIONAL WILDLIFE REFUGE, OREGON¹

GARY L. IVEY, Department of Fisheries and Wildlife, 104 Nash Hall, Oregon State University, Corvallis, OR 97331, USA

BRUCE D. DUGGER, Department of Fisheries and Wildlife, 104 Nash Hall, Oregon State University, Corvallis, OR 97331, USA

Abstract: We used logistic regression to model the effects of weather, habitat, and management variables on Greater Sandhill Crane (*Grus canadensis tabida*) nest success at Malheur National Wildlife Refuge in southeast Oregon. We monitored 506 nests over 9 breeding seasons. Mean apparent nest success was $72\% \pm 4\%$ and varied from 51 to 87%. Nest success was lower one year after a field was burned and declined with nest initiation date. Nest success was higher during warmer springs, in deeper water, and in years with moderate precipitation. Haying, livestock grazing, and predator control did not influence nest success. We suggest the short-term consequence of burning on nest success is outweighed by its long term importance and that water level management is the most important tool for managing crane nest success. Finally, studies of brood ecology are needed to develop a more complete picture of crane nesting ecology.

PROCEEDINGS OF THE NORTH AMERICAN CRANE WORKSHOP 10:157

Key words: breeding ecology, *Grus canadensis tabida*, land management, Oregon, predator control, sandhill crane, weather.

¹See Ivey, G. L., and B. D. Dugger. 2008. Factors influencing nest success of greater sandhill cranes at Malheur National Wildlife Refuge, Oregon. *Waterbirds* 31:52-61.