University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Virology Papers

Virology, Nebraska Center for

1986

HTLV-III Serology in Hemophilia: Relationship with Immunologic Abnormalities

Joan C. Gill

Children's Hospital of Wisconsin

Jay E. Menitove

Community Blood Center, gch@cbckc.org

Philip R. Anderson

James T. Casper

Children's Hospital of Wisconsin

Sushilkumar G. Devare

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/virologypub>

Part of the [Virology Commons](#)

Gill, Joan C.; Menitove, Jay E.; Anderson, Philip R.; Casper, James T.; Devare, Sushilkumar G.; Wood, Charles; Adair, Stuart; Casey, James; Scheffel, Christi; and Montgomery, Robert R., "HTLV-III Serology in Hemophilia: Relationship with Immunologic Abnormalities" (1986). *Virology Papers*. 175.

<https://digitalcommons.unl.edu/virologypub/175>

This Article is brought to you for free and open access by the Virology, Nebraska Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Virology Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Joan C. Gill, Jay E. Menitove, Philip R. Anderson, James T. Casper, Sushilkumar G. Devare, Charles Wood, Stuart Adair, James Casey, Christi Scheffel, and Robert R. Montgomery

HTLV-III Serology in Hemophilia: Relationship with Immunologic Abnormalities

Joan C. Gill, M.D., Jay E. Menitove, M.D.,
Philip R. Anderson, Ph.D., James T. Casper, M.D.,
Sushilkumar G. Devare, Ph.D., Charles Wood, Ph.D.,
Stuart Adair, M.D., James Casey, B.S.,
Christi Scheffel, B.S., and Robert R. Montgomery, M.D.

From the Department of Pediatrics of the Medical College of Wisconsin; Blood Center of Southeastern Wisconsin; Great Lakes Hemophilia Foundation, Milwaukee; Department of Molecular Biology, Abbott Laboratories, North Chicago, Illinois

Supported in part by Grant HL 31015 of the National Heart, Lung, and Blood Institute.
Submitted for publication July 24, 1985; accepted October 3, 1985.

We investigated the relationship of the presence of antibodies to HTLV-III and immunologic abnormalities in patients with hemophilia. Serum antibodies to HTLV-III were analyzed by ELISA assay, immunoprecipitation of labeled cell extracts, and immunoprecipitation of purified HTLV-III p24. Thirty-four (61%) of the total group ($n = 56$) had antibody to HTLV-III; 34 (76%) of 45 patients given commercial factor VIII preparations were seropositive, compared with none of 11 patients treated exclusively with cryoprecipitate obtained from volunteer blood donors. Of patients who were seropositive for HTLV-III antibody, 94% had abnormal T4/T8 ratios, and 33% of those whose serum was antibody negative had abnormal T4/T8 ratios; five patients, each antibody positive, have lymphadenopathy syndrome. Sequential studies in a subset of patients indicate that there is a changing pattern of antibody production to HTLV-III antigens after seroconversion.

Prior to reports linking human T cell lymphotropic virus type III^{1,2} or lymphadenopathy-associated virus^{3,4} with acquired immune deficiency syndrome, we⁵ and others^{6,7} described laboratory abnormalities of immune regulation, particularly lymphocyte subset alteration, in asymptomatic patients with hemophilia. Subsequently, thrombocytopenia, lymphadenopathy syndrome, and other

AIDS-related abnormalities were noted in hemophilic patients.⁸⁻¹⁰ Furthermore, hemophilic patients given concentrates from the United States have been found to have a high prevalence of antibody to LAV/HTLV-III,^{4,11-15} and the appearance of antibody appears to coincide with the AIDS epidemic. To determine whether the immunologic abnormalities are associated with evidence of infection or exposure to the retrovirus, we investigated the presence of serum antibody to one of the isolates, HTLV-III, in patients with hemophilia.

See related article, p. 504, and letter, p. 631.

AIDS	Acquired immune deficiency syndrome
HTLV-III	Human T cell lymphotropic virus type III
ELISA	Enzyme-linked immunosorbent assay
LAV	Lymphadenopathy-associated virus
SDS-PAGE	Sodium dodecylsulfate-polyacrylamide gel electrophoresis

METHODS

All patients enrolled in the comprehensive care program of the Great Lakes Hemophilia Foundation who live within a 60-mile radius of Milwaukee were contacted to participate in the study; 43 patients were selected on

Table 1. HTLV-III antibody in hemophilia

Treatment	Type	n	Positive	
			n	%
Concentrate				
Factor VIII	A severe	32	27	84
	A moderate	3	2	67
	A mild	4	2	50
Factor VIII-IX	A inhibitor	4	3	75
Factor IX	B severe	1	0	0
	B moderate	<u>1</u>	<u>0</u>	0
Total		45	34	76
Cryoprecipitate				
	A severe	10	0	0
	A moderate	<u>1</u>	<u>0</u>	0
Total		11	0	0
Total hemophilia		56	34	61

the basis of availability to have a sample of blood taken during the 2-week period of sampling. Thirteen additional patients were selected from the Green Bay satellite center when they were evaluated for other reasons during the same period. The patients were divided into two groups, those in whom hemorrhages were treated with commercial lyophilized concentrate and those exclusively given cryoprecipitate prepared from the plasma of volunteer blood donors at the Blood Center of Southeastern Wisconsin. Histories were obtained from all patients, and all were specifically examined for the presence of lymphadenopathy and hepatosplenomegaly. Control groups included normal adults, 15 parents of hemophilic boys, five spouses of hemophilic men, and 50 homosexual men seen at a Milwaukee community medical clinic.

Clinical laboratory studies. Complete blood counts and platelet counts were performed on an ELT-8DS Blood Analyzer (Ortho Diagnostic Systems, Westwood, Mass.). White blood cell differential counts were performed by visualization of cells on a Wright-stained blood smear. Serum IgG, IgM, and IgA concentrations were determined by radial immunodiffusion (Helena Laboratories, Beaumont, Texas). Enumeration of lymphocyte subsets was determined by indirect immunofluorescence with OKT monoclonal antibodies incubated with the Ficoll-Hypaque separated mononuclear cell preparation from venous blood, as previously described.⁵

Virus and cells. Uninfected HT-9 cells and HT-9 cells chronically infected with HTLV-III were kindly provided by R. C. Gallo, National Cancer Institute, Bethesda, Maryland. Virus was concentrated one thousandfold, purified by density gradient centrifugation, and disrupted by sonication in the presence of 1% Triton X-100 and 0.05 M

Tris-HCl (pH 9.0). The solution of disrupted virus was clarified by centrifugation at $100,000 \times g$ for 1 hour.

HTLV-III ELISA. In brief, the assay was performed by incubation of test sera with solubilized HTLV-III antigen bound to wells of microtiter plates or antigen-coated polystyrene beads (Abbott Laboratories, North Chicago, Illinois). After washing the plates or beads, alkaline phosphatase-conjugated goat anti-human IgG was added, and after further washing the substrate *p*-nitrophenyl phosphate was used to quantitate the antibody bound.

Immunoprecipitation of labeled cell extracts. Cultures of 10^7 HT-9 cells and 10^7 HT-9 cells chronically infected with HTLV-III were centrifuged and washed twice in methionine- and cysteine-free medium, resuspended in 10 ml of the same medium, and allowed to incubate at 37° C for 30 minutes. Cells were then collected by centrifugation and placed in 10 ml of medium containing 100 μ Ci/ml 35-S methionine and 100 μ Ci/ml 35-S cysteine. Cells were incubated for 4 hours at 37° C in this medium, harvested by centrifugation, and washed twice in complete media. Cells were resuspended in lysis buffer containing 10 mM sodium phosphate, pH 7.5, 100 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, and 0.1% sodium dodecyl sulfate at a concentration of 4×10^6 cells/ml. Ten microliters of test serum was incubated with 200 μ l aliquots of lysate for 30 minutes at 0° C; *Staphylococcus aureus* protein A bound to Sepharose beads (Pharmacia, Piscataway, N.J.) was then used to recover antigen-antibody complexes, which were then analyzed by SDS-PAGE as previously described.¹⁷

p-24 Radioimmunoassay. Double-antibody precipitation with ¹²⁵I-labeled HTLV-III p24 was performed as previously described.¹⁸ In brief, the p24 protein was purified to homogeneity by phosphocellulose column chromatography¹⁸ and labeled with ¹²⁵I (30 μ Ci/ μ g) by Iodo-beads (Pierce Chemical Co., Rockford, Ill.). Serial dilutions of test sera were incubated with ¹²⁵I-labeled purified p24, followed by precipitation of antigen-antibody complexes by incubation with anti-human IgG antisera, centrifugation to recover precipitates, and counting in an ANSA gamma counter (Abbott Laboratories). A sample was considered positive if precipitated counts were five times background.

RESULTS

Antibody to HTLV-III was detected by ELISA in 34 (61%) of 56 hemophilic patients (Table I). Among 45 patients who received commercial lyophilized concentrate, 34 (76%) had antibody to HTLV-III, compared with none of 11 patients treated exclusively with cryoprecipitate. Neither of the two patients with hemophilia B treated with factor IX concentrate had antibody, but three of four patients with inhibitors to factor VIII, all treated with

Fig. 1. Immunoprecipitation of ^{35}S methionine- and ^{35}S cysteine-labeled HTLV-III antigens by sera from patients with hemophilia. Molecular weight standards (Kd) are in lane A, with corresponding molecular weight to left of autoradiogram; molecular weights of HTLV-III antigens are shown on right. Immunoprecipitation of extract of HT-9 cells without HTLV-III is shown in lane J. Note variable patterns of reactivity among patients. All patient sera immunoprecipitate gp 160/120. Patient D serum has antibody to p24 but decreased reactivity to pp55; patient B serum precipitates gp 120/160 only.

both factor VIII and factor IX concentrate, had antibody.

All patients with positive results by ELISA had antibody to viral glycoproteins gp120 and gp 160, as demonstrated by radioimmunoprecipitation of labeled cell extracts (Figure 1). Similarly, all patients who had positive ELISA results had antibodies capable of recognizing ^{125}I -labeled HTLV-III p24; however, titers of these anti-p24 antibodies varied as much as a thousandfold. Those with low antibody titers to purified p24 also lacked detection of anti-p24 by radioimmunoprecipitation of labeled cell extracts. Furthermore, the ability of these sera to precipitate pp55 varied (Figure 1). This result is not unexpected, because pp55 is a precursor protein consisting of the p24, p17, and p15 *gag* proteins of the virus. Thus, sera with a low titer to p24 may be unable to precipitate quantities of pp55 if antibodies to other *gag* proteins are not present.

Chi-square analysis of immunologic status and HTLV-III serology revealed five statistically significant factors (Table II). Each of five patients with persistent generalized lymphadenopathy had HTLV-III antibody. Decreased platelet counts, decreased T4 lymphocytes, decreased T4/T8 ratios, and elevated serum IgG concentrations were associated with antibody to HTLV-III, whereas abnormal IgM and IgA and elevated T8 lymphocytes were not associated with HTLV-III antibodies.

phocytes were not associated with HTLV-III antibodies.

Sequential studies were performed on samples stored since January 1982 from 17 of the patients who were seropositive for HTLV-III; only three had results on earlier samples. In November 1982, patient 1 was seronegative for HTLV-III antibody and had a normal T4/T8 ratio. Two months later, antibody was detectable and his T4/T8 ratio decreased. Antibody and T4/T8 ratio have remained abnormal, and in May 1984, 17 months after the appearance of HTLV-III antibodies, persistent lymphadenopathy developed. Patient 2 had a similar course, with seroconversion in October 1982, a decrease in T4/T8 ratio at his next evaluation, and development of generalized lymphadenopathy 22 months later. After the appearance of antibody to HTLV-III in January 1983, a third patient developed an abnormal T4/T8 ratio in November 1983, and had an episode of localized herpes zoster in February 1985.

The pattern in these three patients of serum reactivity to various HTLV-III viral proteins by immune precipitation of HTLV-III proteins is demonstrated in Figure 2. None of the initial samples precipitated viral proteins, whereas all three precipitated the gp 160-120, pp55, and p24 viral proteins when studied most recently. The patients

Fig. 2. Immunoprecipitation in four hemophilic patients studied sequentially. *Early* indicates first serum sample obtained, and *late*, most recent. *Lanes A, C, and D* contain serum from patients 3, 1, and 2, respectively. Note absence of reactivity with viral proteins in early samples. *Lane B* contains serum from patient with initially positive HTLV-III ELISA in whom lymphadenopathy and thrombocytopenia developed; note decreased reactivity with pp55 and p24 in late sample.

with lymphadenopathy precipitated relatively more pp55 and p24 than did the patient with herpes zoster. In contrast, a fourth patient, initially seropositive at the onset of the study, had an early response to p24, which was absent 18 months later; lymphadenopathy also has and more recently the onset of significant thrombocytopenia (platelet count $<50,000/\mu\text{l}$).

Selected family members were examined to some insight into the transmissibility of HTLV-III. Fifteen parents of hemophilic boys (the sons of 10 were seropositive for HTLV-III antibody) who prepared and administered treatment product were seronegative for HTLV-III antibody. Similarly, none of five spouses of hemophilic patients (two positive for HTLV-III antibody) had HTLV-III antibody. In contrast, 14 (28%) of 50 homosexual men seen at a community medical clinic in Milwaukee were seropositive for HTLV-III by ELISA.

DISCUSSION

Evidence for a human retroviral cause for AIDS has recently accumulated. The original isolates, HTLV-III and LAV, are likely the same virus¹⁹ and have been cultured from blood, semen, and saliva from patients with AIDS or lymphadenopathy syndrome and from asymptomatic individuals from high-risk groups.^{1,3,4,11,20-27} Antibody to these viruses has been detected in the serum of

Table II. Relationship of HTLV-III antibody to immunologic abnormalities in hemophilia

	HTLV-III positive	HTLV-III negative
Platelet count/ μl		
<150,000	5	0
$\geq 150,000$	16	24
	$\chi^2 = 6.4; P < 0.02$	
T4/T8 ratio		
≤ 1.5	30	8
> 1.5	2	16
	$\chi^2 = 22.9; P < 0.01$	
Total T4 lymphocytes/ μl		
< 630	13	3
≥ 630	14	21
	$\chi^2 = 7.5; P < 0.01$	
Serum IgG (mg/dl)		
> 1500	9	0
≤ 1500	13	18
	$\chi^2 = 9.5; P < 0.01$	
Lymphadenopathy		
Present	5	0
Absent	21	22
	$\chi^2 = 4.7; P < 0.05$	

both asymptomatic and ill individuals from these risk groups.^{1,3,4,12-16} Antibody to HTLV-III/LAV has been detected in 34%¹¹ to 94%¹² asymptomatic hemophilic patients, a group at high risk for AIDS. In this study, 61% of hemophilic patients had evidence of prior exposure to HTLV-III/LAV. It is not known whether antibody seroconversion occurred in response to exposure to live virus and reflects or resolved infection, or resulted from exposure to inactivated virus in clotting factor replacement products. However, the presence of lymphadenopathy syndrome, thrombocytopenia, herpes zoster, and laboratory evidence of immune dysregulation similar to that associated with AIDS suggests that these patients have been exposed to live virus. Inasmuch as none of these patients has developed the full clinical symptoms of AIDS, and may have future resolution rather than progression of immunologic dysfunction, any speculation regarding the protective nature of various antibody patterns would be premature. Nevertheless, whereas only 4% of antibody-positive patients have normal T4/T8 ratios, 33% of antibody-negative patients have abnormal T4/T8 ratios (Table II). This implies that a spectrum of *in vitro* laboratory abnormalities, associated with factor VIII concentrate therapy, occurs independent of HTLV-III antibody positivity. In the light of evidence suggesting that antigenemia can exist in asymptomatic antibody-negative individuals,²⁶ this interpretation is somewhat tentative pending the availability of antigen testing.

HTLV-III/LAV appears to be transmitted by intimate sexual contact and by exposure to blood products, and not by casual contact such as typical exposure to family members other than spouses. None of the parents of hemophilic patients tested had antibody to HTLV-III. The five spouses of hemophilic patients (two antibody positive) reported had negative HTLV-III serologic findings. However, continued follow-up of this group is needed so that accurate counseling can be provided in the future; evidence for infection with HTLV-III via heterosexual transmission is now emerging.²⁸⁻³⁰

Hemophilic patients given nonheated commercial factor VIII concentrates are at high risk for development of antibody to HTLV-III and associated immunologic abnormalities. Patients given cryoprecipitate prepared from the plasma of Southeastern Wisconsin blood donors appear to be at lower risk. Preliminary evidence suggests that heat treatment of clotting factor concentrates significantly reduces retrovirus infectivity,³¹⁻³² nonheated concentrates should not be used for treatment of in patients with hemophilia.

REFERENCES

- Popovic M, Sarngadharan MG, Read E, Gallo RC. Detection, isolation, and continuous production of cytopathic retroviruses (HTLV-III) from patients with AIDS and pre-AIDS. *Science* 1984; 224: 497-500.
- Sarngadharan MG, Popovic M, Bruch L, Schupbach J, Gallo RC. Antibodies reactive with human T-lymphotropic retroviruses (HTLV-III) in the serum of patients with AIDS. *Science* 1984; 224: 506-508.
- Barré-Sinoussi F, Chermann JC, Rey F, et al. Isolation of a lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science* 1983; 220: 868-871.
- Moniagnier L, Chermann JC, Barré-Sinoussi F, et al. A new human T-lymphotropic retrovirus: characterization and possible role in lymphadenopathy and acquired immune deficiency syndrome. In: Gallo RC, Essex M, Gross L, eds. *T-Cell Leukemia/Lymphoma Virus* 1984. Cold Spring Harbor, N.J. Cold Spring Harbor Laboratory, 1984: 363-379.
- Menitove JE, Aster RH, Casper JT, et al. T-lymphocyte subpopulations in patients with classic hemophilia treated cryoprecipitate and lyophilized concentrates. *N Engl J Med* 1983; 308: 83-86.
- Lederman MM, Ratnoff OD, Scillian JJ, Jones PK, Schacter B. Impaired cell-mediated immunity in patients with classic hemophilia. *N Engl J Med* 1983; 308: 79-83.
- Goldsmith JC, Moseley PL, Monick M, Brady M, Hunningbake GW. T-lymphocyte subpopulation abnormalities in apparently healthy patients with hemophilia. *Ann Intern Med* 1983; 98: 294-296.
- Ratnoff OD, Menitove JE, Aster RH, Lederman MM. Coincident classic hemophilia and "idiopathic" thrombocytopenic purpura in patients under treatment with concentrates anti-hemophilic factor (factor VIII). *N Engl J Med* 1983; 308: 439-432.
- Ragni MV, Lewis JH, Spero JA, Bontempo FA. Acquired immunodeficiency-like syndrome in two hemophiliacs. *Lancet* 1983; 7: 213-214.
- Gill JE, Menitove JE, Wheeler D, Aster RH, Montgomery RR. Generalized lymphadenopathy and T-cell abnormalities in hemophilia A. *J Pediatr* 1983; 103: 18-22.
- Cheingsong-Popov R, Weiss RA, Dalgleish A, et al. Prevalence of antibody to human T-lymphotropic virus type III in AIDS and AIDS-risk patients in Britain. *Lancet* 1984; 2: 477-480.
- Melbye M, Froebel KS, Madhok R, et al. HTLV-III seropositivity in European haemophiliacs exposed to factor VIII concentrate imported from the USA. *Lancet* 1984; 2: 1,444-1,446.
- Evatt BL, Gomperts ED, McDougal SJ, Ramsey RB. Coincidental appearance of LAV/HTLV-III antibodies in hemophiliacs and the onset of the AIDS epidemic. *N Engl J Med* 1985; 312: 483-486.
- Goedert JJ, Sarngadharan M, Eyster ME, et al. Antibodies reactive with human T-cell leukemia viruses in the serum of hemophiliacs receiving factor VIII concentrate. *Blood* 1982; 312: 483-486.
- Lederman MM, Ratnoff OD, Evatt BL, McDougal JS. Acquisition of antibody to lymphadenopathy-associated virus in patients with classic hemophilia (factor VIII deficiency). *Ann Intern Med* 1985; 102: 753-757.
- Eyster ME, Goedert JJ, Sarngadharan MG, Weiss SH, Gallo RC, Blattner WA. Development and early natural history of HTLV-III antibodies in persons with hemophilia. *JAMA* 1985; 253: 2,219-2,223.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227: 680-685.
- Devare SG, Stephenson JR, Sarma PS, Aaronson SA, Chander S. Bovine lymphosarcoma: development of a radioimmunologic technique for detection of the immunologic agent. *Science* 1976; 194: 1,428-1,430.
- Ratner L, Gallo RC, Wong-Staal F. HTLV-III, LAV, ARV are variants of same AIDS virus. *Nature* 1985; 313: 636-637.
- Barré-Sinoussi F, Mathur-Wagh U, Rey F, et al. Isolation of lymphadenopathy-associated virus (LAV) and detection of LAV antibodies from U.S. patients with AIDS. *JAMA* 1985; 253: 1,737-1,739.
- Vilmer F, Barré-Sinoussi F, Rouzioux C, et al. Isolation of new lymphotropic retrovirus from two siblings with hemophilia B, one with AIDS. *Lancet* 1984; 1: 753-757.
- Gallo RC, Salahuddin SZ, Popovic M, et al. Frequent detection and isolation of cytopathic retroviruses (HTLV-III) from patients with AIDS and at risk for AIDS. *Science* 1984; 224: 500-503.
- Feorino PM, Kalyanaraman VS, Haverkas HW, et al. Lymphadenopathy-associated virus infection of a blood donor recipient pair with acquired immunodeficiency syndrome. *Science* 1984; 225: 69-72.
- Zagury D, Bernard J, Leibowitch J, et al. HTLV-III in saliva of people with AIDS-related complex and healthy homosexual men at risk for AIDS. *Science* 1984; 226: 447-451.
- Ho DD, Schooley RT, Rata TR, et al. HTLV-III in the semen and blood of a healthy homosexual man. *Science* 1984; 226: 451-453.
- Salahuddin SZ, Groopman JE, Markham PD, et al. HTLV-III in symptom-free seronegative persons. *Lancet* 1984; 2: 1,418-1,420.

27. Gomperts ED, Feorino P, Evatt BL, *et al.* LAV/HTLV-III presence in peripheral blood lymphocytes of seropositive young hemophiliacs. *Blood* 1985; 65: 1,549-1,552.
28. Pitchenik AE, Shafron RD, Glasser RM, Spira TJ. The acquired immunodeficiency syndrome in the wife of a hemophiliac. *Ann Intern Med* 1984; 100: 62-65.
29. Redfield RR, Markham PD, Salahuddin SZ. Frequent transmission of HTLV-III among spouses of patients with AIDS-related complex and AIDS. *JAMA* 1985; 253: 1,571-1,573.
30. Kreiss JK, Kitchen LW, Prince HE, Kasper CK, Essex M. Antibody to human T-lymphotropic virus type III in wives of hemophiliacs: evidence for heterosexual transmission. *Ann Intern Med* 1985; 102: 623-626.
31. Levy JA, Mitra G, Mozen MM. Recovery and inactivation of infectious retrovirus from factor VIII concentrates. *Lancet* 1984; 1: 722-723.
32. Spire B, Dormont D, Barré-Sinoussi F, Montagnier L, Chermann JC. Inactivation of lymphadenopathy-associated virus by heat, gamma rays, and ultraviolet light. *Lancet* 1984; 1: 188-189.