

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Nebraska Swine Reports

Animal Science Department

1996

The Effects of Tallow Addition to the Diets of Lactating Sows on Hormone and Metabolite Concentrations

Paul Ermer

University of Nebraska-Lincoln

Scott Tilton

University of Nebraska-Lincoln

Phillip S. Miller

University of Nebraska-Lincoln, pmiller1@unl.edu

Austin Lewis

University of Nebraska-Lincoln, alewis2@unl.edu

Cynthia Wolverton

University of Nebraska-Lincoln, cynthia.wolverton@unl.edu

Follow this and additional works at: https://digitalcommons.unl.edu/coopext_swine



Part of the [Animal Sciences Commons](#)

Ermer, Paul; Tilton, Scott; Miller, Phillip S.; Lewis, Austin; and Wolverton, Cynthia, "The Effects of Tallow Addition to the Diets of Lactating Sows on Hormone and Metabolite Concentrations" (1996). *Nebraska Swine Reports*. 195.

https://digitalcommons.unl.edu/coopext_swine/195

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Swine Reports by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



There was a limited number of sows in this study, and sow and litter performance was quite variable (Table 2). These data are included primarily to aid in interpretation of the metabolite responses.

No differences were observed in baseline concentrations of NEFA or glucose among dosages of epinephrine (Table 3). This was expected because epinephrine is metabolized rapidly, therefore the previous dosage should not affect baseline concentrations on the next day.

Peak NEFA concentration in plasma exhibited a quadratic response ($P < .05$) to epinephrine dosage, with a decline from the .1 $\mu\text{g}/\text{kg}$ dosage to the .4 $\mu\text{g}/\text{kg}$ dosage, followed by increases in NEFA concentration to the 1.6 $\mu\text{g}/\text{kg}$ dosage. When adjusted for

baseline, peak height increased linearly ($P < .01$), suggesting that variation in baseline concentration (although not significant) was contributing to the quadratic effect observed with the unadjusted peak concentrations. Response area for NEFA increased linearly ($P < .01$) with some tendency ($P < .06$) for a quadratic response to increasing doses of epinephrine, following a pattern similar to that observed for peak height.

Peak glucose concentration increased linearly with increasing dosage of epinephrine ($P < .01$). This was also observed for the adjusted peak concentration ($P < .01$) and the glucose response area ($P < .01$). These data show that the sow responds to increasing dosages of epinephrine by increasing glucose release into plasma from glycogen stores and from increased

gluconeogenesis.

Conclusions

The lactating sow is able to increase energy mobilization from body tissues in response to administration of increasing dosages of epinephrine. The optimal dosage of epinephrine was not established. This experiment has shown that the optimal dosage is higher than the dosage used previously in sows or the optimal dosage for the dairy cow (.4 $\mu\text{g}/\text{kg}$).

¹Scott L. Tilton and Paul M. Ermer are graduate students, Austin J. Lewis is a Professor, Phillip S. Miller is an Assistant Professor, and Cynthia K. Wolverton is a Research Technologist, Department of Animal Science.

The Effects of Tallow Addition to the Diets of Lactating Sows on Hormone and Metabolite Concentrations

Paul M. Ermer
Scott L. Tilton
Phillip S. Miller
Austin J. Lewis
Cynthia K. Wolverton¹

was observed for sows consuming either diet. These results show the addition of tallow to lactation diets does not affect the concentrations of glucose, NEFA, insulin or glucagon in the fed state.

The objective of the following experiment was to measure changes in feed and energy intake and meal patterns associated with the addition of tallow to lactation diets. In addition, nonfasting (fed) concentrations of hormones and metabolites were measured.

Summary and Implications

The metabolic responses of sows fed a corn-soybean meal or a corn-soybean meal-10% tallow diet were measured. The addition of tallow to lactating sows diets had no effect on feed or energy intake. In addition, there were no effects on the concentrations of glucose, nonesterified fatty acids (NEFA), insulin, or glucagon. No differences in either the time spent consuming feed or the number of meals consumed were observed. Finally, no linear association between eating time and area under the curve for insulin

Introduction

In previous research (Nebraska Swine Report, 1995) we have reported several behavioral and physiological responses to the addition of tallow to lactating sow diets. Although there were no differences in feed or energy intake, the addition of tallow resulted in increased rate of feed consumption and decreased the percentage of time spent consuming feed. Fasting concentrations of glucose were increased and nonesterified fatty acids (NEFA) and glucagon were decreased in sows fed the tallow diet.

Methods

Eight first-litter crossbred gilts were used. Gilts were randomly and equally allotted within room (two rooms) to receive either a corn-soybean meal (C-SBM) or a corn-soybean meal-10% tallow diet (Tallow, Table 1). Dietary treatments were initiated after parturition. Sow weight postpartum averaged 389 lb. Farrowing room temperature averaged 72° F and continuous lighting was provided. Sow and litter weights were obtained after parturition and on d 21 of lactation. Feed intake was



Table 1. Diet composition (as-fed basis)^a

Ingredient, %	C-SBM	Tallow
Corn	67.90	56.80
Soybean meal, 46.5%	28.00	29.00
Dicalcium phosphate	2.10	2.30
Salt	.50	.50
Limestone	.40	.30
Vitamin mix	1.00	1.00
Trace mineral mix	.10	.10
Tallow	—	10.00
Formulated composition		
Metabolizable energy, Mcal/lb	1.48	1.69
Crude protein, %	18.8	18.3
Lysine, %	1.02	1.02
Calcium, %	.88	.89
Phosphorous, %	.76	.77

^aDiets are corn-soybean meal (C-SBM) and corn-soybean meal-10% tallow (Tallow).

measured daily for 21 days. Litter size was standardized within 3 days of parturition.

On d 7 and 21 of lactation, meal patterns were measured continuously from 6:00 a.m. to 2:00 p.m. Sows were considered to be feeding when observed chewing with their head in the feeder. Meals were considered to be periods of feeding separated by at least 20 minutes. All feeding periods meeting this criterion were considered to be meals.

Catheters were inserted into the vena cava to enable frequent blood collection. Blood was collected every 15 minutes from 6:00 a.m. to 2:00 p.m. on d 7 and 21 of lactation. Plasma was separated and analyzed for insulin,

glucagon, glucose, and NEFA.

Average hormone and metabolite concentrations were calculated from values obtained at all time points. Areas under the curve (AUC) for insulin are the sum of the average concentrations of two consecutive samples multiplied by the elapsed time. All samples collected on d 7 and 21 were used to calculate AUC for insulin.

Data were analyzed as a randomized complete block. Because of catheter failure, one sow each on d 7 and 21 was not included in the data set.

Results

There were no differences in feed or energy intake (Figure 1). Numerically, both feed and energy intake were lower (23 and 10%, respectively) in

sows fed the Tallow diet than in sows fed the C-SBM diet.

Because of the small number of sows on each treatment, sow and litter performance was averaged across treatments. These data are included to aid in interpretation of hormone and metabolite responses. On average, sows lost 39.3 lb during lactation. Litter weight gain averaged 109.1 lb and litter size at weaning was 9.7 pigs.

No differences in the time spent consuming feed or the number of meals consumed were observed (Table 2).

Average concentrations of glucose or NEFA were not affected by diet (Table 3). Glucose concentrations decreased from d 7 to 21 in sows fed both diets ($P = .01$). Although there was no effect of day of lactation on NEFA

(Continued on next page)

Table 2. Meal patterns of gilts fed a corn-soybean meal (C-SBM) or a corn-soybean meal-10% tallow (Tallow) diet^a

Criteria	Day 7		Day 21		P ^b (Diet)	P (Day)	SE ^c
	C-SBM	Tallow	C-SBM	Tallow			
No. of sows	3	4	4	3			
Time spent consuming feed, min	40.50	37.78	42.00	39.88	NS	NS	8.63
Number of meals consumed	3.25	4.88	4.00	4.13	NS	NS	.79

^aResponses were measured from 6:00 am to 2:00 pm

^bNS is not significant ($P > .05$)

^cSE = standard error




Figure 1. Feed and energy intake of sows fed either a corn-soybean meal (C-SBM) or a corn-soybean meal-10% tallow (Tallow) diet.




Table 3. Average hormone and metabolite concentrations of gilts fed a corn-soybean meal (C-SBM) or a corn-soybean meal-10% tallow (Tallow) diet^a

Criteria	Day 7		Day 21		P ^b (Diet)	P (Day)	SE ^c
	C-SBM	Tallow	C-SBM	Tallow			
No. of sows	3	4	4	3			
Glucose, mmol/L	5.24	5.39	4.55	4.90	NS	.01	.17
NEFA ^d , μ Eq/L	415.92	703.53	352.22	418.50	NS	NS	82.99
Insulin, pmol/L	199.39	202.19	201.34	178.18	NS	NS	27.57
Glucagon, ng/L	68.17	67.88	65.24	78.73	NS	NS	9.71
Insulin:Glucagon, mol:mol	11.58	12.26	13.64	10.24	NS	NS	3.04

^aConcentrations are averages of samples drawn every 15 min from 6:00 am to 2:00 pm

^bNS is not significant ($P > .05$)

^cSE = standard error

^dNEFA is nonesterified fatty acids

concentration, there was a 37% decrease from d 7 to 21 of lactation. Average concentrations of insulin, glucagon, and the insulin:glucagon ratio were not affected by diet or day of lactation (Table 3).

No correlation was detected between AUC for insulin and time spent eating ($r = .38$, Figure 2).

Discussion

Feed and energy intake was lower in this experiment than in a previous experiment. Low feed and energy intakes were associated with increased sow weight loss.

Contrary to results of a previous experiment, no effect of diet on per-

centage of time spent consuming feed was observed. This may be due to the shorter period of time in which meal patterns were measured (8 versus 24 h).

Time spent consuming feed, when expressed as a percentage (8.3%), is similar to values reported previously over a 24-hour period. We expected the percentage of time spent consuming feed to be higher because previous research indicates that sows consume a larger proportion of meals during the day than at night.

There were several discrepancies in hormone and metabolite concentrations observed between this and a previous experiment. First, there were no effects of diet on average concen-

trations of glucose, NEFA, or glucagon in this experiment. Second, greater weight loss in the present experiment was associated with lower concentrations of glucagon and higher insulin:glucagon ratios. It is difficult to compare results of experiments where sows are in a fed versus a fasted state. Both feeding and energy balance affect hormones and metabolites in the fed state, whereas only energy balance affects hormones and metabolites in the fasted state. Whether hormone concentrations in the fed or fasted state are a better indicator of changes in metabolic status during lactation is unknown.

Average concentration of NEFA decreased from d 7 to 21 of lactation. This indicates a decrease in fat mobilization from body reserves as lactation progressed. The large sow weight losses during lactation indicate that body fat reserves may have been significantly depleted.

Several factors may have contributed to the low correlation between time spent feeding and AUC for insulin. First, time spent eating may not reflect feed intake across treatments. In a previous experiment, differences in rate of feed consumption were observed. Secondly, factors other than eating, such as energy balance, influence insulin concentration.

These results, and those of a previous experiment, indicate inconsistent responses in feed and energy intake to the addition of tallow to lactating sow diets. Changes in fasting hormones and metabolites previously observed in sows fed diets containing tallow were not observed in the fed state. Future work will examine hormone and metabolite profiles in response to feeding and investigate other metabolic signals that may influence feeding behavior or energy intake in lactating sows.




Figure 2. Correlation of area under the curve (AUC) for insulin to eating time for sows fed either a corn-soybean meal (C-SBM) or a corn-soybean meal-10% tallow (Tallow) diet.

¹Paul M. Ermer and Scott L. Tilton are graduate students, Phillip S. Miller is an Assistant Professor, Austin J. Lewis is a Professor, and Cynthia K. Wolverton is a Research Technologist, Department of Animal Science.