

University of Nebraska - Lincoln

## DigitalCommons@University of Nebraska - Lincoln

Erforschung biologischer Ressourcen der  
Mongolei / Exploration into the Biological  
Resources of Mongolia, ISSN 0440-1298

Institut für Biologie der Martin-Luther-  
Universität Halle-Wittenberg

2016

### New Data on Feeding Ecology of *Bubo bubo* and *Asio otus* (Aves: Strigidae) in Mongolia

Michael Stubbe

*Martin-Luther-Universität*

Nayamsuren Batsaikhan

*National University of Mongolia*, microtus\_b@yahoo.com

Oliver Lindecke

*Martin-Luther-Universität*

Ravčigijn Samjaa

*National University of Mongolia*, r.samjaa@yahoo.com

Annegret Stubbe

*Martin-Luther-Universität*, annegret.stubbe@zoologie.uni-halle.de

Follow this and additional works at: <https://digitalcommons.unl.edu/biolmongol>

 Part of the Asian Studies Commons, Biodiversity Commons, Ecology and Evolutionary Biology Commons, Environmental Sciences Commons, Nature and Society Relations Commons, Ornithology Commons, and the Other Animal Sciences Commons

Stubbe, Michael; Batsaikhan, Nayamsuren; Lindecke, Oliver; Samjaa, Ravčigijn; and Stubbe, Annegret,  
"New Data on Feeding Ecology of *Bubo bubo* and *Asio otus* (Aves: Strigidae) in Mongolia" (2016).

*Erforschung biologischer Ressourcen der Mongolei / Exploration into the Biological Resources of Mongolia*, ISSN 0440-1298. 183.

<https://digitalcommons.unl.edu/biolmongol/183>

This Article is brought to you for free and open access by the Institut für Biologie der Martin-Luther-Universität Halle-Wittenberg at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Erforschung biologischer Ressourcen der Mongolei / Exploration into the Biological Resources of Mongolia, ISSN 0440-1298 by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Erforsch. biol. Ress. Mongolei (Halle/Saale) 2016 (13): 301-311

## New data on feeding ecology of *Bubo bubo* and *Asio otus* (Aves: Strigidae) in Mongolia<sup>1</sup>

M. Stubbe, N. Batsajchan, O. Lindecke, R. Samjaa & A. Stubbe

### Abstract

New materials of 10 Eagle owl *Bubo bubo* localities from Mongolia were analyzed. Feeding ecology of Long-eared owl *Asio otus* could be studied by investigation of pellets from 4 roosting or nesting places in S-Mongolia.

All together in the prey of *Bubo bubo* were analysed 1098 vertebrates (1017 mammals, 39 birds, 41 reptiles 1 fish), involved about 22 mammal species. In the pellets were dominant Lagomorpha with 23.3 % in SW-Mongolia, Dipodidae in W-, SW- and S-Mongolia (37.3 %, 23.4 %, 30.2 %), Cricetidae in SW- and S-Mongolia (13.6 %, 17.6 %), Gerbillidae in W- and S-Mongolia (19 %, 32.1%) and Arvicolidae in W- and SW-Mongolia (23.8%, 23.8%). The percentage of birds was registered between 3.2 and 4.5 %. Rests of evertebrates (Arachnida, Solifugae, Scorpiones, Coleoptera, Orthoptera) were noticed in rare cases.

For *Asio otus* 747 vertebrates (726 mammals, 12 birds, 9 reptiles) as prey were determined. There were analysed about 13 mammal species. In the prey of the Longeard Owl are dominant the small Cricetidae species (*Cricetus* and *Phodopus* with 50.5 %) as well as Gerbillidae (*Meriones* spec. 29 %) and the small Dipodidae (*Salpingotus* and *Cardiocranus*) with 12.3 %.

**Key words:** *Bubo bubo*, *Asio otus*, Mongolia, feeding ecology

### Introduction

First results on feeding ecology of *Bubo bubo* were published by STUBBE & CHOTOLCHU (1968), PIECHOCKI et al. (1977), STUBBE et al. (1989) and HOFMANN et al. (2005). The new pellets were collected from 10 localities, especially in W-Mongolia. In 2007 JAVKHLANTSETSEG et al. published results on nest site and food composition of the Eagle Owl in Central and Northern Mongolia. They identified 276 prey from pellets and remains from nests. 72.1 % were mammals, 26.1 % birds, 0.7 % amphibians, and 1.1 % beetles. *Bubo bubo* is nesting in rocky mountains, steppe hills with broken rocks, lake shores with cliffs and river banks, forested areas and river valleys in all natural zones and belts in the country (GOMBOBAATAR & MONKS 2011). All nesting places which we found were in rock crevices or small caves. TUMURBAT et al. (2009) found five nests in cliffs, three in rock columns, two in sandy precipices and one in a rock ledge.

*Asio otus* is nesting in tree holes, deserts nests of Carrion Crow, Black-billed Magpie and raptors in various types of forests including coniferous and mixed forests in mountain taiga forest, forest steppe, patchy woodland in the steppe and lake and river valleys (GOMBOBAATAR & MONKS 2011). The species is also breeding in the arid Gobi zone.

The feathers of the Eagle Owl and Long-eared Owl have a high value in traditional custom and use, especially by the Kazakh inhabitants in the most western Aimag Bajan-Ölgij (see also PIECHOCKI 1980). As mascot you can find the feathers nearly in every ger at predestined places, near altarpieces or above the baby cradle. In Bajan-Ölgij it's possible to buy the feathers on the market or in shops (fig. 1). It means also that we have a conflict between old traditional behaviour and modern nature conservation.

<sup>1</sup> Results of the Mongolian-German Biological Expeditions since 1962, No. 337.



Fig. 1: Feathers, claws and whole skin of Eagle owl (in Mongolian: Shar shuvuu) and Long-eard owl (in Mongolian: Sooton guivanga) on the market and in gers - talismans of the Kazakh people in West-Mongolia (photos: A. & M. STUBBE).

## Material and results

New materials from 4 of our old and pellets from 5 new Eagle owl localities, especially from S-Mongolia, were analysed now. At first, all skull remains were selected and determined.

Feeding ecology of *Asio otus* could be studied by investigation of pellets from 4 roosting or nesting places in S-Mongolia. *Asio otus* is distributed and breeding there in all greater dry desert valleys (sajrs, fig. 3) and oasis, stocked with *Ulmus pumila* or *Populus diversifolia*, maybe also in saxaul woodlands (*Haloxylon ammodendron*) of the desert. In Mongolia, the Long-eared Owl is a breeding visitor and a partial migrant. Migration routes and wintering localities are not investigated. According to GOMBOBAATAR & MONKS (2011), most of the breeding individuals arrive in the breeding sites by late April/early May.



Fig. 2: Typical breeding and feeding habitats of *Bubo bubo* in western (above: formerly valley of Čonocharajchijn-gol 1975, nowadays a hydroelectric power station) and southwestern Mongolia (below: upper course of Bulgan-gol, 2004); photos: M. STUBBE.



Fig. 3: Sajr Šutegijn Bajan-gol (Sum Manlai, South-Mongolia) - breeding and feeding habitat of Eagle owl and Long-eared owl (photo: M. STUBBE).




Fig. 4: Breeding localities of *Bubo bubo*: 1 - Ulaan-uul, 20 km W Chovd-chot, 2 - Čonocharajchijn-gol, 3 - Urd-Tamir-gol (Cencher Somon), 4 - Ugji-nuur, 5 - Arc-bogd (Gobi-Altai), 6 - Sajchany-Zaravč (50 km ENE Ulaanbaatar), 7 - Bodončijn-gol (upper course), Mongolian Altai; 8 - Bodončijn-gol (lower course), Dzungarian Gobi; 9 - Bulgan-gol (upper course), 10 - Guntamga, Dzungarian Gobi; 11 - Minž-gol, 12 - Ich Gazarijn Čuluu, 13 - Šutegijn Bajan-gol, 14 - 10 km SE Chanbogd, 15 - Undagijn-gol (Galbyn-gobi), 16 - Galbyn-gobi.

Table 1: Analyzed feeding composition of *Bubo bubo*

finding locations	2	8	9	10	11	12	12	12	13	13	13	14	15	16
year	2002	2002	2002	2002	2002	2004	2005	2010	2009	2010	2011	2005	2009	2004
<b>MAMMALIA</b>														
<b>Insectivora</b>														
<i>Hemiechinus auritus</i>	1	-	1	-	-	-	1	1	5	2	1	1	11	-
<i>Sorex</i> spec.	1	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>Lagomorpha</b>														
<i>Ochotona</i> spec.	35	14	19	-	-	-	-	-	3	1	-	-	-	-
<i>Lepus tolai</i>	9	6	2	-	-	-	1	1	2	2	1	1	1	-
<b>Chiroptera</b>														
<i>Myotis</i> spec.	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Chiroptera indet.	3	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>Rodentia</b>														
DIPODIDAE														
<i>Dipus/Allactaga</i> spec.	143	5	30	2	-	-	4	5	52	25	22	8	9	-
<i>Salpingotus/Cardiocranius</i>	22	-	4	-	-	-	-	-	33	7	4	1	-	5
CRICETIDAE														
<i>Cricetus/Phodopus</i> spec.	19	-	13	-	-	-	-	3	65	2	4	-	1	5
<i>Allocricetus</i> spec.	4	-	11	-	-	-	-	-	3	-	-	-	-	-
GERBILLIDAE														
<i>Rhombomys opimus</i>	-	-	-	-	-	-	-	-	9	1	2	-	3	-
<i>Meriones tamariscinus</i>	-	-	9	-	-	-	-	-	-	-	-	-	-	-
<i>Meriones</i> spec.	84	-	5	-	-	12	-	17	73	16	2	8	2	14
ARVICOLIDAE														
<i>Ondatra zibethicus</i>	1	-	9	-	-	-	-	-	-	-	-	-	-	-
<i>Ellobius tancrei</i>	8	-	24	-	-	-	-	-	1	-	-	-	-	-
<i>Eolagurus przewalskii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	2
<i>Microtus oeconomus</i>	96	1	-	-	-	-	-	-	-	-	-	-	-	-
<i>Microtus</i> spec.	-	1	4	3	7	-	1	-	-	-	-	-	-	-
SCIURIDAE														
<i>Marmota sibirica</i>	-	1	-	-	-	-	-	1	-	-	-	-	-	-
<i>Citellus</i> spec.	-	3	-	-	1	-	-	-	-	-	-	-	-	-
<b>AVES</b>														
<b>REPTILIA</b>														
LACERTIDAE														
	1	-	-	-	-	-	-	-	39	-	-	-	-	1
<b>PISCES</b>														
<b>Arachnida</b>														
SOLIFUGAE														
SCORPIONES														
<b>INSECTA</b>														
	++	-	-	-	-	-	-	-	-	-	-	-	-	-




Fig. 5: 1 - Baruun chajlaastaj, ca. 20 km SE Cogt-Ovoo, 2 – Mandach Sum, 3 - Galbyn-gobi, 4 Bordzongijn-gobi, Ulaan Eregijn Sajr.

Table 2: *Asio otus* (finding locations 1-4)

MAMMALIA	n	%
<b>Insectivora</b>		
<i>Hemiechinus auritus</i>	1	0.1
<i>Crocidura</i> spec.	4	0.5
<b>Lagomorpha</b>		
<i>Lepus tolai</i>	1	0.1
<b>Rodentia</b>		
<b>DIPODIDAE</b>		
<i>Dipus/Allactaga</i> spec.	24	3.2
<i>Salpingotus/Cardiocranus</i>	92	12.2
<b>CRICETIDAE</b>		
<i>Cricetulus/Phodopus</i> spec.	373	49.3
<i>Allocricetulus</i> spec.	6	0.8
<b>GERBILLIDAE</b>		
<i>Rhombomys opimus</i>	13	1.7
<i>Meriones</i> spec.	217	28.7
<b>ARVICOLIDAE</b>		
<i>Eolagurus przewalskii</i>	4	0.5
<b>AVES</b>	12	1.6
<b>REPTILIA</b>	9	1.2
Vertebrata	756	
Insecta	23	



Fig. 6: Fledgling of *Asio otus* in the sajr Baruun chajlaastaj (photo: A. STUBBE).








Fig. 7: Food composition of *Bubo bubo*: **a** - in West-Mongolia (locality 2), **b** - Southwest-Mongolia (localities 8 - 10) and **c** - South-Mongolia (localities 13 - 16); **d** - *Asio otus* (1 - 4).

In total in pellets of *Bubo bubo* were analysed 1 098 vertebrates (1 017 mammals, 39 birds, 41 reptiles, 1 fish), the mammals belonging to 22 species. In the pellets Lagomorpha (mainly *Ochotona pallasi*) were dominant with 23.3 % in SW-Mongolia, Dipodidae were dominating in W-, SW- and S-Mongolia (37.3 %, 23.4 %, 30.9 %), Cricetidae in SW- and S-Mongolia (13.6 %, 17.0 %) and Arvicolidae in W- and SW-Mongolia (23.8 %, 23.9 %). The percentage of birds was between 3.2 and 4.5 %. In N-Mongolia (Minž-gol) one fish was found as prey of *Bubo bubo*. Rests of evertebrates (Arachnida, Solifugae, Scorpiones, Coleoptera, Orthoptera) were noticed in rare cases.

In the pellets of *Asio otus* 756 vertebrates (735 mammals, 12 birds, 9 reptiles) were determined. The diet contained about 13 mammal species. Small Cricetidae (*Cricetus* and *Phodopus*) were dominant with 50.5 %, followed by the families Gerbillidae (*Meriones* spec. - 29 %) and small Dipodidae (*Salpingotus* and *Cardiocranus*) with 12.3 %. In one nest - far from every human housing and activity - was found a recently caught *Mus musculus*.



Fig. 8: *Bubo bubo* fledgling ringed in the sajr Šutegijn Bajan-gol (photo: F.-U. MICHLER, 23.07.2011).

## Discussion

Mammals are the main prey of *Bubo bubo*, but also many bird species could be found in the pellet analyses (PIECHOCKI et al. 1977, STUBBE et al. 1989). Micromammalia as Soricidae, the little jerboas *Cardiocranus* and *Salpingotus*, *Mus musculus* as well as small hamsters of the genus *Phodopus*, *Cricetus* and *Allocricetus* can be analyzed in various percentages in the prey. To the biggest prey belong hares (*Lepus tolai*) and *Ondatra zibethicus*.

Rare prey objects are carnivores and fishes. PIECHOCKI et al. (1977) and STUBBE et al. (1989) mention two juvenile Red foxes (*Vulpes vulpes*) and three *Mustela erminea* for the Čono-charajchijn-gol as prey of *Bubo bubo*.

Also in West-Mongolia, Čonocharajchijn-gol, we found two fishes (*Oreoleuciscus* spec.) and at the river Tuul-gol one *Brachymystax lenok* (locality 6) as prey of the Eagle owl (PIECHOCKI et al. 1977).

Remarkable are a lot of lower jaws of Lacertidae (s. table 1) in the pellets from Šutegijn Bajan-gol. There are living species of the genus *Phrynocephalus* and *Eremias*.

In table 3 and 4 we have summarized our results on feeding ecology of *Bubo bubo* in West- and Southwest-Mongolia between 1974 and 2002 in order to show also the dynamic in the biodiversity, differences from year to year as well as in the different ecosystems.

Table 3: Summarized data (vertebrates) of qualitative pellet analyses from feeding/breeding localities of *Bubo bubo* in West-Mongolia (localities 1 – 2), in brackets - year of collection

species/groups	PIECHOCK I et al. 1977 (1974/75)		STUBBE et al. 1989 (1982)		HOFMANN et al. 2005 (1984/85)		STUBBE et al. 2002 (2002)		total	
	n	%	n	%	n	%	n	%	n	%
<b>MAMMALIA</b>										
<i>Hemiechinus auritus</i>	13	0.9	-	-	2	0.8	1	0.2	16	0.5
Soricidae	20	1.4	6	0.6	2	0.8	1	0.2	29	1.0
Chiroptera	1	-	2	0.2	-	-	5	1.1	8	0.3
<i>Ochotona</i>	85	6.2	256	27.6	15	6.1	35	7.9	391	13.0
<i>Lepus tolai</i>	35	2.5	11	1.2	11	4.4	9	2.0	66	2.2
<i>Dipus/Allactaga</i>	241	17.5	186	20.1	71	28.6	143	32.4	641	21.4
<i>Cardiocranus/Salpingotus</i>	50	3.6	72	7.8	33	13.3	22	5.0	177	5.9
<i>Citellus</i>	-	-	1	0.1	-	-	-	-	1	-
<i>Phodopus/Cricetus</i>	146	10.6	88	9.5	7	2.8	19	4.3	260	8.7
<i>Allocricetus</i>	4	0.3	3	0.3	19	7.7	4	0.9	30	1.0
<i>Mus musculus</i>	-	-	1	0.1	-	-	-	-	1	-
<i>Meriones</i>	54	3.9	72	7.8	11	4.4	84	19.0	221	7.4
<i>Alticola</i>	4	0.3	-	-	-	-	-	-	4	0.1
<i>Eolagurus przewalskii</i>	2	0.1	3	0.3	5	2.0	-	-	10	0.3
<i>Ellobius tancrei</i>	77	5.8	14	1.5	6	2.4	8	1.8	105	3.5
<i>Microtus oeconomus</i>	529	38.3	151	16.3	50	20.2	96	21.7	826	27.6
<i>Ondatra zibethicus</i>	-	-	5	0.5	-	-	1	0.2	6	0.2
Carnivora	3	0.2	2	0.2	-	-	-	-	5	0.2
<b>Mammalia total</b>	<b>1264</b>	<b>91.6</b>	<b>873</b>	<b>94.2</b>	<b>232</b>	<b>93.6</b>	<b>428</b>	<b>96.8</b>	<b>2797</b>	<b>93.3</b>
<b>AVES</b>	<b>114</b>	<b>8.3</b>	<b>54</b>	<b>5.8</b>	<b>15</b>	<b>6.1</b>	<b>14</b>	<b>3.2</b>	<b>197</b>	<b>6.6</b>
<b>REPTILIA</b>	-	-	-	-	1	0.4	-	-	1	-
<b>PISCES</b>	2	0.1	-	-	-	-	-	-	2	0.1
<b>Vertebrates total</b>	<b>1380</b>	<b>100</b>	<b>927</b>	<b>100</b>	<b>248</b>	<b>100</b>	<b>442</b>	<b>100</b>	<b>2997</b>	<b>100</b>

In the period from 1974 until 2002, we collected about 3.000 remains of vertebrates in the pellets of Eagle owls in West-Mongolia in the valleys of Čonocharajchijn- and Chovd-gol. The dominating mammals were species of the genus *Ochotona*, the great Dipodidae *Dipus sagitta* and *Allactaga sibirica* as well as especially *Microtus oeconomus* at the first place with 27.6% (16.3-38.3 %). Table 3 gives a reflection of dynamic processes in the mammalian community. *Ondatra zibethicus* was introduced in the Great Westmongolian Lakes (Char-us-nuur) in 1967 (DAWAA et al. 1977) and was also observed at Čonocharajchijn-gol.

Table 4: Summarized data (vertebrates) of qualitative pellet analyses from feeding/breeding localities of *Bubo bubo* in Southwest-Mongolia; in brackets - year of collection

species/groups	STUBBE et al. 1989 (1980 & 1982)		HOFMANN et al. 2005 (1985 & 1988, 1991/92)		STUBBE et al. 2002 (2002)		total	
	n	%	n	%	n	%	n	%
<b>MAMMALIA</b>								
<i>Hemiechinus auritus</i>	1	0.2	1	0.2	1	0.6	3	0.3
Soricidae	1	0.2	1	0.2	-	-	2	0.2
<i>Ochotona</i>	41	8.2	74	15.3	33	18.9	148	12.8
<i>Lepus tolai</i>	19	3.8	10	2.1	8	4.6	37	3.2
<i>Marmota sibirica</i>	-	-	-	-	1	0.6	1	0.1
<i>Citellus undulatus</i>	8	1.6	-	-	3	1.7	11	0.9
<i>Dipus/Allactaga</i>	57	11.4	44	9.1	37	21.1	138	11.9
<i>Alactagulus/Scirtopoda</i>	13	2.6	37	7.6	-	-	50	4.3
<i>Cardiocranius/Salpingotus</i>	12	2.4	5	1.0	4	2.3	21	1.8
<i>Mus musculus</i>	1	0.2	26	5.5	-	-	27	2.3
<i>Dryomys nitedula</i>	-	-	2	0.4	-	-	2	0.2
<i>Phodopus/Cricetus</i>	88	17.7	117	24.1	24	13.7	229	19.8
<i>Rhombomys opimus</i>	-	-	1	0.2	-	-	1	0.1
<i>Meriones</i>	3	0.6	22	4.5	14	8.0	39	3.4
<i>Eolagurus</i>	3	0.6	55	11.3	-	-	58	5.0
<i>Ellobius tancrei</i>	115	23.1	50	10.3	24	13.7	189	16.3
<i>Alticola</i>	63	12.6	2	0.4	-	-	65	5.6
<i>Microtus</i>	33	6.6	10	2.1	9	5.1	52	4.4
<i>Ondatra zibethicus</i>	-	-	8	1.7	9	5.1	17	1.4
Carnivora	1	0.2	1	0.2	-	-	2	0.2
<b>Mammalia total</b>	<b>459</b>	<b>92.0</b>	<b>466</b>	<b>96.1</b>	<b>167</b>	<b>95.4</b>	<b>1092</b>	<b>94.2</b>
<b>AVES</b>	<b>40</b>	<b>8.0</b>	<b>19</b>	<b>3.9</b>	<b>8</b>	<b>4.6</b>	<b>67</b>	<b>8.8</b>
<b>Vertebrates total</b>	<b>499</b>	<b>100</b>	<b>485</b>	<b>100</b>	<b>175</b>	<b>100</b>	<b>1159</b>	<b>100</b>

In the period from 1980 until 2002 were 1.159 remains of vertebrates collected in the pellets of *Bubo bubo* in Southwest-Mongolia. Dzungarian gobi and Mongolian Altai (locality 7 by STUBBE et al. 1989). At the upper course of Bodončijn-gol therefore we found more remains of *Alticola* and *Citellus*. In total *Ochotona*, the great jerboas *Dipus* and *Allactaga* as well as small hamsters (*Phodopus*, *Cricetus*, *Allocricetus*) and the subterrestrial living *Ellobius tancrei* were the main prey of the Eagle-owl.

Some peculiarities in the prey of Dzungarian region are *Dryomys nitedula* and under the gerbillids *Meriones tamariscinus*. *Ondatra zibethicus* was acclimatized in the Bulgan-gol in 1971 (DAWAA et al. 1977) and is in the mentioned area now a welcome prey object too.

## Literature

DAWAA. N.; STUBBE. M.; DORŽRAA. O. (1977): Die Bisamratte *Ondatra zibethica* (L.. 1758) in der Mongolischen Volksrepublik. – Beitr. Jagd- u. Wildforsch. **10**: 342-352.

GOMBOBAATAR. S.; MONKS. E.M. (comp.) (2011): Mongolian Red List of Birds. - Regional Red List Series Vol. **7**. Birds (Zool. Soc. London). - Ulaanbaatar.

- HOFMANN. Th.; STUBBE. M.; PIECHOCKI. R.; HEIDECKE. D.; SAMJAA. R.; ERFURT. J.; SUM-JAA. D. (2005): Zur Nahrungsökologie des Uhus *Bubo bubo* in der Mongolei. - Erforsch. biol. Ress. Mongolei ( Halle/Saale) **9**: 413-417.
- JAVKHLANTSTSEG. T.; GOMBOBAATAR. S.; YOSEF. R. (2007): Prey selection in Eurasian Eagle-owl (*Bubo bubo*) in Mongolia. - Proceedings of the World Owl Conference. 31 Oct. – 4 Nov. 2007. Groningen. p.72.
- PIECHOCKI. R. (1980): Der Uhu in der Mongolischen Volksrepublik. - Der Falke **27**: 375-380.
- PIECHOCKI. R.; STUBBE. M.; UHLENHAUT. K.; DAWAA. N. (1977): Die Ernährungsökologie des Uhus *Bubo bubo yenisseensis* BUTURLIN in der Mongolischen Volksrepublik. - Zool. Jb. Syst. **104**: 539-559.
- SCOTT. D.M.; GLADWIN. K.; BARTON. N. (2005): Comparison of the diet of Two Desert-living Owls. The Long-eared Owl (*Asio otus*) and Little Owl (*Athene noctua*) from Southern Mongolia. - Mongolian Journal of Biological Science **3** (1): 31-37.
- STUBBE. M.; ANSORGE. H.; PIECHOCKI. R.; LANGE. U.; SAMJAA. R.; BAARTUU. D. (1989): Weitere Beiträge zur Ernährung des Uhus *Bubo bubo* (L.. 1758) in der Westmongolei. - Erforsch. biol. Ress. MVR (Halle/Saale) **6**: 25-32.
- STUBBE. M.; CHOTOLCHU. N. (1968): Zur Säugetierfauna der Mongolei. - Mitt. Zool. Mus. Berlin **44**: 5-121.
- TUMURBAT. J.; GOMBOBAATAR. S.; YOSEF. R. (2009): Nest site and food composition of the Eurasian Eagle-owl *Bubo bubo* in Mongolia. - Ardea **97** (4): 519-523.

### Addresses:

Michael Stubbe\*  
 Annegret Stubbe  
 Oliver Lindecke  
 Martin-Luther-University  
 Halle-Wittenberg  
 Institute of Biology  
 Department of Zoology  
 Domplatz 4  
 Halle/Saale  
 D-06108

E-mail: annegret.stubbe@zoologie.uni-halle.de

Nyamsuren Batsajchan  
 Ravčigijn Samjaa  
 Nationale University of Mongolia  
 School of Arts and Sciences  
 Department of Biology  
 P.O.Box 348  
 Ulaanbaatar 210646  
 Mongolia

\* Corresponding author