

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

UCARE Research Products

UCARE: Undergraduate Creative Activities &  
Research Experiences

---

4-2020

## Mitochondria as a Potential Antifungal Target for Isocyanide Compounds

Nancy Nguyen

*University of Nebraska - Lincoln*

Virendra Tiwari

*University of Nebraska - Lincoln, vtiwari2@unl.edu*

Medhanjali DasGupta

*University of Nebraska - Lincoln, medhanjali.dasgupta@huskers.unl.edu*

David Berkowitz

*University of Nebraska-Lincoln, dberkowitz1@unl.edu*

Mark A. Wilson

*University of Nebraska-Lincoln, mwilson13@unl.edu*

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/ucareresearch>

 Part of the [Microbiology Commons](#)

---

Nguyen, Nancy; Tiwari, Virendra; DasGupta, Medhanjali; Berkowitz, David; Wilson, Mark A.; and Riekhof, Wayne R., "Mitochondria as a Potential Antifungal Target for Isocyanide Compounds" (2020). *UCARE Research Products*. 187.

<https://digitalcommons.unl.edu/ucareresearch/187>

This Poster is brought to you for free and open access by the UCARE: Undergraduate Creative Activities & Research Experiences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in UCARE Research Products by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Nancy Nguyen, Virendra Tiwari, Medhanjali DasGupta, David Berkowitz, Mark A. Wilson, and Wayne R. Riekhof




# Mitochondria as Potential Antifungal Target for Isocyanide Compounds

Nancy Nguyen<sup>2</sup>, Virendra Tiwari<sup>1</sup>, Medhanjali DasGupta<sup>2</sup>, David Berkowitz<sup>1</sup>, Mark Wilson<sup>2</sup>, and Wayne Riekhof<sup>3</sup>  
Departments of Chemistry<sup>1</sup>, Biochemistry<sup>2</sup>, and School of Biological Sciences<sup>3</sup>  
University of Nebraska – Lincoln, Lincoln, NE, 68588 USA



## Background

- Antibacterial and antifungal resistance has created a need for new antimicrobial compounds with different mechanisms of action relative to established drugs.
- Fungi and mammals are both eukaryotes, so it is difficult to find a compound that is effective in fungi and is not toxic to mammals.
- Natural isocyanide such as the fungal-natural product Xanthocillin have antimicrobial properties<sup>1</sup> and are generally non-toxic in mammalian cell culture models.
- Para-nitrophenyl isocyanide (p-NPIC), was synthesized as a “model” isocyanide and tested for its ability to inhibit microbial growth.



1. Lim FY, Won TH, Raffa N, Baccile JA, Wisecaver J, Rokas A, ... Keller NP. Fungal Isocyanide Synthases and Xanthocillin Biosynthesis in *Aspergillus fumigatus*. *mBio*. 2018; 9:e00785-18.  
2. Center for Disease and Control. Countries from which *Candida auris* cases have been reported, as of May 31, 2019. <https://www.cdc.gov/fungal/candida-auris/tracking-c-auris.html#world>. July 18, 2019.

## Xanthocillin and Para-nitrophenyl Isocyanide Structure




Figure 1: The chemical structure of the natural compound, Xanthocillin (left), produced by *Penicillium notatum*. The synthetic compound, para-nitrophenyl isocyanide (p-NPIC) was synthesized by Dr. David Berkowitz from the Department of Chemistry at the University of Nebraska–Lincoln (right).

## Research Questions

- Is para-nitrophenyl isocyanide an effective anti-microbial compound?
- If so, what is the minimum inhibitory concentration of para-nitrophenyl isocyanide to deletion mutants in solid and liquid media?
- What is the molecular mechanism of action of para-nitrophenyl isocyanide?

## Testing p-NPIC on *S. cerevisiae* Mutants on Solid Medium

### Determining MIC on *S. cerevisiae* and *C. albicans* on Solid Medium





Figure 4: Determining the MIC of *Saccharomyces cerevisiae* strain (BY4741) and *Candida albicans* strain (SC5314) on solid YPD media containing various concentrations of p-NPIC. Cultures were serially diluted 5-fold and transferred using a 48 pin multi-spot replicator. Plates were grown at 30°C for 24 hours.

## Genetic Screening on Solid Media Methods




## Hypersensitive Deletion Mutants on Solid Media

Gene Classes	# of Hypersensitive Mutant Strains	List of Affected Genes
Mitochondrial function	60	AIM10, MRPL, RPO41, ...
Vacuolar membrane ATPase	10	VMA4, VMA6, VPH2, ...
Iron-sulfur and copper binding	7	FRE2, GRX5, SCO1, ...
Cytochrome C function	5	COX7, OXA1, QCR2, ...
Ribosomal function	4	RPS23B, RPP2B, BUD21, ...
Translation factors	3	HCR1, TIF3, MRN1
Other	76	

## Testing p-NPIC on *S. cerevisiae* Mutants in Liquid Medium


### p-NPIC Serial Dilution Method



## Hypersensitive Deletion Mutants in Liquid Media

Gene Classes	# of Hypersensitive Mutant Strains	Top 3 Most Affected Genes
Mitochondrial function	21	GCV3, HTD2, OAR1
Vacuolar membrane ATPase	7	VMA6, VMA10, VMA16
Iron-sulfur and copper binding	3	NFU1, GRX5, IBA57
Cytochrome C function	2	QCR2, COX19
Translation factors	1	TIF3
Other	13	HMF1, MOT3, NUP133

## MIC of Hypersensitive Mitochondria Deletion Mutants in Liquid Media



## Conclusions

- Disrupting genes involved in mitochondrial function, iron and copper homeostasis, and the vacuolar membrane ATPase cause hypersensitivity to this compound.
- We hypothesize that p-NPIC and other isocyanides are disrupting metal homeostasis and the function of Complex IV, leading to cell growth arrest.

## Future Questions

- Does p-NPIC disrupt metal homeostasis?
- Is p-NPIC an enhancer of ROS production at Complex IV?
- Do other synthetic isocyanide compounds share the same mechanism of action?

## Acknowledgments

This work was supported by the UNL UCARE Research program and an NIH Jumpstart seed award from the UNL Office of Research.