University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from the Center for Plant
Science Innovation

Plant Science Innovation, Center for

8-12-2004

Disabling surveillance: bacterial type III secretion system effectors that suppress innate immunity

Avelina Espinosa

University of Nebraska – Lincoln, aespinosa@rwu.edu

James R. Alfano

University of Nebraska at Lincoln, jalfano2@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/plantscifacpub>

Part of the [Plant Biology Commons](#), [Plant Breeding and Genetics Commons](#), and the [Plant Pathology Commons](#)

Espinosa, Avelina and Alfano, James R., "Disabling surveillance: bacterial type III secretion system effectors that suppress innate immunity" (2004). *Faculty Publications from the Center for Plant Science Innovation*. 204.

<https://digitalcommons.unl.edu/plantscifacpub/204>

This Article is brought to you for free and open access by the Plant Science Innovation, Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Center for Plant Science Innovation by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Microrreview

Disabling surveillance: bacterial type III secretion system effectors that suppress innate immunity

Avelina Espinosa[†] and James R. Alfano^{*}

Plant Science Initiative and The Department of Plant Pathology, University of Nebraska, Lincoln, NE 68588-0660, USA.

Summary

Many Gram-negative bacterial pathogens of plants and animals are dependent on a type III protein secretion system (TTSS). TTSSs translocate effector proteins into host cells and are capable of modifying signal transduction pathways. The innate immune system of eukaryotes detects the presence of pathogens using specific pathogen recognition receptors (PRRs). Plant PRRs include the FLS2 receptor kinase and resistance proteins. Animal PRRs include Toll-like receptors and nucleotide-binding oligomerization domain proteins. PRRs initiate signal transduction pathways that include mitogen-activated protein kinase (MAPK) cascades that activate defence-related transcription factors. This results in induction of proinflammatory cytokines in animals, and hallmarks of defence in plants including the hypersensitive response, callose deposition and the production of pathogenesis-related proteins. Several type III effectors from animal and plant pathogens have evolved to counteract innate immunity. For example, the *Yersinia* YopJ/P cysteine protease and the *Pseudomonas syringae* HopPtoD2 protein tyrosine phosphatase inhibits defence-related MAPK kinase activity in animals and plants respectively. Thus, type III effectors can suppress signal transduction pathways activated by PRR surveillance systems. Understanding targets and activities of type III effectors will reveal much about bacterial pathogenicity and the innate immune system in plants and animals.

Introduction

A number of Gram-negative bacterial plant and animal pathogens depend on type III secretion systems (TTSSs) to inject virulence effector proteins into host cells (Hueck, 1998; Galán and Collmer, 1999). These pathogens include the animal pathogens *Yersinia* spp., *Salmonella typhimurium*, *Shigella flexneri*, *Escherichia coli* and *Pseudomonas aeruginosa* and the plant pathogens *Pseudomonas syringae*, *Erwinia amylovora*, *Xanthomonas campestris* and *Ralstonia solanacearum*. The diseases they cause run the whole gamut from Black Death in humans (i.e. the bubonic plague) caused by *Yersinia pestis* to fire blight of pear and apple caused by *E. amylovora* (Alfano and Collmer, 1996; Cornelis, 2000). Moreover, TTSSs have also been recently identified in insect pathogens and eukaryotic-associated bacteria (Dale *et al.*, 2001; Marie *et al.*, 2001). Certain animal pathogens have two distinct TTSSs (three if the flagellar TTSS is included), whereas plant pathogens, excluding the flagellar TTSS, appear to have one (Shea *et al.*, 1996; Haller *et al.*, 2000).

The TTSSs and the proteins they secrete have been given a variety of names depending on the species of origin. For example, the well-characterized TTSS from *Yersinia* spp. is called the Ysc system (Yop secretion) and the effector proteins it secretes are Yops (*Yersinia* outer protein). In plant pathogens the TTSS is called the Hrp (hypersensitive response and pathogenicity) system because the original mutants lost the ability to elicit the hypersensitive response (HR) (Lam *et al.*, 2001), a programmed cell death (PCD) associated with plant defence, as well as their pathogenic ability. We now know these original mutants that displayed Hrp phenotypes were defective in components of the Hrp TTSS apparatus (Alfano and Collmer, 1997). The proteins secreted by Hrp TTSSs have been given a variety of names including Hop (for Hrp outer protein), Xop (*Xanthomonas* outer protein), Pop (*Pseudomonas* outer protein, which actually are *R. solanacearum* proteins based on its earlier genus name) and Avr (for avirulence) because several of these were originally identified based on their property of limiting the host range of the

Received 12 July, 2004; revised 11 August, 2004; accepted 12 August, 2004. *For correspondence. E-mail jalfano2@unl.edu; Tel. (+1) 402 4720395; Fax (+1) 402 4723139. [†]Present address: Biology and Marine Biology Department, Roger Williams University, Bristol, RI 02809, USA.

pathogen (Alfano and Collmer, 1997; Bonas and Lahaye, 2002).

The activities of many of the TTSS effectors have been well characterized in animal pathogens where the majority of them co-opt the host cytoskeleton by either inhibiting or promoting actin polymerization to block or induce phagocytosis by host cells (Cornelis, 2002a; Cossart and Sansonetti, 2004). The TTSS effectors accomplish this by modulating the signal transduction pathways controlling actin polymerization in host cells. There are excellent reviews describing the activities of animal type III effectors (Cornelis, 2002b; Holden, 2002; Zaharik *et al.*, 2002). Plant pathogen effectors are less understood. However, this outlook has improved with the recent identification of many effectors from plant pathogens (mostly from *P. syringae*), providing additional insights into their potential activities (Collmer *et al.*, 2002; Guttman *et al.*, 2002; Petnicki-Ocwieja *et al.*, 2002; Greenberg and Vinatzer, 2003; Chang *et al.*, 2004). The large increase in the effector inventory has been spurred by genomic investigations and similar studies on other TTSS-containing plant pathogens suggest that the effector list for these microbes will be considerable as well (Salanoubat *et al.*, 2002; da Silva *et al.*, 2002; Buttner *et al.*, 2003). What are these effectors doing inside the plant cell to favour pathogenesis? As we describe below several *P. syringae* effectors have been shown to suppress plant defence responses. Interestingly, several effectors from animal pathogens also suppress the innate immune system in mammals. Thus, an emerging role of type III effectors appears to be suppression of eukaryotic innate immunity.

One of the primary goals of this review is to highlight the plant pathogen effectors that suppress plant defences and compare them with animal pathogen effectors that act in similar ways. Plant defences have been categorized differently based on how the resistance is manifested and a brief introduction to some of these terms is warranted. Induced defences that successfully stop non-pathogenic bacteria (along with premade defences, which are not discussed here) from growing in the intercellular spaces of plants have been referred to as basal defences, localized induced resistance or innate immunity (Nurnberger *et al.*, 2004). In addition, plants can protect themselves against pathogens that cause diseases in other plant species, referred to as non-host resistance (Heath, 2000; Thordal-Christensen, 2003). Finally, specific cultivars or lines of a given plant species can be resistant to specific races or strains of a pathogenic microorganism. These defences are referred to as cultivar-specific or gene-for-gene resistance because they result from the presence of Avr signals in the pathogen, which are recognized by resistance (R) proteins in the host plant (Flor, 1971; Keen, 1990). As described below, most of the bacterial Avr signals are type III effectors, which are recognized inside the

plant cell by R protein-mediated defence pathways. Here, we will use the terms basal resistance, non-host resistance and cultivar-specific resistance to refer to the different types of plant resistances. Collectively, we will refer to them as the plant innate immune system because they seem to be different layers of the same defence system and have similarities to innate immunity in animals.

The innate immune systems of plants, mammals and insects are dependent on a hardwired sensor-based surveillance system that recognizes indicators of infection by microorganisms (Imler and Hoffmann, 2001; Nurnberger *et al.*, 2004). Thus, to succeed as pathogens bacteria must first disable these defences before they gain access to their biochemical loot – the nutrients available in the host. This review will focus primarily on the recent findings that indicate plant pathogens use TTSS effectors to circumvent plant innate immunity. Because TTSS effectors from animal pathogens are also capable of suppressing innate immunity, we will compare these effectors with their plant pathogen counterparts. First, we introduce some of the salient features of the innate immune systems in both plants and animals that constitute the 'alarm' sensors that trigger host defence responses against bacteria.

Tripping the switch: pathogen-associated molecular patterns (PAMPs), general elicitors and Avr proteins activate innate immunity

Plants induce basal defence responses upon sensing conserved molecules produced by microorganisms (Boller, 1995). These include chitin (Felix *et al.*, 1993) and ergosterol (Granado *et al.*, 1995) from fungi and lipopolysaccharide (LPS) (Dow *et al.*, 2000), cold shock protein (Felix and Boller, 2003), and flagellin from bacteria (Gomez-Gomez and Boller, 2002). These molecules have been referred to as general elicitors, but they are conceptually equivalent to pathogen-associated molecular patterns (PAMPs), molecules present in microbes that are recognized by the innate immune system in animals (Boller, 1995; Medzhitov and Janeway, 1997). PAMPs are recognized in animals by specific host pattern recognition receptors (PRRs), which include the Toll-like receptors (TLRs) and nucleotide binding oligomerization domain (NOD) proteins (Medzhitov, 2001; Inohara and Nunez, 2003). TLRs generally sense extracellular PAMPs, whereas NODs recognize PAMPs intracellularly (Girardin *et al.*, 2002).

A similar dichotomy is seen with plant recognition proteins. A plant receptor-like kinase (RLK), FLS2, recognizes bacterial flagellin from the outside of plant cells, whereas R proteins recognize bacterial Avr proteins intracellularly. The similarities between these proteins and animal TLRs and NODs have been described (Dangl and Jones, 2001; Medzhitov, 2001; Nurnberger and Brunner,

2002; Inohara and Nunez, 2003; Jones and Takemoto, 2004; Nurnberger *et al.*, 2004) and here, we group plant proteins that sense either PAMPs or Avr proteins together with animal proteins that sense PAMPs as PRRs (Fig. 1). FLS2 is a plant PRR that recognizes flagellin PAMP (Gomez-Gomez and Boller, 2000). Bacterial molecules other than flagellin induce similar basal defences, suggesting that other plant PRRs might recognize additional bacterial PAMPs. Moreover, bacterial extracts without elicitor-active flagellin induce basal resistance in *Arabidopsis* implying that plants can recognize other bacterial PAMPs (Zipfel *et al.*, 2004). *Arabidopsis* possesses 173 RLKs that contain predicted extracellular domains and 10 candidate RLKs are grouped phylogenetically with FLS2 (Shiu and Bleecker, 2003). Thus, at least a subset of these may encode PRRs. However, RLKs also play a prominent role in plant development and hormone perception (Torii, 2000).

FLS2 has a predicted extracellular leucine-rich repeat (LRR) domain which is probably involved in protein–

protein interactions (Kobe and Kajava, 2001) (see Fig. 1). LRR domains are present in plant R proteins as well as in Toll, TLRs and NOD proteins from insects and mammals. Another key domain of FLS2 is a Ser/Thr kinase domain, which transduces the PAMP signal to the cytoplasm (Gomez-Gomez *et al.*, 2001). Similar to FLS2, the rice R protein Xa21 possesses an extracellular LRR domain and a cytoplasmic Ser/Thr kinase domain (Fig. 1) (Song *et al.*, 1995). Xa21 confers resistance from the bacterial plant pathogen *Xanthomonas oryzae* pv. *oryzae* race 6. This is interesting because known bacterial Avr proteins are recognized inside plant cells by cytoplasmic R proteins. While the Avr determinant, AvrXa21, which is recognized by Xa21 has yet to be identified, *X. o. oryzae* mutants lacking AvrXa21 activity are defective in sulphur assimilation hinting that Xa21 recognizes a sulphated molecule (Shen *et al.*, 2002). Moreover, *X. o. oryzae* genes required for AvrXa21 avirulence are similar to *Sinorhizobium meliloti* nodulation genes needed for sulphate decoration of carbohydrate Nod factors. Legume RLKs involved in recognition of rhizobia Nod factors have been identified (Limpens *et al.*, 2003; Madsen *et al.*, 2003; Radutoiu *et al.*, 2003) and they possess a cytoplasmic kinase domain similar to FLS2 and Xa21 (Fig. 1). However, instead of LRR domains they have LysM domains, which are thought to interact with carbohydrates (Spaink, 2004). The similarities between FLS2 and Xa21 suggest AvrXa21 may be more similar to a PAMP than an Avr protein, which blurs the distinction between what constitutes a PAMP and an Avr.

Plant R proteins that perceive bacterial Avr proteins can be categorized into four different classes (Fig. 1). The largest, referred to as CC-NBS-LRR R proteins, have a N-terminal coiled-coil (CC) domain, a central nucleotide binding site (NBS) domain (equivalent to NOD domains in animal PRRs) and a C-terminal LRR domain (Dangl and Jones, 2001; Jones and Takemoto, 2004). A second class, TIR-NBS-LRR, possesses a different N-terminal domain, TIR (for Toll and IL-1 receptor) instead of a CC domain. Only one member of this class is known to recognize a bacterial Avr, AvrRps4 (Gassmann *et al.*, 1999), and several other members recognize fungal and viral Avr proteins (Nimchuk *et al.*, 2003). A third type of NBS-LRR R protein, RRS1, is a TIR-NBS-LRR R protein that has a C-terminal tryptophan-arginine-lysine-tyrosine (WRKY) domain. WRKY domains are predicted to act as transcription factors (Eulgem *et al.*, 2000; Deslandes *et al.*, 2002). Another R protein is the tomato Pto protein, which is a Ser/Thr protein kinase required for the recognition of the *P. syringae* Avr proteins AvrPto and AvrPtoB (Pedley and Martin, 2003). Pto is not an NBS-LRR type R protein. However, Pto is dependent on Prf, a CC-NBS-LRR type R protein, for recognition of Avr proteins. Thus, Pto is not a 'classic' PRR like FLS2 and NBS-LRR R

Fig. 1. Representative mammalian and plant pattern recognition receptors that recognize bacterial molecules. In mammals, TLR2 recognizes several different microbial PAMPs including diacyl and triacyl lipopeptides with the assistance of TLR1 and TLR6, respectively, which are not shown (Takeda and Akira, 2003). TLR4 recognizes the lipopolysaccharide component of the outer membrane of Gram-negative bacteria in conjunction with CD14 (not shown). TLR5 recognizes bacterial flagellin and TLR9 recognizes bacterial CpG DNA. NOD1 and NOD2 both recognize components of peptidoglycan. NOD1 recognizes *N*-acetylglucosamine-*N*-acetylmuramic acid (G-M) with a tripeptide containing mesodiaminopimelate as the terminal amino acid, which is found mainly in Gram-negative bacteria (Chamaillard *et al.*, 2003). NOD2 recognizes G-M dipeptide, which is found in all bacteria containing peptidoglycan. In plants, FLS2 recognizes flagellin, receptor kinases such as NFR1 recognize carbohydrate NOD factors from rhizobia, and the Xa21 receptor kinase recognizes AvrXa21, an undefined Avr signal. Representative R proteins that recognize different Avr proteins are shown (see text for details). LPS, lipopolysaccharide; LRR, leucine-rich repeats; TIR, toll and IL-receptor; PK, serine/threonine protein kinase; LysM, lysin motif; NOD/NBS, nucleotide-binding oligomerization domain/nucleotide-binding site; CARD, amino terminal caspase recruitment effector domain; CC, coiled-coil; WRKY, tryptophan-arginine-lysine-tyrosine domain. This figure was modified from Jones and Takemoto (2004) and Takeda and Akira (2003).

proteins. Thus far, all the R proteins that recognize identified bacterial Avr proteins are located inside the plant cell with several of them localized to the plant plasma membrane (Boyes *et al.*, 1998; Nimchuk *et al.*, 2001). RRS1 is unusual because it becomes nuclear-localized only in the presence of its cognate Avr protein, PopP2, which contains nuclear localization signals (Deslandes *et al.*, 2003). The long standing model for how R proteins recognize their cognate Avr proteins is by directly interacting with them (Gabriel and Rolfe, 1990). However, recent evidence, at least for specific bacterial Avr proteins, supports another model called the guard hypothesis, in which R proteins sense an Avr protein indirectly by detecting its enzymatic activity (Schneider, 2002; Innes, 2004). Because the activities of the majority of plant pathogen effectors are unknown, it remains possible that some are not enzymes. Thus, some bacterial Avr proteins may still be recognized directly by their cognate R proteins.

Letting out the dogs: outputs of innate immunity

Signal transduction pathways are triggered once animal or plant PRRs perceive PAMPs or Avr proteins and defence-related outputs are deployed. The signal transduction components acting downstream of Avr protein recognition have been recently reviewed (Martin *et al.*, 2003; Nimchuk *et al.*, 2003). Many of these are dependent on salicylic acid (SA) and are also important in systemic acquired resistance (SAR) (Durrant and Dong, 2004), a form of resistance that is triggered in plants away from the local site of infection. The emerging picture is that a complex signal network is triggered upon perception of an Avr protein. One distinction that can be made is that CC-NBS-LRR R proteins are dependent on NDR1 (Century *et al.*, 1997), whereas TIR-NBS-LRR R proteins are dependent on EDS1 (Aarts *et al.*, 1998; Falk *et al.*, 1999).

Another plant protein known to play a prominent role in R protein-mediated defence pathways is NPR1/NIM1, which was first identified for its role in SAR (Cao *et al.*, 1994; Delaney *et al.*, 1995). NPR1 interacts with several members of the TGA family of basic leucine zipper transcription factors, is nuclear-localized when activated and is a positive regulator of defence genes (Durrant and Dong, 2004). NPR1 shares some similarity with I κ B, a negative regulator of the NF- κ B transcription factor utilized in signal transduction pathways in animals (Ryals *et al.*, 1997). The key role that NPR1 plays in local resistance is highlighted by the fact that overexpression of NPR1 in *Arabidopsis* enhances resistance to *P. syringae*, the oomycete *Peronospora parasitica*, and fungus *Erysiphe cichoracearum* (Friedrich *et al.* 2001; Cao *et al.*, 1998) and in rice conferred enhanced resistance to *X. o. oryzae* (Chern *et al.*, 2001). As NPR1 activation requires SA, this

emphasizes the importance of SA-dependent defences at infection sites.

Components of signal transduction pathways operating downstream of PAMP perception in both plants and animals are not well understood. However, there are common features shared by both, which have been noted (Cohn *et al.*, 2001; Nurnberger and Brunner, 2002; Holt *et al.*, 2003; Nurnberger *et al.*, 2004). For example, mitogen-activated protein kinase (MAPK) cascades are activated in plants and animals after PAMP perception (Nurnberger *et al.*, 2004). In both hosts, transcription factors are activated after activation of MAPK – the NF- κ B transcription factor in animals and WRKY transcription factors in plants are good examples (Lee *et al.*, 1997; Asai *et al.*, 2002; Kim and Zhang, 2004; Zipfel *et al.*, 2004). In plants, MAPK pathways appear to be points of convergence between multiple elicitors of plant defence including PAMPs and Avr proteins. For example in tobacco, the MAPKs SIPK (salicylate-inducible protein kinase) and WIPK (wound-inducible protein kinase) are activated by fungal, viral and bacterial elicitors (Zhang and Klessig, 2001). Silencing of a tobacco MAPK kinase kinase (MAPKKK), NPK1, stops the plant from responding to several Avr proteins, including the *X. c. vesicatoria* Avr protein AvrBs2, consistent with a requirement of MAPK cascades in these R protein-mediated responses (Jin *et al.*, 2002). A MAPKKK that acts upstream of SIPK in *Nicotiana benthamiana* and tomato, MAPKKK α , has recently been identified. This MAPKKK is triggered by different elicitors including AvrPto (del Pozo *et al.*, 2004). A complete MAPK cascade [MKK1 (MAPKKK), MKK4/5 (MAPK kinases) and MPK3/6 (MAPKs)] has been identified in *Arabidopsis*. Among these, MPK6, a SIPK MAPK orthologue, is activated by the flagellin PAMP (Asai *et al.*, 2002; Nuhse *et al.*, 2000). Interestingly, this MAPK cascade results in the activation of several WRKY transcription factors and at least one of these factors protects *Arabidopsis* from fungal and bacterial infections (Asai *et al.*, 2002; Navarro *et al.*, 2004; Zipfel *et al.*, 2004).

There is much overlap between the outputs of both PAMP-triggered and Avr-triggered signal transduction pathways in plants. Some of these physiological responses include generation of reactive oxygen species (ROS) and nitric oxide (NO), changes in cytoplasmic Ca²⁺ levels, production of pathogen-related gene expression, phytoalexin production and callose deposition in the cell wall (Heath, 2000; Dangl and Jones, 2001; Gomez-Gomez and Boller, 2002; Jones and Takemoto, 2004). Among the more dramatic defence responses triggered is the aforementioned HR, more frequently associated with non-host and cultivar-specific resistance than with basal resistance. Even though we know many of the outputs of a successful defence response, it remains unclear which outputs are effective against bacteria. The failure of

PAMPs to trigger an HR may be more of a threshold effect than differences in signal transduction as most of the other responses are shared between cultivar-specific and basal resistances. Supporting this, the non-specific oxidative burst triggered by both saprophytes and pathogens is less substantial than the one that is specifically induced by the presence of Avr proteins (Baker and Orlandi, 1995).

Clearly, a better understanding of the differences in R protein pathways and PAMP-triggered pathways is needed. Currently, one way to separate plant defence outputs is to test whether the output in question is dependent on SA. This can be easily done by determining whether the response is produced in *Arabidopsis* mutant plants impaired in SA biosynthesis (Wildermuth *et al.*, 2001). Alternatively, *nahG* plants could be utilized to test whether a plant response is dependent on SA. These plants carry the *P. putida nahG* gene that encodes salicylate hydroxylase (Gaffney *et al.*, 1993), an enzyme that degrades SA, which is generally required for R protein-dependent defences (Dangl and Jones, 2001; Bonas and Lahaye, 2002). Nevertheless, experiments using *nahG* plants should be interpreted with caution because a recent report suggests that the loss of non-host resistance to *P. syringae* by *nahG*-containing *Arabidopsis* plants results from SA degradation products and not from the inhibition of SA-dependent defences (van Wees and Glazebrook, 2003).

In animals, PAMPs are recognized by TLRs and NOD PRRs, which activate signalling pathways that induce innate proinflammatory responses (Barton and Medzhitov, 2003; Inohara and Nunez, 2003). Several of these responses, once over a threshold, will activate the adaptive immune system in animals. Identifying signal transduction components downstream of animal PRRs is an active area of research (Barton and Medzhitov, 2004). All known TLRs utilize MyD88 TIR-domain adaptor proteins. Recently, certain TLRs have been shown to utilize other TIR-domain-containing adaptor proteins called Trif (or TICAM-1) (Yamamoto *et al.*, 2003) and TIRAP (Horng *et al.*, 2002). MyD88 recruits members of the IL-1 receptor-associated kinases IRAK-1 and IRAK-4 (Takeda *et al.*, 2003). The plant R protein Pto shares some similarities with IRAKs and Pelle, a kinase involved in innate immunity in *Drosophila* highlighting another similarity in plant and animal innate immunity (Cohn *et al.*, 2001). After IRAK is activated by phosphorylation, it associates with TRAF6, an E3 ligase, which in turn activates MAP kinases. This kinase cascade results in the activation of both the Ap-1 and NF- κ B transcription factors, which activate a large number of proinflammatory genes (Pasare and Medzhitov, 2003; Takeda *et al.*, 2003).

Signalling pathways acting downstream of NOD PRRs are less understood than TLRs probably because they have only recently been associated with PAMP percep-

tion. However, activation of both NOD1 and NOD2 results in NF- κ B and caspase activation (Inohara and Nunez, 2003). In addition, both also associate with RICK (also known as RIP2), a CARD-containing protein kinase, which in turn associates with NEMO, a regulatory subunit of the IKK complex (Carneiro *et al.*, 2004). Ultimately, triggering NOD1 and NOD2 results in the secretion of cytokines (e.g. IL-8) from antigen-presenting cells and the expression of other costimulatory molecules that activate antigen-specific effector T cells of the adaptive immune system (Inohara and Nunez, 2003).

A myriad of responses occur after triggering either animal or plant PRRs. The identification of these responses has been greatly facilitated by mRNA expression profiling after exposure to a PAMP or pathogen. For example, Tao *et al.* (2003) profiled the changes in mRNA expression in *Arabidopsis* infected with a virulent *P. syringae* strain versus *P. syringae* strains containing Avr proteins recognized by R protein PRRs present in the test plant. The presence of specific Avr proteins in *P. syringae* (i.e. making *P. syringae* avirulent) produced a similar mRNA pattern as the virulent pathogen except that the expression of defence genes was earlier and increased. This supports the hypothesis that bacterial plant pathogens have strategies to successfully delay and/or suppress defence responses. Recent gene profiling experiments found that many genes are activated upon challenging *Arabidopsis* plants with flagellin (Navarro *et al.*, 2004; Zipfel *et al.*, 2004). These experiments revealed important observations and new directions for experimentation. One particularly interesting result is that the recognition of flagellin by FLS2 induces the expression of many other RLKs suggesting that recognition of the flagellin PAMP potentiates the plant to recognize other PAMPs. Studies such as these should continue to reveal the global changes that occur upon PAMP recognition.

Disabling surveillance: type III effectors that act as suppressors of innate immunity

Given the elaborate surveillance strategies plants and animals use to defend against microorganisms, it is not surprising that microorganisms have countered with their own strategies to thwart their hosts. It has become increasingly clear within the last couple of years that plant pathogen type III effectors suppress plant defence responses (Alfano and Collmer, 2004). Indeed, there were clues in the early 1990s that the TTSSs of plant pathogens were capable of suppressing plant defences. For example, *X. c. pv. campestris hrp* mutants induced an HR-like response in the vascular system of crucifers (called the vascular HR), which the compatible wild-type strain did not elicit, suggesting that wild-type strains suppressed the vascular HR (Kamoun *et al.*, 1992). In pioneering

research, Jakobek *et al.* (1993) showed that a compatible *P. syringae* pathovar, *P. s. pv. phaseolicola*, suppressed induction of defence-related mRNA and phytoalexins in bean that were separately induced by an incompatible *P. syringae* pathovar and non-pathogenic *E. coli*. In retrospect, the induction of defence responses by *E. coli* probably resulted from PAMP-triggered defences. Another study, used electron microscopy to show that *X. c. pv. vesicatoria* suppressed papillae formation in plants and this ability was dependent on *hrp* genes (Brown *et al.*, 1995).

These early studies suggested the TTSS of bacterial plant pathogens was involved in defence suppression. More recent studies suggested that specific type III effectors may be altering defence pathways in plants. For example, VirPphA, a type III effector from *P. s. pv. phaseolicola*, blocked other 'masked' Avr proteins from eliciting the HR (Jackson *et al.*, 1999). *P. s. pv. phaseolicola virPphA* mutants elicited HRs on certain bean cultivars that were normally infected by the wild-type strain. Thus, the *virPphA* mutation effectively converted a virulent pathogen to an avirulent one. Similar phenotypes were observed for *P. s. pv. phaseolicola* mutants defective in AvrPphC and AvrPphF (Tsiamis *et al.*, 2000). Another phenomenon that now makes more sense in the context of effectors acting as suppressors is the observation that AvrRpt2 blocked the ability of AvrRpm1 to elicit an RPM1-dependent HR in *Arabidopsis* plants (Ritter and Dangl, 1996). AvrRpm1 elicits an HR in *Arabidopsis* more quickly than AvrRpt2. However, when both Avr proteins are present in *P. syringae* and infiltrated into *Arabidopsis* Col-0, which contains R proteins that recognize both Avr, only the slower AvrRpt2-dependent HR develops. This suggested that the AvrRpt2 somehow interfered with AvrRpm1 recognition. This effect appears to result from the AvrRpt2-dependent elimination of a plant protein called RIN4, which is monitored by RPM1 such that when RIN4 is eliminated the RPM1 surveillance system is disabled (Mackey *et al.*, 2002; 2003; Axtell and Staskawicz, 2003).

Recently, several *P. syringae* effectors have been identified as suppressors of the HR and other plant responses associated with defence. The HR elicited by several different Avr proteins were suppressed by the *P. syringae* effectors AvrPphE_{Pto}, AvrPpiB1_{Pto}, AvrPtoB, AvrRpt2, HopPtoD2, HopPtoE, HopPtoF (an AvrPphF homologue) and HopPtoN (Lopez-Solanilla *et al.*, 2004; Abramovitch *et al.*, 2003; Axtell and Staskawicz, 2003; Bretz *et al.*, 2003; Espinosa *et al.*, 2003; Mackey *et al.*, 2003; Jamir *et al.*, 2004). A subset of these also has been shown to suppress other hallmarks of plant defence (Bretz *et al.*, 2003; Chen *et al.*, 2004; Jamir *et al.*, 2004). Moreover, AvrPphE_{Pto}, HopPtoE, AvrPtoB, HopPtoF and HopPtoG suppressed Bax-induced PCD in yeast and plants, and

induction of the PR transcript *PR1a* (Jamir *et al.*, 2004). AvrPtoB also suppressed the Bax-dependent PCD in plants and as well as stress-induced PCD in yeast (Abramovitch *et al.*, 2003). While the mechanism of suppression is unknown, suppression of PCD pathways in yeast suggests that these effectors may act on conserved pathways in eukaryotes.

Some type III effectors are able to suppress basal defences triggered by PAMPs. For example, Hauck *et al.* (2003) found that transgenically expressed AvrPto in *Ara-bidopsis* suppressed the expression of a set of genes predicted to encode proteins that are secreted cell wall and defence proteins, which are normally expressed in a manner independent of SA. Moreover, when *P. syringae* TTSS-defective mutants were infected into AvrPto- or AvrRpt2-expressing plants these mutants grew to significantly higher levels than control strains suggesting that plant defences induced by these mutants were suppressed by both AvrRpt2 and AvrPto (Hauck *et al.*, 2003; Chen *et al.*, 2004). Recently, DebRoy *et al.* (2004) have shown that the *P. syringae* effectors AvrE and HopPtoM suppress SA-dependent basal defences. Characterization of other type III effectors to determine whether they suppress SA-dependent or -independent defence responses (or both) will help identify the plant targets of these effectors. A listing of plant pathogen type III effectors that appear to have a role in suppression of innate immunity has been recently published (Alfano and Collmer, 2004).

Several caveats should be noted regarding effectors that possess suppressor activity: many of the experiments were carried out in heterologous systems (e.g. yeast); or in conditions where the effector displaying the suppressing activity was overexpressed; and, finally, many of the mutants defective in effectors that suppress the HR do not obviously alter their host range (although, in many cases this was not tested), which would be predicted if individual suppressors were required for growth on a specific host plant. Another paradox associated with several of the effectors that suppress defence response is that they can also form necrotic lesions in compatible hosts (Badel *et al.*, 2003; DebRoy *et al.*, 2004; Lopez-Solanilla *et al.*, 2004). These lesions may represent a late HR resulting from latent recognition of these effectors by the plant's surveillance system. A recent report demonstrates that both the HR and necrotic lesions caused by *P. syringae* require a MAPK pathway suggesting that they may be related responses (del Pozo *et al.*, 2004).

There has been substantial progress on the identification of enzymatic activities of plant pathogen type III effectors (Innes, 2003; Alfano and Collmer, 2004; Chang *et al.*, 2004). A growing number of these are active or predicted cysteine proteases (Shao *et al.*, 2002; Axtell *et al.*, 2003; Hotson *et al.*, 2003; Lopez-Solanilla *et al.*, 2004). While the roles that many of these cysteine proteases play in

virulence are unknown, there is evidence that one role for this group of effectors is defence suppression. For example, the *P. s. pv. phaseolicola* AvrPphB cysteine protease targets the *Arabidopsis* PBS1 kinase, a protein involved in R protein-mediated defences (Shao *et al.*, 2003). The *P. s. tomato* HopPtoN effector, a cysteine protease belonging to the same clade as AvrPphB, suppresses an Avr-triggered HR (Lopez-Solanilla *et al.*, 2004).

One plant pathogen type III effector that is particularly noteworthy is HopPtoD2 from *P. s. pv. tomato* DC3000. When HopPtoD2 is expressed in *P. s. pv. phaseolicola*, it suppresses the non-host HR, PR1 gene expression and the oxidative burst (Bretz *et al.*, 2003; Espinosa *et al.*, 2003). Importantly, HopPtoD2 was also demonstrated to possess protein tyrosine phosphatase (PTP) activity. We are not aware of any reports of tyrosine phosphorylated proteins in plants other than the MAPKs (Zhang and Klessig, 2001), which are phosphorylated at both threonine and tyrosine. As noted above, MAPK pathways play important roles in signal transduction pathways controlling plant defence (Zhang and Klessig, 2001). An HR-like response is produced when a constitutively active tobacco MAPKK (NtMEK2^{DD}) is transiently expressed in plants (Yang *et al.*, 2001). Interestingly, when HopPtoD2 is co-expressed in plants with NtMEK2^{DD} the HR-like response is suppressed suggesting that the host cell target(s) of HopPtoD2 is downstream of this MAPKK (Espinosa *et al.*, 2003). This MAPK pathway is activated by a number of Avr proteins and the *Arabidopsis* MAPK homologues AtMPK3/AtMPK6 are also involved in the response to the PAMP flagellin (Zhang and Klessig, 2001; Asai *et al.*, 2002). Thus, it is plausible that HopPtoD2 acts at a point of convergence of signal transduction pathways that are utilized by both Avr-triggered and PAMP-triggered defence responses. Figure 2 shows possible sites of action in the plant defence signal transduction pathways for type III effectors that suppress innate immunity.

Animal pathogen type III effectors have also been shown to suppress innate immunity. Two type III effectors, SptP and YopH, from *S. typhimurium* and *Yersinia* spp., respectively, are also PTPs similar to HopPtoD2 (Guan and Dixon, 1990; Kaniga *et al.*, 1996). Both of these effectors, like many animal pathogen type III effectors, suppress actin polymerization, which inhibits phagocytosis and prevents bacterial uptake by animal cells (Andersson *et al.*, 1996; Fu and Galán, 1998). YopH is also known to downregulate the proinflammatory response (Boland and Cornelis, 1998; Viboud *et al.*, 2003), which is probably at least partially due to the suppression of the synthesis of the monocyte chemoattractant protein 1, a chemokine involved in macrophage recruitment to the sites of infection (Sauvonnet *et al.*, 2002a). YopH is also responsible for the ability of *Yersinia* to prevent T and B cell activation and therefore inhibiting adaptive immunity as well (Yao

Fig. 2. Potential sites of action of plant pathogen type III effectors in innate immunity signal transduction pathways in plants. Bacterial plant pathogens inject many different type III effectors (i.e. Hops) into plant cells. Some type III effectors (i.e. Avr proteins depicted here as a green object) are recognized by plant R proteins triggering SA-dependent and SA-independent defences. The plant cell is capable of recognizing the flagellin PAMP with the receptor-like kinase (RLK) FLS2. Other PAMPs are probably recognized by other plant RLKs. Type III effectors that act as defence suppressors can act at unique parts of the innate immunity signal transduction pathways (e.g. AvrRpt2 suppression of the AvrRpm1-triggered defences) (1); or others may act at convergence point of different signal transduction pathways (e.g. HopPtoD2 suppression of a MAPK pathway) (2); certain type III effectors such as AvrBs3 are localized to the nucleus and modify eukaryotic transcription (Buttner and Bonas, 2003) (3); type III effectors may also act post-transcriptionally to suppress Avr-triggered (4) and PAMP-triggered outputs (5), both of which result from SA-dependent and SA-independent responses. See text for additional details. This figure was modified from Alfano and Collmer (2004).

et al., 1999). Recently, the Lck kinase, a protein involved in T cell receptor signalling, was shown to be a substrate of YopH (Alonso *et al.*, 2004). This is consistent with suppression of innate and adaptive immunity by preventing T cell activation.

SptP is a multifunctional effector with a GTPase activating protein domain in its N-terminal half and a PTP domain in its C-terminal half (Kaniga *et al.*, 1996). Murli *et al.* (2001) showed that *S. typhimurium* sptP mutants activated Erk MAPK pathways to higher levels suggesting that SptP modulated this MAPK pathway. SptP appears to inhibit the Erk MAPK pathway by interfering with Raf kinase activation (Lin *et al.*, 2003). *Salmonella* is known to trigger proinflammatory cytokines in animal cells through the activation of Erk, Jnk and p38 MAPK pathways (Hobbie *et al.*, 1997). SptP also downregulates secretion of the proinflammatory cytokine IL-8 (Haraga and Miller, 2003), which is consistent with it acting as a suppressor of animal innate immunity. Interestingly, SptP also inhibits the p38 MAPK pathway in *Caenorhabditis elegans* and a *Salmonella* sptP mutant is reduced in its ability to kill *C. elegans* suggesting a role in virulence in this heterologous pathosystem (Tenor *et al.*, 2004). These

results indicate that SptP targets MAPK pathways involved in innate immunity (Kim *et al.*, 2002; Aballay *et al.*, 2003).

The *Yersinia* YopJ/P type III effector also suppresses animal innate immunity. YopJ belongs to a large family of type III effectors that are cysteine proteases that are well distributed in animal and plant pathogens (Orth, 2002). The first clue that YopJ suppressed innate immunity was that *Yersinia* was capable of suppressing the proinflammatory cytokine tumour necrosis factor α (TNF- α) that was induced by several MAPK pathways activated after infection (Ruckdeschel *et al.*, 1997). YopJ/P was shown by several groups to be capable of suppressing both NF- κ B and TNF- α in macrophages (Boland and Cornelis, 1998; Palmer *et al.*, 1998; Ruckdeschel *et al.*, 1998; Schesser *et al.*, 1998). YopJ also blocks several signal transduction pathways that activate the CREB transcription factor (Meijer *et al.*, 2000). This transcription factor is activated by the LPS PAMP indicating that YopJ suppresses PAMP-triggered innate immunity. How does YopJ cause these effects? YopJ blocks members of a superfamily of MAPKs needed to activate MAPK pathways induced in mammalian cells by pathogens, and the I κ B complex, which is required for activation of NF- κ B (Orth *et al.*, 1999). YopJ is an isopeptidase that can use proteins conjugated to small ubiquitin-related modifier (SUMO) as substrates cleaving the isopeptide bond that links SUMO to the target protein (Orth *et al.*, 2000). Cleaving a SUMO group from a protein can alter the protein's activity. However, why YopJ/P's cysteine protease activity is required for MAPK and NF- κ B signalling and how this contributes to its ability to suppress the mammalian immune system is currently unknown.

There are other animal pathogen type III effectors that are capable of suppressing innate immunity. Recently, the *Salmonella* SspH 1 type III effector was shown to localize to the nucleus where it inhibits NF- κ B resulting in down-regulation of IL-8 production (Haraga and Miller, 2003). SspH 1 belongs to a family of type III effectors that contain LRR predicted to be involved in protein-protein interactions. Several of these, including the *Yersinia* YopM and the *Shigella* IpaH9.8, have also been shown to localize the nucleus (Skrzypek *et al.*, 1998; Toyotome *et al.*, 2001). YopM's LRRs are required for localization to the nucleus (Benabdillah *et al.*, 2004). IpaH9.8, like SspH 1, inhibits NF- κ B suggesting that this group of effectors may modify eukaryotic transcription (Haraga and Miller, 2003). Consistent with this idea is the observation that a *Yersinia* *yopM* mutant alters the transcription profile of macrophages compared with the transcription profile of macrophages challenged with wild-type bacteria (Sauvonnet *et al.*, 2002b). Type III effectors that act as eukaryotic defence suppressors do not appear to be limited to pathogens. Recently, NopL, a type III effector from the symbiont

Rhizobium sp. NGR234, was found to suppress the induction of PR genes in tobacco suggesting that symbionts may also benefit from suppressing plant innate immunity (Bartsev *et al.*, 2004).

Concluding remarks

The ability to avoid eukaryotic surveillance systems of the innate immune system was probably a critical development in the evolution of bacterial pathogenicity. In the animal world, this was made doubly complex by an adaptive immune system that could recognize a bewildering array of molecules and specific immune cells (i.e. macrophages) that have the capacity to engulf microorganisms. At first glance, plants seem to be an easier mark because their defence systems rely only on premade defences and a surveillance system that recognizes both conserved molecules on microorganisms (PAMPs) and specific products of pathogens (Avr proteins). However, this surveillance system is surprisingly complex and successful, which may explain why there are relatively few bacterial species that can invade plants. It seems logical to assume that the ability to recognize PAMPs came before Avr recognition in the evolution of plant immunity. Figure 3 shows a possible sequence of events that allowed Gram-negative bacterial pathogens to evolve centred on the horizontal acquisition of a TTSS. In this model, acquisition of the TTSS and type III effectors that disable the PAMP-based innate immune system is the central event that allowed these bacteria to evade plant defences. It seems likely that after pathogens were successful in disabling PAMP-triggered surveillance systems, the plant countered by evolving the ability to recognize Avr proteins of pathogens. To counteract, the pathogen acquired many type III effectors, which are probably functionally redundant, and the plant evolved many PRRs that could recognize as many possible 'looks' the pathogen gives the plant. Understanding plant surveillance systems and type III effectors that disable it is imperative if we are to understand the pathogenicity of plants.

In a larger context, the study of bacterial pathogenicity of plants and animals has benefited from a comparative pathobiology approach. For example, the similarity of NODs to NBS-LRR R proteins probably aided the discovery that NODs were involved in pathogen recognition. Further comparative analyses of the protein complexes involved in PRR perception and their downstream signal transduction pathways will probably increase our understanding of both pathosystems. And many other points of comparison remain. For example, is the YopJ/P-induced apoptosis of macrophages (Mills *et al.*, 1997; Monack *et al.*, 1997) equivalent to the Avr-induced PCD (i.e. HR) in plants? That is, is the PCD of a macrophage a form of defence or a virulence strategy of the pathogen? The

Fig. 3. Scenario for the evolution of bacterial plant pathogens based on suppression of plant innate immunity with type III effectors.

A. A possible scenario for the evolution of Gram-negative bacterial plant pathogenicity predicts a stage in the evolution of plants that allowed non-pathogens to grow in the apoplast unperturbed because plants lacked an innate immune system.

B. Plants that evolved or possessed an innate immune system that allowed for the detection of PAMPs were successful in stopping the growth of non-pathogens.

C. Bacteria that acquired a TTSS through horizontal transfer or evolved a specialized TTSS from the flagellar TTSS and delivered type III effectors that suppressed the PAMP-triggered defence responses were successful in growing in the intercellular spaces of plants.

D. Plants responded to this biotic stress by evolving R proteins that could recognize certain type III effects as Avr protein to trigger a rapid intense defence response which included the HR.

E. The acquisition of additional type III effectors conferred the ability to suppress Avr-triggered defences.

F. This put additional evolutionary pressure on plants to expand their R protein surveillance system to recognize the activity of other type III effectors.

The scenario predicts that the first targets of type III effectors were plant components that were involved in the PAMP-triggered defences (i.e. basal defences). Continued co-evolution of plant PRRs and type III effector suppressors resulted in a highly polymorphic collection of effectors and plant PRRs.

CW, plant cell wall; PM, plant plasma membrane.

Yersinia YopB type III translocator does trigger proinflammatory responses in host cells similar to PAMPs (Viboud *et al.*, 2003), and potentially analogous to Avr proteins. Plant and animal bacterial pathogens now have groups of type III effectors that have similar activities (e.g. cysteine

proteases and tyrosine phosphatases) and a subset of defence suppressors. Thus, comparing the activities and targets of type III suppressors in plant and animal pathosystems should lead to a greater understanding of the innate immune system in eukaryotes.

Acknowledgements

We thank Tanja Petnicki-Ocwieja, Zhenq Qing Fu and Byeong-ryool Jeong for critical reading of the manuscript. Work in the Alfano laboratory is currently supported by NSF Plant Genome Research Program Cooperative Agreement DBI-0077622, NSF Grant MCB-0317165 and USDA/NRI Grant 03-35319-13862.

References

- Aarts, N., Metz, M., Holub, E., Staskawicz, B.J., Daniels, M.J., and Parker, J.E. (1998) Different requirements for EDS1 and NDR1 by disease resistance genes define at least two R gene-mediated signalling pathways in *Arabidopsis*. *Proc Natl Acad Sci USA* **95**: 10306–10311.
- Aballay, A., Drenkard, E., Hilbun, L.R., and Ausubel, F.M. (2003) *Caenorhabditis elegans* innate immune response triggered by *Salmonella enterica* requires intact LPS and is mediated by a MAPK signalling pathway. *Curr Biol* **13**: 47–52.
- Abramovitch, R.B., Kim, Y.J., Chen, S., Dickman, M.B., and Martin, G.B. (2003) *Pseudomonas* type III effector AvrPtoB induces plant disease susceptibility by inhibition of host programmed cell death. *EMBO J* **22**: 60–69.
- Alfano, J.R., and Collmer, A. (1996) Bacterial pathogens in plants: life up against the wall. *Plant Cell* **8**: 1683–1698.
- Alfano, J.R., and Collmer, A. (1997) The type III (Hrp) secretion pathway of plant pathogenic bacteria: trafficking harpins, Avr proteins, and death. *J Bacteriol* **179**: 5655–5662.
- Alfano, J.R., and Collmer, A. (2004) Type III secretion system effector proteins: double agents in bacterial disease and plant defense. *Annu Rev Phytopathol* **42**: 385–414.
- Alonso, A., Bottini, N., Bruckner, S., Rahmouni, S., Williams, S., Schoenberger, S.P., and Mustelin, T. (2004) Lck dephosphorylation at Tyr-394 and inhibition of T cell antigen receptor signaling by *Yersinia* phosphatase YopH. *J Biol Chem* **279**: 4922–4928.
- Andersson, K., Carballeira, N., Magnusson, K.-E., Persson, C., Stendahl, O., Wolf-Watz, H., and Fallman, M. (1996) YopH of *Yersinia pseudotuberculosis* interrupts early phosphotyrosine signalling associated with phagocytosis. *Mol Microbiol* **20**: 1057–1069.
- Asai, T., Tena, G., Plotnikova, J., Willmann, M.R., Chiu, W.-L., Gomez-Gomez, L., *et al.* (2002) MAP kinase signalling cascade in *Arabidopsis* innate immunity. *Nature* **415**: 977–983.
- Axtell, M.J., and Staskawicz, B.J. (2003) Initiation of RPS2-specified disease resistance in *Arabidopsis* is coupled to the AvrRpt2-directed elimination of RIN4. *Cell* **112**: 369–377.
- Axtell, M.J., Chisholm, S.T., Dahlbeck, D., and Staskawicz, B.J. (2003) Genetic and molecular evidence that the *Pseudomonas syringae* type III effector protein AvrRpt2 is a cysteine protease. *Mol Microbiol* **49**: 1537–1546.
- Badel, J.L., Nomura, K., Bandyopadhyay, S., Shimizu, R., Collmer, A., and He, S.Y. (2003) *Pseudomonas syringae*

- pv. *tomato* DC3000 HopPtoM (CEL ORF3) is important for lesion formation but not growth in tomato and is secreted and translocated by the Hrp type III secretion system in a chaperone-dependent manner. *Mol Microbiol* **49**: 1239–1251.
- Baker, C.J., and Orlandi, E.W. (1995) Active oxygen in plant pathogenesis. *Annu Rev Phytopathol* **33**: 299–321.
- Barton, G.M., and Medzhitov, R. (2003) Toll-like receptor signaling pathways. *Science* **300**: 1524–1525.
- Barton, G.M., and Medzhitov, R. (2004) Toll signaling: RIPping off the TNF pathway. *Nat Immunol* **5**: 472–474.
- Bartsev, A.V., Deakin, W.J., Boukli, N.M., McAlvin, C.B., Stacey, G., Malnoe, P., et al. (2004) NopL, an effector protein of *Rhizobium* sp. NGR234, thwarts activation of plant defense reactions. *Plant Physiol* **134**: 871–879.
- Benabdillah, R., Mota, L.J., Lutzelschwab, S., Demoinet, E., and Cornelis, G.R. (2004) Identification of a nuclear targeting signal in YopM from *Yersinia* spp. *Microb Pathog* **36**: 247–261.
- Boland, A., and Cornelis, G.R. (1998) Role of YopP in suppression of tumor necrosis factor alpha release by macrophages during *Yersinia* infection. *Infect Immun* **66**: 1878–1884.
- Boller, T. (1995) Chemoperception of microbial signals in plant cells. *Annu Rev Plant Physiol Plant Mol Biol* **46**: 189–214.
- Bonas, U., and Lahaye, T. (2002) Plant disease resistance triggered by pathogen-derived molecules: refined models of specific recognition. *Curr Opin Microbiol* **5**: 44–60.
- Boyes, D.C., Nam, J., and Dangl, J.L. (1998) The *Arabidopsis thaliana* RPM1 disease resistance gene product is a peripheral plasma membrane protein that is degraded coincident with the hypersensitive response. *Proc Natl Acad Sci USA* **95**: 15849–15854.
- Bretz, J.R., Mock, N.M., Charity, J.C., Zeyad, S., Baker, C.J., and Hutcheson, S.W. (2003) A translocated protein tyrosine phosphatase of *Pseudomonas syringae* pv. *tomato* DC3000 modulates plant defence response to infection. *Mol Microbiol* **49**: 389–400.
- Brown, I., Mansfield, J., and Bonas, U. (1995) *hrp* genes in *Xanthomonas campestris* pv. *vesicatoria* determine ability to suppress papilla deposition in pepper mesophyll cells. *Mol Plant Microbe Interact* **8**: 825–836.
- Buttner, D., and Bonas, U. (2003) Common infection strategies of plant and animal pathogenic bacteria. *Curr Opin Plant Biol* **6**: 312–319.
- Buttner, D., Noel, L., Thieme, F., and Bonas, U. (2003) Genomic approaches in *Xanthomonas campestris* pv. *vesicatoria* allow fishing for virulence genes. *J Biotechnol* **106**: 203–214.
- Cao, H., Bowling, S.A., Gordon, A.S., and Dong, X. (1994) Characterization of an *Arabidopsis* mutant that is nonresponsive to inducers of systemic acquired resistance. *Plant Cell* **6**: 1583–1592.
- Cao, H., Li, X., and Dong, X. (1998) Generation of broad-spectrum disease resistance by overexpression of an essential regulatory gene in systemic acquired resistance. *Proc Natl Acad Sci USA* **95**: 6531–6536.
- Carneiro, L.A., Travassos, L.H., and Philpott, D.J. (2004) Innate immune recognition of microbes through Nod1 and Nod2: implications for disease. *Microbes Infect* **6**: 609–616.
- Century, K.S., Shapiro, A.D., Repetti, P.P., Dahlbeck, D., Holub, E., and Staskawicz, B.J. (1997) *NDR1*, a pathogen-induced component required for *Arabidopsis* disease resistance. *Science* **278**: 1963–1965.
- Chamaillard, M., Girardin, S.E., Viala, J., and Philpott, D.J. (2003) Nods, Nalps and Naip: intracellular regulators of bacterial-induced inflammation. *Cell Microbiol* **5**: 581–592.
- Chang, J.H., Goel, A.K., Grant, S.R., and Dangl, J.L. (2004) Wake of the flood: ascribing functions to the wave of type III effector proteins of phytopathogenic bacteria. *Curr Opin Microbiol* **7**: 11–118.
- Chern, M.S., Fitzgerald, H.A., Yadav, R.C., Canlas, P.E., Dong, X., and Ronald, P.C. (2001) Evidence for a disease-resistance pathway in rice similar to the NPR1-mediated signaling pathway in *Arabidopsis*. *Plant J* **27**: 101–113.
- Chen, Z., Kloek, A.P., Cuzick, A., Moeder, W., Tang, D., Innes, R.W., et al. (2004) The *Pseudomonas syringae* type III effector AvrRpt2 functions downstream or independently of SA to promote virulence on *Arabidopsis thaliana*. *Plant J* **37**: 494–504.
- Cohn, J., Sessa, G., and Martin, G.B. (2001) Innate immunity in plants. *Curr Opin Immunol* **13**: 55–62.
- Collmer, A., Lindeberg, M., Petnicki-Ocwieja, T., Schneider, D., and Alfano, J.R. (2002) Genomic mining type III secretion system effectors in *Pseudomonas syringae* yields new picks for all TTSS prospectors. *Trends Microbiol* **10**: 462–469.
- Cornelis, G.R. (2000) Molecular and cell biology aspects of plague. *Proc Natl Acad Sci USA* **97**: 8778–8783.
- Cornelis, G.R. (2002a) *Yersinia* type III secretion: send in the effectors. *J Cell Biol* **158**: 401–408.
- Cornelis, G.R. (2002b) The *Yersinia* Ysc-Yop 'type III' weaponry. *Nat Rev Mol Cell Biol* **3**: 742–752.
- Cossart, P., and Sansonetti, P.J. (2004) Bacterial invasion: the paradigms of enteroinvasive pathogens. *Science* **304**: 242–248.
- Dale, C., Young, S.A., Haydon, D.T., and Welburn, S.C. (2001) The insect endosymbiont *Sodalis glossinidius* utilizes a type III secretion system for cell invasion. *Proc Natl Acad Sci USA* **98**: 1883–1888.
- Dangl, J.L., and Jones, J.D. (2001) Plant pathogens and integrated defence responses to infection. *Nature* **411**: 826–833.
- DebRoy, S., Thilmony, R., Kwack, Y.B., Nomura, K., and He, S.Y. (2004) A family of conserved bacterial effectors inhibits salicylic acid-mediated basal immunity and promotes disease necrosis in plants. *Proc Natl Acad Sci USA* **101**: 9927–9932.
- Delaney, T.P., Friedrich, L., and Ryals, J.A. (1995) *Arabidopsis* signal transduction mutant defective in chemically and biologically induced disease resistance. *Proc Natl Acad Sci USA* **92**: 6602–6606.
- Deslandes, L., Olivier, J., Theulieres, F., Hirsch, J., Feng, D.X., Bittner-Eddy, P., et al. (2002) Resistance to *Ralstonia solanacearum* in *Arabidopsis thaliana* is conferred by the recessive *RRS1-R* gene, a member of a novel family of resistance genes. *Proc Natl Acad Sci USA* **99**: 2404–2409.
- Deslandes, L., Olivier, J., Peeters, N., Feng, D.X., Khounlotham, M., Boucher, C., et al. (2003) Physical interaction between *RRS1-R*, a protein conferring resistance to bacterial wilt, and PopP2, a type III effector targeted to the plant nucleus. *Proc Natl Acad Sci USA* **100**: 8024–8029.
- Dow, M., Newman, M.A., and von Roepenack, E. (2000) The induction and modulation of plant defense responses by bacterial lipopolysaccharides. *Annu Rev Phytopathol* **38**: 241–261.

- Durrant, W.E., and Dong, X. (2004) Systemic acquired resistance. *Annu Rev Phytopathol* **42**: 185–209.
- Espinosa, A., Guo, M., Tam, V.C., Fu, Z.Q., and Alfano, J.R. (2003) The *Pseudomonas syringae* type III-secreted protein HopPtoD2 possesses protein tyrosine phosphatase activity and suppresses programmed cell death in plants. *Mol Microbiol* **49**: 377–387.
- Eulgem, T., Rushton, P.J., Robatzek, S., and Somssich, I.E. (2000) The WRKY superfamily of plant transcription factors. *Trends Plant Sci* **5**: 199–206.
- Felix, G., and Boller, T. (2003) Molecular sensing of bacteria in plants. The highly conserved RNA-binding motif RNP-1 of bacterial cold shock proteins is recognized as an elicitor signal in tobacco. *J Biol Chem* **278**: 6201–6208.
- Falk, A., Feys, B.J., Frost, L.N., Jones, J.D., Daniels, M.J., and Parker, J.E. (1999) EDS1, an essential component of R gene-mediated disease resistance in *Arabidopsis* has homology to eukaryotic lipases. *Proc Natl Acad Sci USA* **96**: 3292–3297.
- Felix, G., Regenass, M., and Boller, T. (1993) Specific perception of subnanomolar concentrations of chitin fragments by tomato cells. Induction of extracellular alkalization, changes in protein phosphorylation, and establishment of a refractory state. *Plant J* **4**: 307–316.
- Flor (1971) Current status of the gene-for-gene concept. *Annu Rev Phytopathol* **9**: 275–296.
- Friedrich, L., Lawton, K., Dietrich, R., Willits, M., Cade, R., and Ryals, J. (2001) NIM1 overexpression in *Arabidopsis* potentiates plant disease resistance and results in enhanced effectiveness of fungicides. *Mol Plant Microbe Interact* **14**: 1114–1124.
- Fu, Y., and Galán, J.E. (1998) The *Salmonella typhimurium* tyrosine phosphatase SptP is translocated into host cells and disrupts the actin cytoskeleton. *Mol Microbiol* **27**: 359–368.
- Gabriel, D.W., and Rolfe, B.G. (1990) Working models of specific recognition in plant–microbe interactions. *Annu Rev Phytopathol* **28**: 365–391.
- Gaffney, T., Friedrich, L., Vernooij, B., Negrotto, D., Nye, G., Uknes, S., *et al.* (1993) Requirement of salicylic-acid for the induction of systemic acquired-resistance. *Science* **261**: 754–756.
- Galán, J.E., and Collmer, A. (1999) Type III secretion machines: bacterial devices for protein delivery into host cells. *Science* **284**: 1322–1328.
- Gassmann, W., Hinsch, M.E., and Staskawicz, B.J. (1999) The *Arabidopsis* *RPS4* bacterial-resistance gene is a member of the TIR-NBS-LRR family of disease-resistance genes. *Plant J* **20**: 265–277.
- Girardin, S.E., Sansonetti, P.J., and Philpott, D.J. (2002) Intracellular vs extracellular recognition of pathogens – common concepts in mammals and flies. *Trends Microbiol* **10**: 193–199.
- Gomez-Gomez, L., and Boller, T. (2000) FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in *Arabidopsis*. *Mol Cell* **5**: 1003–1011.
- Gomez-Gomez, L., Bauer, Z., and Boller, T. (2001) Both the extracellular leucine-rich repeat domain and the kinase activity of FLS2 are required for flagellin binding and signaling in *Arabidopsis*. *Plant Cell* **13**: 1155–1163.
- Gomez-Gomez, L., and Boller, T. (2002) Flagellin perception: a paradigm for innate immunity. *Trends Plant Sci* **7**: 251–256.
- Granado, J., Felix, G., and Boller, T. (1995) Perception of fungal sterols in plants (subnanomolar concentrations of ergosterol elicit extracellular alkalization in tomato cells). *Plant Physiol* **107**: 485–490.
- Greenberg, J.T., and Vinatzer, B.A. (2003) Identifying type III effectors of plant pathogens and analyzing their interaction with plant cells. *Curr Opin Microbiol* **6**: 20–28.
- Guan, K.L., and Dixon, J.E. (1990) Protein tyrosine phosphatase activity of an essential virulence determinant in *Yersinia*. *Science* **249**: 553–556.
- Guttman, D.S., Vinatzer, B.A., Sarkar, S.F., Ranall, M.V., Kettler, G., and Greenberg, J.T. (2002) A functional screen for the type III (Hrp) secretome of the plant pathogen *Pseudomonas syringae*. *Science* **295**: 1722–1726.
- Haller, J.C., Carlson, S., Pederson, K.J., and Pierson, D.E. (2000) A chromosomally encoded type III secretion pathway in *Yersinia enterocolitica* is important in virulence. *Mol Microbiol* **36**: 1436–1446.
- Haraga, A., and Miller, S.I. (2003) A *Salmonella enterica* serovar *typhimurium* translocated leucine-rich repeat effector protein inhibits NF-kappa B-dependent gene expression. *Infect Immun* **71**: 4052–4058.
- Hauck, P., Thilmony, R., and He, S.Y. (2003) A *Pseudomonas syringae* type III effector suppresses cell wall-based extracellular defense in susceptible *Arabidopsis* plants. *Proc Natl Acad Sci USA* **100**: 8577–8582.
- Heath, M.C. (2000) Nonhost resistance and nonspecific plant defenses. *Curr Opin Plant Biol* **3**: 315–319.
- Hobbie, S., Chen, L.M., Davis, R.J., and Galan, J.E. (1997) Involvement of mitogen-activated protein kinase pathways in the nuclear responses and cytokine production induced by *Salmonella typhimurium* in cultured intestinal epithelial cells. *J Immunol* **159**: 5550–5559.
- Holden, D.W. (2002) Trafficking of the *Salmonella* vacuole in macrophages. *Traffic* **3**: 161–169.
- Holt, B.F., 3rd, Hubert, D.A., and Dangl, J.L. (2003) Resistance gene signaling in plants – complex similarities to animal innate immunity. *Curr Opin Immunol* **15**: 20–25.
- Horng, T., Barton, G.M., Flavell, R.A., and Medzhitov, R. (2002) The adaptor molecule TIRAP provides signalling specificity for Toll-like receptors. *Nature* **420**: 329–333.
- Hotson, A., Chosed, R., Shu, H., Orth, K., and Mudgett, M.B. (2003) *Xanthomonas* type III effector XopD targets SUMO-conjugated proteins *in planta*. *Mol Microbiol* **50**: 377–389.
- Hueck, C.J. (1998) Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol Mol Biol Rev* **62**: 379–433.
- Imler, J.L., and Hoffmann, J.A. (2001) Toll receptors in innate immunity. *Trends Cell Biol* **11**: 304–311.
- Innes, R. (2003) New effects of type III effectors. *Mol Microbiol* **50**: 363–365.
- Innes, R.W. (2004) Guarding the goods. New insights into the central alarm system of plants. *Plant Physiol* **135**: 695–701.
- Inohara, N., and Nunez, G. (2003) NODs: intracellular proteins involved in inflammation and apoptosis. *Nat Rev Immunol* **3**: 371–382.
- Jackson, R.W., Athanassopoulos, E., Tsiamis, G., Mansfield, J.W., Sesma, A., Arnold, D.L., *et al.* (1999) Identification of a pathogenicity island, which contains genes for virulence and avirulence, on a large native plasmid in the bean pathogen *Pseudomonas syringae* pathovar phaseolicola. *Proc Natl Acad Sci USA* **96**: 10875–10880.

- Jakobek, J.L., Smith, J.A., and Lindgren, P.B. (1993) Suppression of bean defense responses by *Pseudomonas syringae*. *Plant Cell* **5**: 57–63.
- Jamir, Y., Guo, M., Oh, H.-S., Petnicki-Ocwieja, T., Chen, S., Tang, X., *et al.* (2004) Identification of *Pseudomonas syringae* type III effectors that suppress programmed cell death in plants and yeast. *Plant J* **37**: 554–565.
- Jin, H., Axtell, M.J., Dahlbeck, D., Ekwenna, O., Zhang, S., Staskawicz, B., and Baker, B. (2002) NPK1, an MEKK1-like mitogen-activated protein kinase kinase, regulates innate immunity and development in plants. *Dev Cell* **3**: 291–297.
- Jones, D.A., and Takemoto, D. (2004) Plant innate immunity – direct and indirect recognition of general and specific pathogen-associated molecules. *Curr Opin Immunol* **16**: 48–62.
- Kamoun, S., Kamdar, H.V., Tola, E., and Kado, C.I. (1992) Incompatible interactions between crucifers and *Xanthomonas campestris* involve vascular hypersensitive response: role of the *hrpX* locus. *Mol Plant Microbe Interact* **5**: 22–33.
- Kaniga, K., Uralil, J., Bliska, J.B., and Galán, J.E. (1996) A secreted protein tyrosine phosphatase with modular effector domains in the bacterial pathogen *Salmonella typhimurium*. *Mol Microbiol* **21**: 633–641.
- Keen, N.T. (1990) Gene-for-gene complementarity in plant-pathogen interactions. *Annu Rev Genet* **24**: 447–463.
- Kim, C.Y., and Zhang, S. (2004) Activation of a mitogen-activated protein kinase cascade induces WRKY family of transcription factors and defense genes in tobacco. *Plant J* **38**: 142–151.
- Kim, D.H., Feinbaum, R., Alloing, G., Emerson, F.E., Garsin, D.A., Inoue, H., *et al.* (2002) A conserved p38 MAP kinase pathway in *Caenorhabditis elegans* innate immunity. *Science* **297**: 623–626.
- Kobe, B., and Kajava, A.V. (2001) The leucine-rich repeat as a protein recognition motif. *Curr Opin Struct Biol* **11**: 725–732.
- Lam, E., Kato, N., and Lawton, M. (2001) Programmed cell death, mitochondria and the plant hypersensitive response. *Nature* **411**: 848–853.
- Lee, F.S., Hagler, J., Chen, Z.J., and Maniatis, T. (1997) Activation of the I κ B kinase complex by MEKK1, a kinase of the JNK pathway. *Cell* **88**: 213–222.
- Limpens, E., Franken, C., Smit, P., Willemse, J., Bisseling, T., and Geurts, R. (2003) LysM domain receptor kinases regulating rhizobial Nod factor-induced infection. *Science* **302**: 630–633.
- Lin, S.L., Le, T.X., and Cowen, D.S. (2003) SptP, a *Salmonella typhimurium* type III-secreted protein, inhibits the mitogen-activated protein kinase pathway by inhibiting Raf activation. *Cell Microbiol* **5**: 267–275.
- Lopez-Solanilla, E., Bronstein, P.A., Schneider, A.R., and Collmer, A. (2004) HopPtoN is a *Pseudomonas syringae* TTSS effector with cysteine protease that suppresses pathogen-induced necrosis associated with both compatible and incompatible plant interactions. *Mol Microbiol* (in press).
- Mackey, D., Holt, B.F., III, Wiig, A., and Dangl, J.L. (2002) RIN4 interacts with *Pseudomonas syringae* type III effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. *Cell* **108**: 743–754.
- Mackey, D., Belkadir, Y., Alonso, J.M., Ecker, J.R., and Dangl, J.L. (2003) *Arabidopsis* RIN4 is a target of the type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance. *Cell* **112**: 379–389.
- Madsen, E.B., Madsen, L.H., Radutoiu, S., Olbryt, M., Rakwalska, M., Szczygłowski, K., *et al.* (2003) A receptor kinase gene of the LysM type is involved in legume perception of rhizobial signals. *Nature* **425**: 637–640.
- Marie, C., Broughton, W.J., and Deakin, W.J. (2001) Rhizobium type III secretion systems: legume charmers or alarmers? *Curr Opin Plant Biol* **4**: 336–342.
- Martin, G.B., Bogdanove, A.J., and Sessa, G. (2003) Understanding the functions of plant disease resistance proteins. *Annu Rev Plant Biol* **54**: 23–61.
- Medzhitov, R. (2001) Toll-like receptors and innate immunity. *Nat Rev Immunol* **1**: 135–145.
- Medzhitov, R., and Janeway, C.A., Jr (1997) Innate immunity: the virtues of a nonclonal system of recognition. *Cell* **91**: 295–298.
- Meijer, L.K., Schesser, K., Wolf-Watz, H., Sassone-Corsi, P., and Pettersson, S. (2000) The bacterial protein YopJ abrogates multiple signal transduction pathways that converge on the transcription factor CREB. *Cell Microbiol* **2**: 231–238.
- Mills, S.D., Boland, A., Sory, M.-P., Van Der Smissen, P., Kerbouch, C., Finlay, B.B., and Cornelis, G.R. (1997) *Yersinia enterocolitica* induces apoptosis in macrophages by a process requiring functional type III secretion and translocation mechanisms and involving YopP, presumably acting as an effector protein. *Proc Natl Acad Sci USA* **94**: 12638–12643.
- Monack, D.M., Mecsas, J., Ghori, N., and Falkow, S. (1997) *Yersinia* signals macrophages to undergo apoptosis and YopJ is necessary for this cell death. *Proc Natl Acad Sci USA* **94**: 10385–10390.
- Murli, S., Watson, R.O., and Galán, J.E. (2001) Role of tyrosine kinases and the tyrosine phosphatase SptP in the interaction of *Salmonella* with host cells. *Cell Microbiol* **3**: 795–810.
- Navarro, L., Zipfel, C., Rowland, O., Keller, I., Robatzek, S., Boller, T., and Jones, J.D. (2004) The transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis. *Plant Physiol* **135**: 1–16.
- Nimchuk, Z., Rohmer, L., Chang, J.H., and Dangl, J.L. (2001) Knowing the dancer from the dance: R-gene products and their interactions with other proteins from host and pathogen. *Curr Opin Plant Biol* **4**: 288–294.
- Nimchuk, Z., Eulgem, T., Holt, B.F., 3rd, and Dangl, J.L. (2003) Recognition and response in the plant immune system. *Annu Rev Genet* **37**: 579–609.
- Nuhse, T.S., Peck, S.C., Hirt, H., and Boller, T. (2000) Microbial elicitors induce activation and dual phosphorylation of the *Arabidopsis thaliana* MAPK 6. *J Biol Chem* **275**: 7521–7526.
- Nurnberger, T., and Brunner, F. (2002) Innate immunity in plants and animals: emerging parallels between the recognition of general elicitors and pathogen-associated molecular patterns. *Curr Opin Plant Biol* **5**: 318–324.
- Nurnberger, T., Brunner, F., Kemmerling, B., and Piater, L. (2004) Innate immunity in plants and animals: striking similarities and obvious differences. *Immunol Rev* **198**: 249–266.
- Orth, K. (2002) Function of the *Yersinia* effector YopJ. *Curr Opin Microbiol* **5**: 38–43.

- Orth, K., Palmer, L.E., Bao, Z.Q., Stewart, S., Rudolph, A.E., Bliska, J.B., and Dixon, J.E. (1999) Inhibition of the mitogen-activated protein kinase superfamily by a *Yersinia* effector. *Science* **285**: 1920–1923.
- Orth, K., Xu, Z., Mudgett, M.B., Bao, Z.Q., Palmer, L.E., Bliska, J.B., *et al.* (2000) Disruption of signaling by *Yersinia* effector YopJ, a Ubiquitin-like protein protease. *Science* **290**: 1594–1597.
- Palmer, L.E., Hobbie, S., Galán, J.E., and Bliska, J.B. (1998) YopJ of *Yersinia pseudotuberculosis* is required for the inhibition of macrophage TNF- α production and downregulation of the MAP kinases p38 and JNK. *Mol Microbiol* **27**: 953–965.
- Pasare, C., and Medzhitov, R. (2003) Toll-like receptors: balancing host resistance with immune tolerance. *Curr Opin Immunol* **15**: 677–682.
- Pedley, K.F., and Martin, G.B. (2003) Molecular basis of Pto-mediated resistance to bacterial speck disease in tomato. *Annu Rev Phytopathol* **41**: 215–243.
- Petnicki-Ocwieja, T., Schneider, D.J., Tam, V.C., Chancey, S.T., Shan, L., Jamir, Y., *et al.* (2002) Genomewide identification of proteins secreted by the Hrp type III protein secretion system of *Pseudomonas syringae* pv. *tomato* DC3000. *Proc Natl Acad Sci USA* **99**: 7652–7657.
- del Pozo, O., Pedley, K.F., and Martin, G.B. (2004) MAP-KKK α is a positive regulator of cell death associated with both plant immunity and disease. *EMBO J* **23**: 3072–3082.
- Radutoiu, S., Madsen, L.H., Madsen, E.B., Felle, H.H., Ume-hara, Y., Gronlund, M., *et al.* (2003) Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. *Nature* **425**: 585–592.
- Ritter, C., and Dangel, J.L. (1996) Interference between two specific pathogen recognition events mediated by distinct plant disease resistance genes. *Plant Cell* **8**: 251–257.
- Ruckdeschel, K., Harb, S., Roggenkamp, A., Hornef, M., Zumbihl, R., Kohler, S., *et al.* (1998) *Yersinia enterocolitica* impairs activation of transcription factor NF-kappaB: involvement in the induction of programmed cell death and in the suppression of the macrophage tumor necrosis factor alpha production. *J Exp Med* **187**: 1069–1079.
- Ruckdeschel, K., Machold, J., Roggenkamp, A., Schubert, S., Pierre, J., Zumbihl, R., *et al.* (1997) *Yersinia enterocolitica* promotes deactivation of macrophage mitogen-activated protein kinases extracellular signal-regulated kinase-1/2, 38, and c-Jun NH2-terminal kinase. Correlation with its inhibitory effect on tumor necrosis factor-alpha production. *J Biol Chem* **272**: 15920–15927.
- Ryals, J., Weymann, K., Lawton, K., Friedrich, L., Ellis, D., Steiner, H.-Y., *et al.* (1997) The *Arabidopsis* NIM-1 protein shows homology to the mammalian transcription factor inhibitor I κ B. *Plant Cell* **9**: 425–439.
- Salanoubat, M., Genin, S., Artiguenave, F., Gouzy, J., Mangenot, S., Arlat, M., *et al.* (2002) Genome sequence of the plant pathogen *Ralstonia solanacearum*. *Nature* **415**: 497–502.
- Sauvonnet, N., Lambermont, I., van der Bruggen, P., and Cornelis, G.R. (2002a) YopH prevents monocyte chemoattractant protein 1 expression in macrophages and T-cell proliferation through inactivation of the phosphatidylinositol 3-kinase pathway. *Mol Microbiol* **45**: 805–815.
- Sauvonnet, N., Pradet-Balade, B., Garcia-Sanz, J.A., and Cornelis, G.R. (2002b) Regulation of mRNA expression in macrophages after *Yersinia enterocolitica* infection. Role of different Yop effectors. *J Biol Chem* **277**: 25133–25142.
- Schneider, D.S. (2002) Plant immunity and film noir: what gumshoe detectives can teach us about plant–pathogen interactions. *Cell* **109**: 537–540.
- Schesser, K., Spiik, A.-K., Dukuzmuremyi, J.-M., Neurath, M.F., Pettersson, S., and Wolf-Watz, H. (1998) The *yopJ* locus is required for *Yersinia*-mediated inhibition of NF- κ B activation and cytokine expression: YopJ contains a eukaryotic SH2-like domain that is essential for its repressive activity. *Mol Microbiol* **28**: 1067–1079.
- Shao, F., Merritt, P.M., Bao, Z., Innes, R.W., and Dixon, J.E. (2002) A *Yersinia* effector and a *Pseudomonas* avirulence protein define a family of cysteine proteases functioning in bacterial pathogenesis. *Cell* **109**: 575–588.
- Shao, F., Golstein, C., Ade, J., Stoutemyer, M., Dixon, J.E., and Innes, R.W. (2003) Cleavage of *Arabidopsis* PBS1 by a bacterial type III effector. *Science* **301**: 1230–1233.
- Shea, J.E., Hensel, M., Gleeson, C., and Holden, D.W. (1996) Identification of a virulence locus encoding a second type III secretion system in *Salmonella typhimurium*. *Proc Natl Acad Sci USA* **93**: 2593–2597.
- Shen, Y., Sharma, P., da Silva, F.G., and Ronald, P. (2002) The *Xanthomonas oryzae* pv. *oryzae* *raxP* and *raxQ* genes encode an ATP sulphurylase and adenosine-5'-phosphosulphate kinase that are required for AvrXa21 avirulence activity. *Mol Microbiol* **44**: 37–48.
- Shiu, S.H., and Bleecker, A.B. (2003) Expansion of the receptor-like kinase/Pelle gene family and receptor-like proteins in *Arabidopsis*. *Plant Physiol* **132**: 530–543.
- da Silva, A.C., Ferro, J.A., Reinach, F.C., Farah, C.S., Furlan, L.R., Quaggio, R.B., *et al.* (2002) Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities. *Nature* **417**: 459–463.
- Skrzypek, E., Cowan, C., and Straley, S.C. (1998) Targeting of the *Yersinia pestis* YopM protein into HeLa cells and intracellular trafficking to the nucleus. *Mol Microbiol* **30**: 1051–1065.
- Song, W.-Y., Wang, G.-L., Chen, L.-L., Kim, H.-S., Pi, L.-Y., Holsten, T., *et al.* (1995) A receptor kinase-like protein encoded by the rice disease resistance gene, *Xa21*. *Science* **270**: 1804–1806.
- Spaink, H.P. (2004) Specific recognition of bacteria by plant LysM domain receptor kinases. *Trends Microbiol* **12**: 201–204.
- Takeda, K., and Akira, S. (2003) Toll receptors and pathogen resistance. *Cell Microbiol* **5**: 143–153.
- Takeda, K., Kaisho, T., and Akira, S. (2003) Toll-like receptors. *Annu Rev Immunol* **21**: 335–376.
- Tao, Y., Xie, Z., Chen, W., Glazebrook, J., Chang, H.S., Han, B., *et al.* (2003) Quantitative nature of *Arabidopsis* responses during compatible and incompatible interactions with the bacterial pathogen *Pseudomonas syringae*. *Plant Cell* **15**: 317–330.
- Tenor, J.L., McCormick, B.A., Ausubel, F.M., and Aballay, A. (2004) *Caenorhabditis elegans*-based screen identifies *Salmonella* virulence factors required for conserved host–pathogen interactions. *Curr Biol* **14**: 1018–1024.
- Thordal-Christensen, H. (2003) Fresh insights into processes of nonhost resistance. *Curr Opin Plant Biol* **6**: 351–357.
- Torii, K.U. (2000) Receptor kinase activation and signal transduction in plants: an emerging picture. *Curr Opin Plant Biol* **3**: 361–367.
- Toyotome, T., Suzuki, T., Kuwae, A., Nonaka, T., Fukuda, H., Imajoh-Ohmi, S., *et al.* (2001) *Shigella* protein IpaH (9.8) is secreted from bacteria within mammalian cells and

- transported to the nucleus. *J Biol Chem* **276**: 32071–32079.
- Tsiamis, G., Mansfield, J.W., Hockenhull, R., Jackson, R.W., Sesma, A., Athanassopoulos, E., *et al.* (2000) Cultivar-specific avirulence and virulence functions assigned to *avrPphF* in *Pseudomonas syringae* pv. *phaseolicola*, the cause of bean halo-blight disease. *EMBO J* **19**: 3204–3214.
- Viboud, G.I., So, S.S., Ryndak, M.B., and Bliska, J.B. (2003) Proinflammatory signalling stimulated by the type III translocation factor YopB is counteracted by multiple effectors in epithelial cells infected with *Yersinia pseudotuberculosis*. *Mol Microbiol* **47**: 1305–1315.
- van Wees, S.C.M., and Glazebrook, J. (2003) Loss of non-host resistance of *Arabidopsis* NahG to *Pseudomonas syringae* pv. *phaseolicola* is due to degradation products of salicylic acid. *Plant J* **33**: 733–742.
- Wildermuth, M.C., Dewdney, J., Wu, G., and Ausubel, F.M. (2001) Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* **414**: 562–565.
- Yamamoto, M., Sato, S., Hemmi, H., Hoshino, K., Kaisho, T., Sanjo, H., *et al.* (2003) Role of adaptor TRIF in the MyD88-independent toll-like receptor signaling pathway. *Science* **301**: 640–643.
- Yang, K.-Y., Liu, Y., and Zhang, S. (2001) Activation of a mitogen-activated protein kinase pathway is involved in disease resistance in tobacco. *Proc Natl Acad Sci USA* **98**: 741–746.
- Yao, T., Mecsas, J., Healy, J.I., Falkow, S., and Chien, Y. (1999) Suppression of T and B lymphocyte activation by a *Yersinia pseudotuberculosis* virulence factor, YopH. *J Exp Med* **190**: 1343–1350.
- Zaharik, M.L., Gruenheid, S., Perrin, A.J., and Finlay, B.B. (2002) Delivery of dangerous goods: type III secretion in enteric pathogens. *Int J Med Microbiol* **291**: 593–603.
- Zhang, S., and Klessig, D.F. (2001) MAPK cascades in plant defense signaling. *Trends Plant Sci* **6**: 520–527.
- Zipfel, C., Robatzek, S., Navarro, L., Oakeley, E.J., Jones, J.D., Felix, G., and Boller, T. (2004) Bacterial disease resistance in *Arabidopsis* through flagellin perception. *Nature* **428**: 764–767.