

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications, Department of
Mathematics

Mathematics, Department of

12-21-2018

Lozenge tilings of a halved hexagon with an array of triangles removed from the boundary, part II

Tri Lai

Follow this and additional works at: <https://digitalcommons.unl.edu/mathfacpub>



Part of the [Applied Mathematics Commons](#), and the [Mathematics Commons](#)

This Article is brought to you for free and open access by the Mathematics, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Mathematics by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Lozenge tilings of a halved hexagon with an array of triangles removed from the boundary, part II

Tri Lai*

University of Nebraska – Lincoln
Lincoln, NE 68588 USA

tlai3@unl.edu

Submitted: Nov 28, 2017; Accepted: Oct 29, 2018; Published: Dec 21, 2018

© The author. Released under the CC BY-ND license (International 4.0).

Abstract

Proctor’s work on staircase plane partitions yields an exact enumeration of lozenge tilings of a halved hexagon on the triangular lattice. Rohatgi later extended this tiling enumeration to a halved hexagon with a triangle cut off from the boundary. In his previous paper, the author proved a common generalization of Proctor’s and Rohatgi’s results by enumerating lozenge tilings of a halved hexagon in the case an array of an *arbitrary* number of triangles has been removed from a non-staircase side. In this paper we consider the other case when the array of triangles has been removed from the staircase side of the halved hexagon. Our result also implies an explicit formula for the number of tilings of a hexagon with an array of triangles removed perpendicularly to the symmetry axis.

Mathematics Subject Classifications: 05A15, 05B45, 05C30

1 Introduction

A *plane partition* is an array of positive integers $p_{i,j}$ (called “*parts*”) so that $p_{i,j} \geq \max(p_{i+1,j}, p_{i,j+1})$. R. Proctor [12] proved a simple product formula for the number of a class of staircase plane partitions. The plane partitions in Proctor’s result are in bijection with lozenge tilings of a hexagon of side-lengths a, b, c, a, b, c (in the counter clockwise order, starting from the northwestern side) on the triangular lattice with a ‘maximal staircase’ cut off, denoted by $\mathcal{P}_{a,b,c}$ (see Figure 1(a)). Here a *lozenge* (or *unit rhombus*) is the union of any two unit equilateral triangles sharing an edge, and a *lozenge tiling* of a region is a covering of the region by lozenges so that there are no gaps or overlaps. This way Proctor’s result yields the following tiling enumeration.

*Supported in part by Simons Foundation Collaboration Grant (#585923).




Figure 1: (a) The halved hexagon (with defects) $\mathcal{P}_{4,7,3}$. (b) The weighted halved hexagon $\mathcal{P}'_{4,7,3}$. (c) and (d) The regions in Rohatgi's paper [13].

Theorem 1 (Proctor [12]). *For any non-negative integers a, b , and c with $a \leq b$, we have*

$$M(\mathcal{P}_{a,b,c}) = \prod_{i=1}^a \left[\prod_{j=1}^{b-a+1} \frac{c+i+j-1}{i+j-1} \prod_{j=b-a+2}^{b-a+i} \frac{2c+i+j-1}{i+j-1} \right], \quad (1)$$

where $M(R)$ denotes the number of lozenge tilings of a region R , and where empty products are taken to be 1.

When $a = b$, the region $\mathcal{P}_{a,b,c}$ above can be viewed as a half of a symmetry hexagon with a zigzag cut along the symmetry axis. In this point of view, we usually call the region $\mathcal{P}_{a,b,c}$ a *halved hexagon (with defects)*. We also note that when $a = b$, Proctor's Theorem 1 implies an exact enumeration for one of the ten symmetry classes of plane partitions, the *transposed-complementary plane partitions* (see e.g. [14]).

Lozenges in a region can carry weights. In the weighted case, we use the notation $M(R)$ for the sum of weights of all lozenge tilings of R , where the *weight* of a lozenge tiling is the weight product of its constituent lozenges. We still call $M(R)$ the (*weighted*) *tiling number* of R . We also consider the weighted counterpart $\mathcal{P}'_{a,b,c}$ of $\mathcal{P}_{a,b,c}$, where all the lozenges along the staircase cut are weighted by $1/2$ (see the lozenges with shaded cores in Figure 1(b)). M. Ciucu [2] proved the following weighted version of Theorem 1.

Theorem 2 (Ciucu [2]). *For any non-negative integers a , b , and c with $a \leq b$*

$$M(\mathcal{P}'_{a,b,c}) = 2^{-a} \prod_{i=1}^{2c+b-a+i} \frac{2c+b-a+i}{c+b-a+i} \prod_{i=1}^a \left[\prod_{j=1}^{b-a+1} \frac{c+i+j-1}{i+j-1} \prod_{j=b-a+2}^{b-a+i} \frac{2c+i+j-1}{i+j-1} \right]. \quad (2)$$

From now on, we use the notations $P_{a,b,c}$ and $P'_{a,b,c}$ for the numbers of tilings of the regions $\mathcal{P}_{a,b,c}$ and $\mathcal{P}'_{a,b,c}$, respectively.

Tiling enumerations of halved hexagons with certain defects have been investigated by a number of authors (see e.g. [2], [4], [13], [7], [8] and the lists of references therein). It is worth noticing that R. Rohatgi [13] generalizes the regions $P_{a,a,c}$ and $P'_{a,a,c}$ to halved hexagons with a triangle removed on the northeastern side (see Figures 1(c) and (d)). In his previous paper [9], the author generalized the tiling enumerations of halved hexagons by Proctor and Rohatgi to halved hexagons in which an array of *an arbitrary number* of adjacent triangles has been removed from the northeastern side (see Figure 2 for examples of the regions). In this paper, we investigate the other case when the array of triangles has been removed from the *western* side of the halved hexagon as shown in Figures 4, 5, and 6. Based on the positions of the array of removed triangles and the weight assignments of the lozenges along the staircase side, we have *eight* families of defected halved hexagons that will be described in detail in Section 2. We will show that the numbers of tilings of these regions are *all* given by simple product formulas.

Our explicit enumerations for these halved hexagons also imply an exact tiling formula for a symmetric hexagon with an array of triangular holes placed perpendicularly to the symmetry axis (see Theorem 12).

It is worth noticing that Ciucu [3], in his effort on finding duals of the MacMahon classical theorem on plane partitions [11], gave a closed form product formula for the tiling number of a hexagon (not necessarily symmetric) with an array of triangular holes in the center. In Ciucu's result, if the array of triangular holes moves far away from the center, the tiling number is *not* a simple product anymore. In contrast, our result (Theorem 12 in Section 2) shows that we have a nice tiling formula for *any* array of triangular holes placed perpendicularly to the symmetric axis of the hexagon.

The rest of the paper is organized as follows. Due to the large number of new regions needed to define, we leave the precise statement of our main results (Theorems 4–12) to Section 2. In Section 3, we quote several fundamental results in the enumeration of tilings and introduce the particular version of Kuo condensation [6] that we will employ in our proofs. Section 4 are devoted to the proofs of our main theorems.

2 Precise statement of the main results

We define the *Pochhammer symbol* $(x)_n$ by

$$(x)_n = \begin{cases} x(x+1) \dots (x+n-1) & \text{if } n > 0; \\ 1 & \text{if } n = 0; \\ \frac{1}{(x-1)(x-2) \dots (x+n)} & \text{if } n < 0. \end{cases} \quad (3)$$




Figure 2: Halved hexagons with an array of triangles removed from the non-staircase boundary in [9].

We also use the ‘skipping’ version of the Pochhammer symbol:

$$[x]_n = \begin{cases} x(x+2)\dots(x+2(n-1)) & \text{if } n > 0; \\ 1 & \text{if } n = 0; \\ \frac{1}{(x-2)(x-4)\dots(x+2n)} & \text{if } n < 0. \end{cases} \quad (4)$$

We also define the following two products:

$$T(x, n, m) = \prod_{i=0}^{m-1} (x+i)_{n-2i} \quad (5)$$

and

$$V(x, n, m) = \prod_{i=0}^{m-1} [x+2i]_{n-2i}. \quad (6)$$

Let $\mathbf{a} = (a_1, a_2, \dots, a_n)$ be a sequence. We will use several operations on sequences as follows:

$$O(\mathbf{a}) = \sum_{i \text{ odd}} a_i, \quad E(\mathbf{a}) = \sum_{i \text{ even}} a_i, \quad (7)$$

$$\mathbf{s}_k(\mathbf{a}) = \sum_{i=1}^k a_i, \quad \mathbf{o}_k(\mathbf{a}) = \sum_{i \geq k} a_{2i-1}, \quad \mathbf{e}_k(\mathbf{a}) = \sum_{i \geq k} a_{2i}. \quad (8)$$

Before going to the statement of our main results, we quote here the tiling formulas of halved hexagons with triangles removed on the base (see Lemma 1.4 in [9]).




Figure 3: (a) The region $\mathcal{Q}(2, 1, 2, 2)$. (b) The region $\mathcal{Q}(0, 1, 1, 1, 2, 2)$. (c) The region $\mathcal{Q}'(2, 1, 2, 2)$. (d) The region $\mathcal{K}(3, 1, 2, 2)$. (e) The region $\mathcal{K}(0, 2, 1, 1, 2, 2)$. (f) The region $\mathcal{K}'(3, 1, 2, 2)$. The lozenges with shaded cores are weighted by $\frac{1}{2}$. This figure first appeared in [9].

Assume that $\mathbf{t} = (t_1, t_2, \dots, t_{2l})$ is a sequence of non-negative integers. Consider a trapezoidal region whose northern, northeastern, and southern sides have lengths $O(\mathbf{t})$, $2E(\mathbf{t})$, and $E(\mathbf{t}) + O(\mathbf{t})$, respectively, and whose western side follows a vertical zigzag lattice paths with $E(\mathbf{t})$ steps. We remove the triangles of sides t_{2i} 's from the base of the latter region so that the distances between two consecutive triangles are t_{2i-1} 's. Denote the resulting region by $\mathcal{Q}(\mathbf{t}) = \mathcal{Q}(t_1, t_2, \dots, t_{2l})$ (see the regions in Figure 3(a) for the case when $t_1 > 0$ and Figure 3(b) for the case when $t_1 = 0$). Inspired by the weighted region $\mathcal{P}'_{a,b,c}$, we consider the weighted counterpart $\mathcal{Q}'(\mathbf{t})$ of the newly defined region, where the vertical lozenges on the western side are weighted by $\frac{1}{2}$ (see Figure 3(c)); the vertical lozenges with shaded cores are weighted by $\frac{1}{2}$.

We are also interested in a variation of the above \mathcal{Q} -type regions as follows. Consider the trapezoidal region whose northern, northeastern, and southern sides have lengths $O(\mathbf{t})$, $2E(\mathbf{t}) - 1$, $E(\mathbf{t}) + O(\mathbf{t})$, respectively, and whose western side follows the vertical zigzag lattice path with $E(\mathbf{t}) - \frac{1}{2}$ steps (i.e. the western side has $E(\mathbf{t}) - 1$ and a half 'bumps'). Next, we also remove the triangles of sides t_{2i} 's from the base so that the

distances between two consecutive ones are t_{2i-1} 's. Denote by $\mathcal{K}(\mathbf{t}) = \mathcal{K}(t_1, t_2, \dots, t_{2l})$ the resulting regions (see the regions in Figure 3(d) for the case when $t_1 > 0$ and Figure 3(e) for the case when $t_1 = 0$). Similar to the case of \mathcal{Q}' -type regions, we also define a weighted version $\mathcal{K}'(\mathbf{t})$ of the $\mathcal{K}(\mathbf{t})$ by assigning to each vertical lozenge on its western side a weight $\frac{1}{2}$ (see Figure 3(f)).

We adopt here the notations $Q(\mathbf{t})$, $Q'(\mathbf{t})$, $K(\mathbf{t})$, and $K'(\mathbf{t})$ from [9] for the numbers of tilings of the regions $\mathcal{Q}(\mathbf{t})$, $\mathcal{Q}'(\mathbf{t})$, $\mathcal{K}(\mathbf{t})$, and $\mathcal{K}'(\mathbf{t})$, respectively.

We define the *hyperfactorial* $H(n)$ by

$$H(n) := 0! \cdot 1! \cdot 2! \cdots (n-1)!,$$

and the '*skipping*' hyperfactorial $H_2(n)$ by

$$H_2(n) = \begin{cases} 0! \cdot 2! \cdot 4! \cdots (n-2)! & \text{if } n \text{ is even;} \\ 1! \cdot 2! \cdot 3! \cdots (n-2)! & \text{if } n \text{ is odd.} \end{cases}$$

Lemma 3. [9, Lemma 1.4] For any sequence of non-negative integers $\mathbf{t} = (t_1, t_2, \dots, t_{2l})$

$$\begin{aligned} Q(\mathbf{t}) &= \frac{\prod_{i=1}^l \frac{(s_{2i}(\mathbf{t}))!}{(s_{2i-1}(\mathbf{t}))!}}{H_2(2E(\mathbf{t})+1)} \prod_{i=1}^l H_2(2s_{2i}(\mathbf{t})+1) H_2(2s_{2i-1}(\mathbf{t})+2) \\ &\times \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ odd}}} \frac{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}{H(s_j(\mathbf{t}) + s_i(\mathbf{t}) + 1)} \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ even}}} \frac{H(s_j(\mathbf{t}) + s_i(\mathbf{t}) + 1)}{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}, \end{aligned} \quad (9)$$

$$\begin{aligned} Q'(\mathbf{t}) &= \frac{2^{-E(\mathbf{t})}}{H_2(2E(\mathbf{t})+1)} \prod_{i=1}^l H_2(2s_{2i}(\mathbf{t})+1) H_2(2s_{2i-1}(\mathbf{t})) \\ &\times \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ odd}}} \frac{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}{H(s_j(\mathbf{t}) + s_i(\mathbf{t}))} \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ even}}} \frac{H(s_j(\mathbf{t}) + s_i(\mathbf{t}))}{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}, \end{aligned} \quad (10)$$

$$\begin{aligned} K(\mathbf{t}) &= \frac{1}{H_2(2E(\mathbf{t}))} \prod_{i=1}^l H_2(2s_{2i}(\mathbf{t})) H_2(2s_{2i-1}(\mathbf{t})+1) \\ &\times \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ odd}}} \frac{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}{H(s_j(\mathbf{t}) + s_i(\mathbf{t}))} \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ even}}} \frac{H(s_j(\mathbf{t}) + s_i(\mathbf{t}))}{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}, \end{aligned} \quad (11)$$

and

$$\begin{aligned} K'(\mathbf{t}) &= \frac{1}{H_2(2E(\mathbf{t}))} \prod_{i=1}^l H_2(2s_{2i}(\mathbf{t})-1) H_2(2s_{2i-1}(\mathbf{t})) \\ &\times \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ odd}}} \frac{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}{H(s_j(\mathbf{t}) + s_i(\mathbf{t}) - 1)} \prod_{\substack{1 \leq i < j \leq 2l \\ j-i \text{ even}}} \frac{H(s_j(\mathbf{t}) + s_i(\mathbf{t}) - 1)}{H(s_j(\mathbf{t}) - s_i(\mathbf{t}))}. \end{aligned} \quad (12)$$




Figure 4: The halved hexagons with an array of holes on the staircase boundary: (a) $H_{3,3,2}^{(1)}(2, 2, 2, 3, 2, 2)$ and (b) $H_{3,3,3}^{(2)}(2, 3, 2, 2, 2, 2)$.

We now go to the definitions of our 8 halved hexagons.

Assume that x, y, z are non-negative integers and that $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers (\mathbf{a} may be an empty sequence). We consider a pentagonal region whose northern, northeastern, southeastern, and south sides have respectively lengths $x + E(\mathbf{a}), y + z + 2O(\mathbf{a}), y + z + 2E(\mathbf{a}), x + O(\mathbf{a})$, and the western side follows a vertical zigzag lattice path with $y + z + E(\mathbf{a}) + O(\mathbf{a})$ steps. Next, we remove an array of alternating up-pointing and down-pointing triangles at level $2z + 2E(\mathbf{a})$ from the bottom. In particular, the array of triangles starts with an up-pointing half triangle of side $2a_1$, then the next triangles have sides a_2, a_3, \dots, a_k , ordered from left to right (see Figure 4(a) for the case $x = 3, y = 3, z = 2, k = 6, a_1 = 2, a_2 = 2, a_3 = 2, a_4 = 3, a_5 = 2, a_6 = 2$). Denote by $H_{x,y,z}^{(1)}(\mathbf{a}) = H_{x,y,z}^{(1)}(a_1, a_2, \dots, a_k)$ the resulting region.

Next, we investigate a variation $H_{x,y,z}^{(2)}(\mathbf{a})$ of $H_{x,y,z}^{(1)}(\mathbf{a})$ as follows. We start with a pentagonal region of side-lengths $x + E(\mathbf{a}), y + z - 1 + 2O(\mathbf{a}), y + z - 1 + 2E(\mathbf{a}), x + O(\mathbf{a}), y + z - 1 + E(\mathbf{a}) + O(\mathbf{a})$ in the same order as in the definition of the $H^{(1)}$ -type region. The array of triangles is now removed at an odd level, $2z + 2E(\mathbf{a}) - 1$, from the bottom (instead of the level $2z + 2E(\mathbf{a})$ in the case of the $H^{(1)}$ -type regions). Figure 4(b) illustrates the region for the case $x = 3, y = 3, z = 3, k = 6, a_1 = 2, a_2 = 3, a_3 = 2, a_4 = 2, a_5 = 2, a_6 = 2$. Lozenge tilings of the two ‘defected’ halved hexagons $H_{x,y,z}^{(1)}(\mathbf{a})$ and $H_{x,y,z}^{(2)}(\mathbf{a})$ are always enumerated by simple product formulas.

Theorem 4. Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(1)}(a_1, a_2, \dots, a_k)$ is given by

$$\begin{aligned} M(H_{x,y,z}^{(1)}(a, b)) &= \frac{P_{y,y+2a,b} P_{z+b,z+b,a} Q(a, b, x, y+z)}{P_{y+z+b,y+z+b,a}} \frac{T(x+b+1, y+a-1, a)}{T(b+1, y+a-1, a)} \\ &\times \frac{T(x+z+a+b+2, y+a-1, a) T(2a+b+2, y+b-1, b) T(z+1, y+b-1, b)}{T(z+a+b+2, y+a-1, a) T(x+2a+b+2, y+b-1, b) T(x+z+1, y+b-1, b)} \end{aligned} \quad (13)$$

for $k \geq 2$

$$\begin{aligned} M(H_{x,y,z}^{(1)}(a_1, a_2, \dots, a_{2k})) &= M(H_{x,y,z}^{(1)}(O(\mathbf{a}), E(\mathbf{a}))) \frac{Q(0, a_1, \dots, a_{2k}, y) Q(a_1, \dots, a_{2k} + z)}{P_{y,y+2O(\mathbf{a}),E(\mathbf{a})} P_{z+E(\mathbf{a}),z+E(\mathbf{a}),O(\mathbf{a})}} \\ &\times \prod_{i=2}^k \frac{T(x+z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(x+y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}, \end{aligned} \quad (14)$$

and

$$M(H_{x,y,z}^{(1)}(a_1, a_2, \dots, a_{2k-1})) = M(H_{x,y,z}^{(1)}(a_1, a_2, \dots, a_{2k-1}, 0)). \quad (15)$$

We can view a halved hexagon with an odd number of triangular holes as a special case of the one with an even number of holes, when the rightmost hole has size 0, as illustrated in the equation (15). For the sake of simplicity, in our next theorems (Theorems 5–11), we only show the tiling formulas of halved hexagons with an even number of holes.

Theorem 5. Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(2)}(a_1, a_2, \dots, a_k)$ is given by

$$\begin{aligned} M(H_{x,y,z}^{(2)}(a, b)) &= \frac{P_{y,y+2a,b} P_{z+b-1,z+b-1,a} K(a, b, x, y+z)}{P_{y+z+b-1,y+z+b-1,a}} \frac{T(x+b+1, y+a-1, a)}{T(b+1, y+a-1, a)} \\ &\times \frac{T(x+z+a+b+1, y+a-1, a) T(2a+b+1, y+b-1, b) T(z+1, y+b-1, b)}{T(z+a+b+1, y+a-1, a) T(x+2a+b+1, y+b-1, b) T(x+z+1, y+b-1, b)}, \end{aligned} \quad (16)$$

and for $k \geq 2$

$$\begin{aligned}
 M(H_{x,y,z}^{(2)}(a_1, a_2, \dots, a_{2k})) &= M(H_{x,y,z}^{(2)}(O(\mathbf{a}), E(\mathbf{a}))) \frac{K(0, a_1, \dots, a_{2k}, y) K(a_1, \dots, a_{2k} + z)}{P_{y,y+2O(\mathbf{a}),E(\mathbf{a})} P_{z+E(\mathbf{a})-1,z+E(\mathbf{a})-1,O(\mathbf{a})}} \\
 &\times \prod_{i=2}^k \frac{T(x + z + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(x + y + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))} \\
 &\times \prod_{i=2}^k \frac{T(y + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(z + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))} \\
 &\times \prod_{i=2}^k \frac{T(x + y + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(x + z + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))} \\
 &\times \prod_{i=2}^k \frac{T(z + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(y + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}. \tag{17}
 \end{aligned}$$




Figure 5: The weighted halved hexagons with an array of holes on the staircase boundary: (a) $H_{3,3,2}^{(3)}(2, 2, 2, 3, 2, 2)$ and (b) $H_{3,3,3}^{(4)}(2, 3, 2, 2, 2, 2)$. The lozenges with shaded cores are weighted by $1/2$.

Similar to the case of original halved hexagons $\mathcal{P}_{a,b,c}$, we are interested in the weighted versions $H_{x,y,z}^{(3)}(\mathbf{a})$ and $H_{x,y,z}^{(4)}(\mathbf{a})$ of the above regions $H_{x,y,z}^{(1)}(\mathbf{a})$ and $H_{x,y,z}^{(2)}(\mathbf{a})$, respectively,

where the vertical lozenges along the western side are weighted by $1/2$ (see Figures 5(a) and (b), respectively; the lozenges with shaded cores are weighted by $1/2$). The tiling numbers of these weighted regions are also given by simple product formulas.

Theorem 6. *Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(3)}(a_1, a_2, \dots, a_k)$ is given by*

$$\begin{aligned} M(H_{x,y,z}^{(3)}(a, b)) &= \frac{P'_{y,y+2a,b} P'_{z+b,z+b,a} Q'(0, a, b, x, y+z)}{P'_{y+z+b,y+z+b,a}} \frac{T(x+b+1, y+a-1, a)}{T(b+1, y+a-1, a)} \\ &\times \frac{T(x+z+a+b+1, y+a-1, a) T(2a+b+1, y+b-1, b) T(z+1, y+b-1, b)}{T(z+a+b+1, y+a-1, a) T(x+2a+b+1, y+b-1, b) T(x+z+1, y+b-1, b)}, \end{aligned} \quad (18)$$

and for $k \geq 2$

$$\begin{aligned} M(H_{x,y,z}^{(3)}(a_1, a_2, \dots, a_{2k})) &= 2^{a_1} M(H_{x,y,z}^{(3)}(O(\mathbf{a}), E(\mathbf{a}))) \\ &\times \frac{Q'(a_1, a_2, \dots, a_{2k}, z) Q'(0, a_1, \dots, a_{2k} + z)}{P'_{y,y+2O(\mathbf{a}),E(\mathbf{a})} P'_{z+E(\mathbf{a}),z+E(\mathbf{a}),O(\mathbf{a})}} \\ &\times \prod_{i=2}^k \frac{T(x+z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(x+y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}. \end{aligned} \quad (19)$$

Theorem 7. *Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(4)}(a_1, a_2, \dots, a_k)$ is given by*

$$\begin{aligned} M(H_{x,y,z}^{(4)}(a, b)) &= \frac{P'_{y,y+2a,b} P'_{z+b-1,z+b-1,a} K'(a, b, x, y+z)}{P'_{y+z+b-1,y+z+b-1,a}} \frac{T(x+b+1, y+a-1, a)}{T(b+1, y+a-1, a)} \\ &\times \frac{T(x+z+a+b, y+a-1, a) T(2a+b, y+b-1, b) T(z+1, y+b-1, b)}{T(z+a+b, y+a-1, a) T(x+2a+b, y+b-1, b) T(x+z+1, y+b-1, b)}, \end{aligned} \quad (20)$$

and for $k \geq 2$

$$\begin{aligned}
M(H_{x,y,z}^{(4)}(a_1, a_2, \dots, a_{2k})) &= 2^{a_1-1} M(H_{x,y,z}^{(4)}(O(\mathbf{a}), E(\mathbf{a}))) \\
&\times \frac{K'(0, a_1, \dots, a_{2k}, y) K'(a_1, \dots, a_{2k} + z)}{P'_{y,y+2O(\mathbf{a}),E(\mathbf{a})} P'_{z+E(\mathbf{a})-1,z+E(\mathbf{a})-1,O(\mathbf{a})}} \\
&\times \prod_{i=2}^k \frac{T(x + z + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(x + y + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(y + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(z + \mathbf{e}_i(\mathbf{a}) + 1, a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(x + y + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}), a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(x + z + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}), a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(z + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}), a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}{T(y + \mathbf{s}_{2i-1}(\mathbf{a}) + \mathbf{s}_{2k}(\mathbf{a}), a_{2i-2} + \mathbf{o}_i(\mathbf{a}) - 1, \mathbf{o}_i(\mathbf{a}))}. \tag{21}
\end{aligned}$$

Finally, we focus on four new families of defected halved hexagons with ‘mixed’ western boundary. In these families, only a half of the lozenges along the western boundary (the ones above the array of holes or the ones below the array of holes) are weighted by $1/2$.

The first ‘mixed boundary’ region $H_{x,y,z}^{(5)}(a_1, a_2, \dots, a_k)$ is obtained from the weighted region $H_{x,y,z+1}^{(4)}(a_1+1, a_2, \dots, a_k)$ by removing all unit triangles running along the southern and the southeastern sides, as well as the ones running along the portion of the western boundary below the array of holes (see Figure 6(a) for the case $x = 3, y = 3, z = 2, k = 6, a_1 = 2, a_2 = 2, a_3 = 2, a_4 = 3, a_5 = 2, a_6 = 2$; the dotted triangles indicate the triangles removed; the lozenges with shaded cores are weighted by $1/2$ as usual). We can see that only the lozenges staying above the array of holes and running along the western boundary side are weighted in the region $H_{x,y,z}^{(5)}(a_1, \dots, a_k)$. The second mixed boundary region $H_{x,y,z}^{(6)}(a_1, a_2, \dots, a_k)$ is created from $H_{x,y,z}^{(3)}(a_1, a_2, \dots, a_k)$ by removing all unit triangles running along the southern side and the portion of the western side below the array of holes (see Figure 6(b) for $x = 3, y = 3, z = 3, k = 6, a_1 = 2, a_2 = 3, a_3 = 2, a_4 = 2, a_5 = 2, a_6 = 2$).

Similarly, the region $H_{x,y,z}^{(7)}(a_1, a_2, \dots, a_k)$ is obtained from the region $H_{x,y,z}^{(3)}(a_1, a_2, \dots, a_k)$ by removing all unit triangles running along the *northern* side and the western side *above* the holes (shown in Figure 6(c) when $x = 3, y = 3, z = 2, k = 6, a_1 = 2, a_2 = 2, a_3 = 2, a_4 = 3, a_5 = 2, a_6 = 2$). Finally, the region $H_{x,y,z}^{(8)}(a_1, a_2, \dots, a_k)$ is obtained by removing the unit triangles running along the *northern* side and the western side *above* the holes in $H_{x,y,z}^{(4)}(a_1, a_2, \dots, a_k)$ (illustrated in Figure 6(d) for $x = 3, y = 3, z = 3, k = 6, a_1 = 2, a_2 = 3, a_3 = 2, a_4 = 2, a_5 = 2, a_6 = 2$).

The tiling numbers of the above four regions with mixed boundary are all given by simple product formulas. These formulas are similar to that in Theorems 4–7 above, except for the appearances of the product V .




Figure 6: The four mixed boundary halved hexagons: (a) $H_{3,3,2}^{(5)}(2, 2, 2, 3, 2, 2)$, (b) $H_{3,3,3}^{(6)}(2, 3, 2, 2, 2, 2)$, (c) $H_{3,3,2}^{(7)}(2, 2, 2, 3, 2, 2)$, and (d) $H_{3,3,3}^{(8)}(2, 3, 2, 2, 2, 2)$.

Theorem 8. Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(5)}(a_1, a_2, \dots, a_k)$ is given by

$$M(H_{x,y,z}^{(5)}(a, b)) = \frac{P'_{y,y+2a+1,b} P_{z+b,z+b,a} Q(a, b, x, y+z)}{P_{y+z+b,y+z+b,a}} \frac{V(2a+2b+3, y+z-1, y)}{V(2x+2a+2b+3, y+z-1, y)} \\ \times \frac{T(x+b+1, y+z+2a, y) T(2a+b+2, y+b-1, b) T(z+1, y+b-1, b)}{T(b+1, y+z+2a, y) T(x+2a+b+2, y+b-1, b) T(x+z+1, y+b-1, b)}, \quad (22)$$

and for $k \geq 2$

$$M(H_{x,y,z}^{(5)}(a_1, a_2, \dots, a_{2k})) = 2^{a_1} M(H_{x,y,z}^{(5)}(O(\mathbf{a}), E(\mathbf{a}))) \\ \times \frac{K'(0, a_1+1, a_2, \dots, a_{2k}, y) Q(a_1, \dots, a_{2k}+z)}{P'_{y,y+2O(\mathbf{a})+1,E(\mathbf{a})} P_{z+E(\mathbf{a}),z+E(\mathbf{a}),O(\mathbf{a})}} \\ \times \prod_{i=2}^k \frac{T(x+z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ \times \prod_{i=2}^k \frac{T(y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ \times \prod_{i=2}^k \frac{T(x+y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ \times \prod_{i=2}^k \frac{T(z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+2, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}. \quad (23)$$

Theorem 9. Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(6)}(a_1, a_2, \dots, a_k)$ is given by

$$M(H_{x,y,z}^{(6)}(a, b)) = \frac{P'_{y,y+2a,b} P_{z+b-1,z+b-1,a} K(a, b, x, y+z)}{P_{y+z+b-1,y+z+b-1,a}} \frac{V(2a+2b+3, y+z-2, y)}{V(2x+2a+2b+3, y+z-2, y)} \\ \times \frac{T(x+b+1, y+z+2a-1, y) T(2a+b+1, y+b-1, b) T(z+1, y+b-1, b)}{T(b+1, y+z+2a-1, y) T(x+2a+b+1, y+b-1, b) T(x+z+1, y+b-1, b)}, \quad (24)$$

and for $k \geq 2$

$$\begin{aligned}
M(H_{x,y,z}^{(6)}(a_1, a_2, \dots, a_{2k})) &= 2^{a_1} M(H_{x,y,z}^{(6)}(O(\mathbf{a}), E(\mathbf{a}))) \\
&\times \frac{Q'(0, a_1, a_2, \dots, a_{2k}, y) K(a_1, \dots, a_{2k} + z)}{P'_{y,y+2O(\mathbf{a}),E(\mathbf{a})} P'_{z+E(\mathbf{a})-1,z+E(\mathbf{a})-1,O(\mathbf{a})}} \\
&\times \prod_{i=2}^k \frac{T(x+z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(x+y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}. \tag{25}
\end{aligned}$$

Theorem 10. Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(7)}(a_1, a_2, \dots, a_k)$ is given by

$$\begin{aligned}
M(H_{x,y,z}^{(7)}(a, b)) &= \frac{P_{y,y+2a-1,b} P'_{z+b,z+b,a} Q'(a, b, x, y+z)}{P'_{y+z+b,y+z+b,a}} \frac{V(2a+2b+1, y+z, y)}{V(2x+2a+2b+1, y+z, y)} \\
&\times \frac{T(x+b+1, y+z+2a-1, y) T(2a+b+1, y+b-1, b) T(z+1, y+b-1, b)}{T(b+1, y+z+2a-1, y) T(x+2a+b+1, y+b-1, b) T(x+z+1, y+b-1, b)}, \tag{26}
\end{aligned}$$

and for $k \geq 2$

$$\begin{aligned}
M(H_{x,y,z}^{(7)}(a_1, a_2, \dots, a_{2k})) &= M(H_{x,y,z}^{(7)}(O(\mathbf{a}), E(\mathbf{a}))) \\
&\times \frac{K(0, a_1, a_2, \dots, a_{2k}, y) Q'(a_1, \dots, a_{2k} + z)}{P_{y,y+2O(\mathbf{a})-1,E(\mathbf{a})} P'_{z+E(\mathbf{a}),z+E(\mathbf{a}),O(\mathbf{a})}} \\
&\times \prod_{i=2}^k \frac{T(x+z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(x+y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\
&\times \prod_{i=2}^k \frac{T(z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}. \tag{27}
\end{aligned}$$

Theorem 11. Assume that x, y, z, k are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_k)$ is a sequence of k non-negative integers. The number of tilings of the defected halved hexagon $H_{x,y,z}^{(8)}(a_1, a_2, \dots, a_k)$ is given by

$$\begin{aligned} M(H_{x,y,z}^{(8)}(a, b)) &= \frac{P_{y,y+2a-2,b} P'_{z+b-1,y+b-1,a} K'(a, b, x, y+z)}{P'_{y+z+b-1,y+z+b-1,a}} \\ &\times \frac{V(2a+2b+1, y+z-1, y) T(x+b+1, y+z+2a-2, y)}{V(2x+2a+2b+1, y+z-1, y) T(b+1, y+z+2a-2, y)} \\ &\times \frac{T(2a+b, y+b-1, b) T(z+1, y+b-1, b)}{T(x+2a+b, y+b-1, b) T(x+z+1, y+b-1, b)}, \end{aligned} \quad (28)$$

and for $k \geq 2$

$$\begin{aligned} M(H_{x,y,z}^{(8)}(a_1, a_2, \dots, a_{2k})) &= M(H_{x,y,z}^{(8)}(O(\mathbf{a}), E(\mathbf{a}))) \\ &\times \frac{Q(0, a_1-1, a_2, \dots, a_{2k}, y) K'(a_1, \dots, a_{2k}+z)}{P_{y,y+2O(\mathbf{a})-2,E(\mathbf{a})} P'_{z+E(\mathbf{a})-1,y+E(\mathbf{a})-1,O(\mathbf{a})}} \\ &\times \prod_{i=2}^k \frac{T(x+z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(y+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(z+e_i(\mathbf{a})+1, a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(x+y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a}), a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(x+z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a}), a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))} \\ &\times \prod_{i=2}^k \frac{T(z+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a}), a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}{T(y+s_{2i-1}(\mathbf{a})+s_{2k}(\mathbf{a}), a_{2i-2}+o_i(\mathbf{a})-1, o_i(\mathbf{a}))}. \end{aligned} \quad (29)$$

Assume that x, y, z are non-negative integers and that $\mathbf{a} = (a_1, a_2, \dots, a_n)$ is a sequence of nonnegative integers as usual. Consider a symmetric hexagon of side-lengths $y+2O(\mathbf{a})-a_1, x+2E(\mathbf{a}), y+2O(\mathbf{a})-a_1, y+2E(\mathbf{a}), x+2E(\mathbf{a}), y+2E(\mathbf{a})$. We remove at level z (from the bottom) a symmetric array of triangles of sides $a_n, a_{n-1}, \dots, a_2, a_1, a_2, \dots, a_{n-1}, a_n$ ordered from left to right, so that the middle triangle is an up-pointing triangle of side a_1 . Denote by $\mathcal{S}_{x,y,z}(\mathbf{a}) = \mathcal{S}_{x,y,z}(a_1, a_2, \dots, a_n)$ the resulting region (see Figure 17 for examples).

We note that, by the symmetry, if the total length of the array, $2(E(\mathbf{a}) + O(\mathbf{a})) - a_1$, and the length of the base of the hexagon, $x + 2O(\mathbf{a}) - a_1$, have the same parity, then z must be even. Thus, z and x always have the same parity.

Theorem 12. Assume that x, y, z are non-negative integers and $\mathbf{a} = (a_1, a_2, \dots, a_n)$ is a sequence of positive integers. If $2E(\mathbf{a}) - 1 \leq z \leq 2y + 2E(\mathbf{a}) + 1$, then the number of tilings of $\mathcal{S}_{x,y,z}(\mathbf{a})$ is always given by a simple product formula as follows.

(1) If x is even (so z is even) and a_1 is even, then

$$\begin{aligned} M(\mathcal{S}_{x,y,z}(\mathbf{a})) &= 2^{y+a_2+a_3+\dots+a_n} M\left(H_{\frac{x}{2}+E(\mathbf{a}),y-\frac{z}{2}+E(\mathbf{a}),\frac{z}{2}-E(\mathbf{a})}^{(2)}\left(\frac{a_1}{2}, a_2, \dots, a_n\right)\right) \\ &\quad \times M\left(H_{\frac{x}{2}+E(\mathbf{a}),y-\frac{z}{2}+E(\mathbf{a}),\frac{z}{2}-E(\mathbf{a})}^{(3)}\left(\frac{a_1}{2}, a_2, \dots, a_n\right)\right). \end{aligned} \quad (30)$$

(2) If x is odd (so z is odd) and a_1 is even, then

$$\begin{aligned} M(\mathcal{S}_{x,y,z}(\mathbf{a})) &= 2^{y+a_2+a_3+\dots+a_n} M\left(H_{\frac{x-1}{2}+E(\mathbf{a}),y-\frac{z}{2}+E(\mathbf{a}),\frac{z}{2}-E(\mathbf{a})+1}^{(2)}\left(\frac{a_1}{2}, a_2, \dots, a_n\right)\right) \\ &\quad \times M\left(H_{\frac{x+1}{2}+E(\mathbf{a}),y-\frac{z-1}{2}+E(\mathbf{a})-1,\frac{z-1}{2}-E(\mathbf{a})}^{(3)}\left(\frac{a_1}{2}, a_2, \dots, a_n\right)\right). \end{aligned} \quad (31)$$

(3) If x is even (so z is even) and a_1 is odd, then

$$\begin{aligned} M(\mathcal{S}_{x,y,z}(\mathbf{a})) &= 2^{y+a_2+a_3+\dots+a_n} M\left(H_{\frac{x}{2}+E(\mathbf{a}),y-\frac{z}{2}+E(\mathbf{a}),\frac{z}{2}-E(\mathbf{a})}^{(5)}\left(\frac{a_1-1}{2}, a_2, \dots, a_n\right)\right) \\ &\quad \times M\left(H_{\frac{x}{2}+E(\mathbf{a}),y-\frac{z}{2}+E(\mathbf{a}),\frac{z}{2}-E(\mathbf{a})}^{(8)}\left(\frac{a_1+1}{2}, a_2, \dots, a_n\right)\right). \end{aligned} \quad (32)$$

(4) If x is odd (so z is odd) and a_1 is odd, then

$$\begin{aligned} M(\mathcal{S}_{x,y,z}(\mathbf{a})) &= 2^{y+a_2+a_3+\dots+a_n} \\ &\quad \times M\left(H_{\frac{x+1}{2}+E(\mathbf{a}),y-\frac{z-1}{2}+E(\mathbf{a})-1,\frac{z-1}{2}-E(\mathbf{a})}^{(5)}\left(\frac{a_1-1}{2}, a_2, \dots, a_n\right)\right) \\ &\quad \times M\left(H_{\frac{x-1}{2}+E(\mathbf{a}),y-\frac{z-1}{2}+E(\mathbf{a}),\frac{z-1}{2}-E(\mathbf{a})+1}^{(8)}\left(\frac{a_1+1}{2}, a_2, \dots, a_n\right)\right). \end{aligned} \quad (33)$$

Finally, we note that when $z < 2E(\mathbf{a}) - 1$ or $z > 2y + 2E(\mathbf{a}) + 1$, then $M(\mathcal{S}_{x,y,z}(\mathbf{a})) = 0$. This will be explained later in Remark 23 of Section 4.

3 Preliminaries

Let G be a simple graph without loops. A *perfect matching* (or simple *matching* in this paper) of G is a collection of vertex-disjoint edges that cover all vertices of G . We use the notation $M(G)$ for the number of matchings of G . The tilings of a region R can be identified with the matchings of its *dual graph* G (the graph whose vertices are unit triangles in R and whose edges connect precisely two unit triangles sharing an edge). In the weighted case, each edge of the dual graph G has the same weight as the corresponding lozenge in the region R . In this case, $M(G)$ denotes the weighted number of matchings of G , i.e. the sum of weights of all matchings in G , where the weight of a matching is the product of all weight of its constituent edges.

If a region admits a tiling, then it has the same number of up- and down-pointing unit triangles. We call the regions satisfying the latter balancing condition *balanced* regions.

Lemma 13 (Region-splitting Lemma). *Let R be a balanced region. Assume that a sub-region S of R satisfies the following two conditions:*

- (i) (Separating Condition) *All unit triangles in S that are adjacent to $R - S$ are of the same type (up-pointing or down-pointing).*
- (ii) (Balancing Condition) *S is balanced.*

Then

$$M(R) = M(S) M(R - S). \quad (34)$$

Proof. ¹ Assume there is a tiling of R which contains boundary-crossing lozenges between S and $R - S$ (i.e., lozenges which consist of a unit triangle from the boundary of S and a unit triangle from the boundary of $R - S$). Since there is only one type of unit triangle on each side of the boundary between S and $R - S$, and since S and $R - S$ are balanced, the regions obtained by removing such boundary-crossing lozenges would no longer be balanced, and hence would have no tilings. Therefore, there can not be any boundary-crossing lozenges, and S and $R - S$ must be tiled independently, giving the factorization (34). \square

One of the main ingredients of our proofs is the following powerful theorem by Eric H. Kuo [6] that is usually mentioned as *Kuo condensation*.

Theorem 14 (Theorem 5.1 in [6]). *Assume that $G = (V_1, V_2, E)$ is a weighted bipartite planar graph with two vertex classes V_1 and V_2 of the same cardinality. Assume in addition that u, v, w, s are four vertices appearing on a cyclic order on a face of G , such that $u, w \in V_1$ and $v, s \in V_2$. Then*

$$M(G) M(G - \{u, v, w, s\}) = M(G - \{u, v\}) M(G - \{w, s\}) + M(G - \{u, s\}) M(G - \{v, w\}). \quad (35)$$

The next lemma is often called *Ciucu's factorization theorem* (Theorem 1.2 in [1]), that allows us write the number of matchings of a symmetric graph as the product of the matching numbers of two disjoint subgraphs.

Lemma 15 (Ciucu's Factorization Theorem). *Let $G = (V_1, V_2, E)$ be a weighted bipartite planar graph with a vertical symmetry axis ℓ . Assume that $a_1, b_1, a_2, b_2, \dots, a_k, b_k$ are all the vertices of G on ℓ appearing in this order from top to bottom². Assume in addition that the vertices of G on ℓ form a cut set of G (i.e. the removal of those vertices separates G into two vertex-disjoint graphs). We reduce the weights of all edges of G lying on ℓ by half and keep the other edge-weights unchanged. Next, we color the two vertex-classes V_1 and V_2 of G by black and white, without loss of generality, assume that a_1 is black. Finally, we remove all edges on the left of ℓ which are adjacent to a black a_i or a white b_j ; we also remove the edges on the right of ℓ which are adjacent to a white a_i or a black*

¹This proof was provided by an anonymous referee for the paper [10].

²It is easy to see that if G admits a perfect matching, then G has an even number of vertices on ℓ .




Figure 7: Ciucu's Factorization Theorem. The edges cut off are illustrated by dotted edges.

b_j . This way, G is divided into two disjoint weighted graphs G^+ and G^- (on the left and on the right of ℓ , respectively). See Figure 7 for an example. Then

$$M(G) = 2^k M(G^+) M(G^-). \quad (36)$$

4 Proof of the main theorems

We first prove Theorem 4.

Proof of Theorem 4. We first prove (13) by induction on $x + y + z$. The base cases are the situations when at least one of the parameters x, y, z is equal to 0.

If $x = 0$, then we split the region $H_{0,y,z}^{(1)}(a, b)$ into two (balanced) subregions along the right side of the a -hole: the lower subregion is the halved hexagon $\mathcal{P}_{b+z, b+z, a}$, and the upper subregion is a union of the halved hexagon $\mathcal{P}_{y, y+a_1, a_2}$ and several forced vertical lozenges (see Figure 8(a)). By Region-splitting Lemma 13, we get

$$M(H_{0,y,z}^{(1)}(a, b)) = M(\mathcal{P}_{b+z, b+z, a}) M(\mathcal{P}_{y, y+a, b}), \quad (37)$$

and (13) follows from Proctor's Theorem 1.

If $y = 0$, then our region has several forced lozenges on the top as in Figure 9(a). By removing these forced lozenges, we obtained an upside-down region $\mathcal{Q}(a, b, x, z)$. Then (13) follows from Lemma 3 in this case.

If $z = 0$, similar to the case when $x = 0$, we also split the region into two halved hexagons and several forced lozenges as in Figure 10(a). Thus, our formula is also implied by Proctor's Theorem 1.




Figure 8: Splitting up a $H^{(1)}$ -type region into two Q -type regions in the case of $x = 0$.




Figure 9: Partitioning a $H^{(1)}$ -type region into two Q -type regions in the case of $y = 0$.




Figure 10: The $H^{(1)}$ -type region can be divided into two Q -type subregions when $z = 0$.




Figure 11: How to apply Kuo condensation to a $H^{(1)}$ -region.




Figure 12: Obtaining a recurrence for the number of tilings of $H^{(1)}$ -type region.

For the induction step, we assume x, y, z are all positive and that (13) holds for any $H^{(1)}$ -type region with two holes whose sum of the x -, y - and z -parameters is strictly less than $x + y + z$. We need to verify (13) for the region $H_{x,y,z}^{(1)}(a, b)$.

We apply Kuo condensation to the dual graph G of the region $H_{x,y,z}^{(1)}(\mathbf{a})$, for a general sequence of nonnegative integers $\mathbf{a} = (a_1, a_2, \dots, a_n)$ with the four vertices u, v, w, s chosen as in Figure 11, each of the four vertices u, v, w, s corresponds to the black unit triangle of the same label. In particular, the u - and v -triangles are respectively the up-pointing and down-pointing black unit triangles at the upper-right corner of the region, and the w - and s -triangles are the up-pointing and down-pointing black unit triangles at the lower-right corner of the region.

Consider the region corresponding to the graph $G - \{u, v, w, s\}$. The removal of the black unit triangles yields several forced lozenges. By removing these forced lozenges, we

get the region $H_{x,y-1,z-1}^{(1)}(\mathbf{a})$ (see the region restricted by the bold contour in Figure 12(a)) and obtain

$$M(G - \{u, v, w, s\}) = M(H_{x,y-1,z-1}^{(1)}(\mathbf{a})). \quad (38)$$

Similarly, we have

$$M(G - \{u, v\}) = M(H_{x,y-1,z}^{(1)}(\mathbf{a})), \quad (39)$$

$$M(G - \{w, s\}) = M(H_{x,y,z-1}^{(1)}(\mathbf{a})), \quad (40)$$

$$M(G - \{u, s\}) = M(H_{x+1,y-1,z-1}^{(1)}(\mathbf{a})), \quad (41)$$

and

$$M(G - \{v, w\}) = M(H_{x-1,y,z}^{(1)}(\mathbf{a})) \quad (42)$$

(see Figures 12(b)–(e), respectively). Plugging the above five identities into the equation (35) in Kuo’s Theorem 14, we get the following recurrence:

$$\begin{aligned} M(H_{x,y,z}^{(1)}(\mathbf{a})) M(H_{x,y-1,z-1}^{(1)}(\mathbf{a})) &= M(H_{x,y-1,z}^{(1)}(\mathbf{a})) M(H_{x,y,z-1}^{(1)}(\mathbf{a})) \\ &\quad + M(H_{x+1,y-1,z-1}^{(1)}(\mathbf{a})) M(H_{x-1,y,z}^{(1)}(\mathbf{a})), \end{aligned} \quad (43)$$

for any sequence of positive integers $\mathbf{a} = (a_1, a_2, \dots, a_n)$. In particular, the number tilings of the region $H_{x,y,z}^{(1)}(a, b)$ satisfies this recurrence (by setting $n = 2$, $a_1 = a$, $a_2 = b$). To finish the proof of (13) we only need to verify that the expression on the right-hand side of (13) satisfies the same recurrence (43). Indeed, denote by $\phi_{x,y,z}(a, b)$ this expression. We need to verify that

$$\phi_{x,y,z}(a, b) \phi_{x,y-1,z-1}(a, b) = \phi_{x,y-1,z}(a, b) \phi_{x,y,z-1}(a, b) + \phi_{x+1,y-1,z-1}(a, b) \phi_{x-1,y,z}(a, b). \quad (44)$$

It is equivalent to show that

$$\frac{\phi_{x,y-1,z}(a, b)}{\phi_{x,y-1,z-1}(a, b)} \frac{\phi_{x,y,z-1}(a, b)}{\phi_{x,y,z}(a, b)} + \frac{\phi_{x+1,y-1,z-1}(a, b)}{\phi_{x,y-1,z-1}(a, b)} \frac{\phi_{x-1,y,z}(a, b)}{\phi_{x,y,z}(a, b)} = 1. \quad (45)$$

We have several claims that are direct consequence of Theorem 1, the definition of the product T , and Lemma 3:

Claim 16.

$$\frac{P_{x,x,a}}{P_{x-1,x-1,a}} = \frac{a+x}{x} \frac{(2a+2x-1)!}{(2a+x)!} \frac{x!}{(2x-1)!}. \quad (46)$$

Claim 17.

$$\frac{T(x, n, m)}{T(x-1, n, m)} = \frac{(x+n-m)_m}{(x-1)_m}. \quad (47)$$

Claim 18. For any sequence $\mathbf{t} = (t_1, t_2, \dots, t_{2l})$

$$\frac{Q(t_1, \dots, t_{2l} + 1)}{Q(t_1, \dots, t_{2l})} = \frac{(s_{2l}(\mathbf{t}) + 1)(2s_{2l}(\mathbf{t}) + 1)!}{(2E(\mathbf{t}) + 1)!} \times \prod_{i=1}^l \frac{(s_{2l}(\mathbf{t}) - s_{2i-1}(\mathbf{t}))!}{(s_{2l}(\mathbf{t}) + s_{2i-1}(\mathbf{t}) + 1)!} \prod_{i=1}^{l-1} \frac{(s_{2l}(\mathbf{t}) + s_{2i}(\mathbf{t}) + 1)!}{(s_{2l}(\mathbf{t}) - s_{2i}(\mathbf{t}))!}. \quad (48)$$

Claim 19. For any sequence $\mathbf{t} = (t_1, t_2, t_3, t_4)$

$$\frac{Q(t_1, t_2, t_3, t_4)}{Q(t_1, t_2, t_3 - 1, t_4)} = \frac{s_4(\mathbf{t})}{s_3(\mathbf{t})} (2s_4(\mathbf{t}) - 1)! (2s_3(\mathbf{t}))! \frac{(s_4(\mathbf{t}) - s_1(\mathbf{t}) - 1)!}{(s_3(\mathbf{t}) - s_1(\mathbf{t}) - 1)!} \times \frac{(s_3(\mathbf{t}) - s_2(\mathbf{t}) - 1)! (s_4(\mathbf{t}) + s_2(\mathbf{t}))! (s_3(\mathbf{t}) + s_1(\mathbf{t}))!}{(s_4(\mathbf{t}) + s_1(\mathbf{t}))! (s_3(\mathbf{t}) + s_2(\mathbf{t}))! (s_4(\mathbf{t}) + s_3(\mathbf{t}) - 1)! (s_4(\mathbf{t}) + s_3(\mathbf{t}))!}. \quad (49)$$

We now simplify the first fraction of the first term on the left-hand side of (45) by using the above claim as

$$\begin{aligned} \frac{\phi(x, y - 1, z)}{\phi(x, y - 1, z - 1)} &= \frac{\Gamma(x + z + a + b + 2, y + a - 2, a)}{\Gamma(x + z + a + b + 1, y + a - 2, a)} \frac{\Gamma(z + 1, y + b - 2, b)}{\Gamma(z, y + b - 2, b)} \\ &\times \frac{\Gamma(z + a + b + 1, y + a - 2, a)}{\Gamma(z + a + b + 2, y + a - 2, a)} \frac{\Gamma(x + z, y + b - 2, b)}{\Gamma(x + z + 1, y + b - 2, b)} \\ &\times \frac{Q(a, b, x, y + z - 1)}{Q(a, b, x, y + z - 2)} \frac{P_{z+b, z+b, a}}{P_{z+b-1, z+b-1, a}} \frac{P_{y+z+b-2, y+z+b-2, a}}{P_{y+z+b-1, y+z+b-1, a}} \\ &= \frac{(x + y + z + a + b)_a (y + z - 1)_b (z + a + b + 1)_a (x + z)_b}{(x + z + a + b + 1)_a (z)_b (y + z + a + b)_a (x + y + z - 1)_b} \\ &\times \frac{(x + y + z + a + b - 1)(2x + 2y + 2z + 2a + 2b - 3)!}{(2y + 2z + 2b - 3)!} \\ &\times \frac{(y + z - 2)!(x + y + z + b - 2)!(x + y + z + 2a + 2b - 1)!}{(x + y + z - 2)!(x + y + z + 2a + b - 1)!(2x + y + z + 2a + 2b - 1)!} \\ &\times \frac{a + b + z (2z + 2a + 2b - 1)! (b + z)!}{b + z (z + b + 2a)! (2z + 2b - 1)!} \\ &\times \frac{y + z + b - 1 (y + z + 2a + b - 1)! (2y + 2z + 2b - 3)!}{y + z + a + b - 1 (2y + 2z + 2a + 2b - 3)! (y + z + b - 1)!}. \end{aligned} \quad (50)$$

Working similarly for the second fraction, $\frac{\phi_{x, y, z-1}(a, b)}{\phi_{x, y, z}(a, b)}$, of the first term and multiplying the result by the above simplified form of the first fraction, we obtain

$$\frac{\phi_{x, y-1, z}(a, b)}{\phi_{x, y-1, z-1}(a, b)} \frac{\phi_{x, y, z-1}(a, b)}{\phi_{x, y, z}(a, b)} = \frac{(2x + y + z + 2a + 2b)(2y + 2z + 2a + 2b - 1)}{(2x + 2y + 2z + 2a + 2b - 1)(x + y + z + 2a + 2b)}. \quad (52)$$

Next, we work on the second term on the left-hand side of (45). The first fraction can be written as

$$\begin{aligned}
\frac{\phi_{x-1,y,z}(a,b)}{\phi_{x,y,z}(a,b)} &= \frac{T(x+b,y+a-1,a)}{T(x+b+1,y+a-1,a)} \frac{T(x+z+a+b+1,y+a-1,a)}{T(x+z+a+b+2,y+a-1,a)} \\
&\times \frac{T(x+2a+b+2,y+b-1,b)}{T(x+2a+b+1,y+b-1,b)} \frac{T(x+z+1,y+b-1,b)}{T(x+z,y+b-1,b)} \frac{Q(a,b,x-1,y+z)}{Q(a,b,x,y+z)} \\
&= \frac{(x+b)_a}{(x+y+b)_a} \frac{(x+z+a+b+1)_a}{(x+y+z+a+b+1)_a} \frac{(x+y+2a+b+1)_b}{(x+2a+b+1)_b} \frac{(x+y+z)_b}{(x+z)_b} \\
&\times \frac{(x+a+b)}{(x+y+z+a+b)(2x+2y+2z+2a+2b-1)!(2x+2a+2b)!} \\
&\times \frac{(x+y+z-1)!(x+b-1)!}{(x+y+z+b-1)!(x-1)!} \\
&\times \frac{(x+y+z+2a+b)!(x+2a+2b)!(2a+2b+2x+y+z-1)!(2a+2b+2x+y+z)!}{(x+y+z+2a+2b)!(x+2a+b)!}.
\end{aligned} \tag{53}$$

Working similarly for the second fraction $\frac{\phi_{x-1,y,z}(a,b)}{\phi_{x,y,z}(a,b)}$ and multiplying by the above simplification of the first one, we get

$$\frac{\phi_{x+1,y-1,z-1}(a,b)}{\phi_{x,y-1,z-1}(a,b)} \frac{\phi_{x-1,y,z}(a,b)}{\phi_{x,y,z}(a,b)} = \frac{x(2x+2a+2b+1)}{(2x+2y+2z+2a+2b-1)(x+y+z+2a+2b)}. \tag{55}$$

We have now

$$\begin{aligned}
\frac{\phi_{x,y-1,z}(a,b)}{\phi_{x,y-1,z-1}(a,b)} \frac{\phi_{x,y,z-1}(a,b)}{\phi_{x,y,z}(a,b)} &+ \frac{\phi_{x+1,y-1,z-1}(a,b)}{\phi_{x,y-1,z-1}(a,b)} \frac{\phi_{x-1,y,z}(a,b)}{\phi_{x,y,z}(a,b)} \\
&= \frac{(2x+y+z+2a+2b)(2y+2z+2a+2b-1)}{(2x+2y+2z+2a+2b-1)(x+y+z+2a+2b)} \\
&+ \frac{x(2x+2a+2b+1)}{(2x+2y+2z+2a+2b-1)(x+y+z+2a+2b)} = 1.
\end{aligned} \tag{56}$$

This finishes the proof of (13).

We now prove (14) by induction on $x+y+z$. The three base cases here are still the cases of $x=0$, $y=0$, and $z=0$.

Similar to the case when $k=1$ above, when $x=0$, we can partition our region into two \mathcal{Q} -type regions and several forced lozenges (shown in Figure 8(b)). By Region-splitting Lemma 13, we have

$$M(H_{0,y,z}^{(1)}(a_1, \dots, a_{2k})) = M(\mathcal{Q}(0, a_1, a_2, \dots, a_{2k}, y)) M(\mathcal{Q}(a_1, a_2, \dots, a_{2k} + z)). \tag{57}$$

Our tiling formula (14) follows from (13) and Lemma 3.

If $y = 0$, we partition our regions into two \mathcal{Q} -type regions (and several forced lozenges) along the line containing the bases of the triangular holes (see Figure 9(b)). By Region-splitting Lemma, we can also write the number of tilings of our region here as the product of these \mathcal{Q} -type regions as follows:

$$M(H_{x,0,z}^{(1)}(a_1, \dots, a_{2k})) = M(\mathcal{Q}(0, a_1, a_2, \dots, a_{2k-1})) M(\mathcal{Q}(a_1, a_2, \dots, a_{2k}, x, z)). \quad (58)$$

Then (14) follows again from (13) and Lemma 3.

The last base case is the case when $z = 0$. Similar to the cases of $x = 0$ and $y = 0$, we can break our region into two \mathcal{Q} -type regions as in Figure 10(b). Region-splitting Lemma gives us the following formula:

$$M(H_{x,0,z}^{(1)}(a_1, \dots, a_{2k})) = M(\mathcal{Q}(0, a_1, a_2, \dots, a_{2k-1}, a_{2k} + x, y)) M(\mathcal{Q}(a_1, a_2, \dots, a_{2k})). \quad (59)$$

Then (14) is implied by (13) and Lemma 3.

For the induction step, we assume that x, y, z are all positive and that (14) holds for any $H^{(1)}$ -type region with $2k$ holes ($k \geq 2$) whose sum of the x -, y - and z -parameters is strictly less than $x + y + z$. We need to show that (14) holds for the general region $H_{x,y,z}^{(1)}(\mathbf{a})$, where $\mathbf{a} = (a_1, a_2, \dots, a_{2k})$ is a sequence of positive integers with $k \geq 2$.

As proved by using Kuo condensation above, the tiling number of the region $H_{x,y,z}^{(1)}(\mathbf{a})$ satisfies the recurrence (43). Therefore, our work is now verifying that the formula in (14) satisfies the same recurrence.

Denote by $\Phi_{x,y,z}(\mathbf{a})$ the formula on the right-hand side of (14). We now need to verify that

$$\frac{\Phi_{x,y-1,z}(\mathbf{a}) \Phi_{x,y,z-1}(\mathbf{a})}{\Phi_{x,y-1,z-1}(\mathbf{a}) \Phi_{x,y,z}(\mathbf{a})} + \frac{\Phi_{x+1,y-1,z-1}(\mathbf{a}) \Phi_{x-1,y,z}(\mathbf{a})}{\Phi_{x,y-1,z-1}(\mathbf{a}) \Phi_{x,y,z}(\mathbf{a})} = 1. \quad (60)$$

We write $\Phi_{x,y,z}(\mathbf{a}) = \phi_{x,y,z}(O(\mathbf{a}), E(\mathbf{a})) \cdot f_{x,y,z}(\mathbf{a})$, where $\phi_{x,y,z}(a, b)$ was defined as in the verification of (13). We need to show that:

$$\begin{aligned} & \frac{f_{x,y-1,z}(\mathbf{a}) f_{x,y,z-1}(\mathbf{a}) \phi_{x,y-1,z}(O(\mathbf{a}), E(\mathbf{a})) \phi_{x,y,z-1}(O(\mathbf{a}), E(\mathbf{a}))}{f_{x,y-1,z-1}(\mathbf{a}) f_{x,y,z}(\mathbf{a}) \phi_{x,y-1,z-1}(O(\mathbf{a}), E(\mathbf{a})) \phi_{x,y,z}(O(\mathbf{a}), E(\mathbf{a}))} \\ & + \frac{f_{x+1,y-1,z-1}(\mathbf{a}) f_{x-1,y,z}(\mathbf{a}) \phi_{x+1,y-1,z-1}(O(\mathbf{a}), E(\mathbf{a})) \phi_{x-1,y,z}(O(\mathbf{a}), E(\mathbf{a}))}{f_{x,y-1,z-1}(\mathbf{a}) f_{x,y,z}(\mathbf{a}) \phi_{x,y-1,z-1}(O(\mathbf{a}), E(\mathbf{a})) \phi_{x,y,z}(O(\mathbf{a}), E(\mathbf{a}))} = 1. \end{aligned} \quad (61)$$

For given sequence $\mathbf{a} = (a_1, a_2, \dots, a_n)$, the function $f_{x,y,z}(\mathbf{a})$ only depends on the parameters y and z and the sums $x + y$ and $x + z$. The values of these quadruples for the four occurrences of f in the first fraction in the first terms are:

$$\begin{aligned} & (y - 1, z, x + y - 1, x + z), & (y, z - 1, z + y, x + z - 1) \\ & (y - 1, z - 1, x + y - 1, x + z - 1), & (y, z, x + y, x + z), \end{aligned} \quad (62)$$

where the two quadruples on the top row correspond to the two f -functions on the numerator, and the two quadruples on the bottom row correspond to the two f -functions

on the denominator of our fraction. In particular, the values of these parameters in each row form the same set. This means that the product of two f -functions on the numerator is equal to the product of the ones on the denominator. It means that

$$\frac{f_{x,y-1,z}(\mathbf{a})f_{x,y,z-1}(\mathbf{a})}{f_{x,y-1,z-1}(\mathbf{a})f_{x,y,z}(\mathbf{a})} = 1. \quad (63)$$

Similarly, we have

$$\frac{f_{x+1,y-1,z-1}(\mathbf{a})f_{x-1,y,z}(\mathbf{a})}{f_{x,y-1,z-1}(\mathbf{a})f_{x,y,z}(\mathbf{a})} = 1, \quad (64)$$

and (61) becomes

$$\begin{aligned} & \frac{\phi_{x,y-1,z}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))}{\phi_{x,y-1,z-1}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))} \frac{\phi_{x,y,z-1}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))}{\phi_{x,y,z}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))} \\ & + \frac{\phi_{x+1,y-1,z-1}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))}{\phi_{x,y-1,z-1}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))} \frac{\phi_{x-1,y,z}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))}{\phi_{x,y,z}(\mathbf{O}(\mathbf{a}), \mathbf{E}(\mathbf{a}))} = 1. \end{aligned} \quad (65)$$

However, this is exactly a consequence of (13), then (14) follows.

Finally, as mentioned before, any $H^{(1)}$ -type region with an odd number of holes can be viewed as a region with an even number of holes, when the rightmost hole has size 0. This means (15) follows. \square

The proofs of Theorems 5–7 are essentially the same as that of Theorem 4, and will be omitted.

We devote the next part of this section to the proof of Theorem 8.

Proof of Theorem 8. This proof follows the lines in the proof of Theorem 4.

We first prove (22) by induction on $x + y + z$. The base cases are still: $x = 0$, $y = 0$, and $z = 0$.

Similar to the case of $H^{(1)}$ -type regions, if $x = 0$, we can split our regions into two disjoint subregions (and several forced lozenges as in Figure 13). By Region-splitting Lemma 13, we get

$$M(H_{0,y,z}^{(5)}(a_1, \dots, a_{2k})) = 2^{a_1} M(\mathcal{K}'(0, a_1 + 1, a_2, \dots, a_{2k}, y)) M(\mathcal{Q}(a_1, a_2, \dots, a_{2k} + z)). \quad (66)$$

Then (22) follows from the above identity (when specializing $k = 1$) and Lemma 3.

If $y = 0$, Figure 14 and Region-splitting Lemma 13 tell us

$$M(H_{x,0,z}^{(5)}(a_1, \dots, a_{2k})) = 2^{a_1} M(\mathcal{K}'(0, a_1 + 1, a_2, \dots, a_{2k-1})) M(\mathcal{Q}(a_1, a_2, \dots, a_{2k}, x, z)). \quad (67)$$

Finally, when $z = 0$, we have as in Figure 15

$$M(H_{x,y,0}^{(5)}(a_1, \dots, a_{2k})) = 2^{a_1} M(\mathcal{K}'(0, a_1 + 1, a_2, \dots, a_{2k-1}, a_{2k} + x, y)) M(\mathcal{Q}(a_1, a_2, \dots, a_{2k})). \quad (68)$$

Then (22) follows again from Lemma 3 in the cases $y = 0$ and $z = 0$.




Figure 13: Spitting up a $H^{(5)}$ -type region in the case of $x = 0$.




Figure 14: The base case $y = 0$ for the $H^{(5)}$ -type regions.




Figure 15: Applying the Region-splitting Lemma 13 to a $H^{(5)}$ -type region in the case of $z = 0$.

For the induction step, we assume that $x, y, z \geq 1$ and that (22) holds for any $H^{(5)}$ -type region with two holes whose sum of the x -, y -, and z -parameters is strictly less than $x + y + z$. We need to show (22) for any region $H_{x,y,z}^{(5)}(a, b)$.

We apply Kuo condensation to the dual graph G of the region $H_{x,y,z}^{(5)}(\mathbf{a})$ for a general sequence of nonnegative integers $\mathbf{a} = (a_1, a_2, \dots, a_n)$, based on Figure 16. The figure tells us that the product of the tiling numbers of the two regions in the top row equals the product of the tiling numbers of the two regions in the middle row plus the product of the tiling numbers of the two regions in the bottom row. In particular, we have

$$\begin{aligned} M(H_{x,y,z}^{(5)}(\mathbf{a})) M(H_{x,y-1,z-1}^{(5)}(\mathbf{a})) &= M(H_{x,y-1,z}^{(5)}(\mathbf{a})) M(H_{x,y,z-1}^{(5)}(\mathbf{a})) \\ &\quad + M(H_{x+1,y-1,z-1}^{(5)}(\mathbf{a})) M(H_{x-1,y,z}^{(5)}(\mathbf{a})), \end{aligned} \quad (69)$$

for any sequence $\mathbf{a} = (a_1, a_2, \dots, a_k)$. To verify (22) we only need to show that the expression on the right-hand side of its satisfies the same recurrence.

We denote by $\psi_{x,y,z}(a, b)$ the product on the right-hand side of (22), we would like to show that

$$\psi_{x,y,z}(a, b) \psi_{x,y-1,z-1}(a, b) = \psi_{x,y-1,z}(a, b) \psi_{x,y,z-1}(a, b) + \psi_{x+1,y-1,z-1}(a, b) \psi_{x-1,y,z}(a, b), \quad (70)$$

or

$$\frac{\psi_{x,y-1,z}(a, b)}{\psi_{x,y-1,z-1}(a, b)} \frac{\psi_{x,y,z-1}(a, b)}{\psi_{x,y,z}(a, b)} + \frac{\psi_{x+1,y-1,z-1}(a, b)}{\psi_{x,y-1,z-1}(a, b)} \frac{\psi_{x-1,y,z}(a, b)}{\psi_{x,y,z}(a, b)} = 1. \quad (71)$$

We have several additional claims (that follow directly from the definition of the products T and V) as follows:

Claim 20.

$$\frac{V(x, n, m)}{V(x, n - 1, m)} = [x + 2n - 2m]_m. \quad (72)$$

Claim 21.

$$\frac{V(x, n, m)}{V(x - 2, n, m)} = \frac{[x + 2n - 2m]_m}{[x - 2]_m}. \quad (73)$$

Claim 22.

$$\frac{T(x, n, m)}{T(x, n - 1, m)} = (x + n - m)_m. \quad (74)$$

Using the above claims, we now have the first fraction in the first term on the left-hand side of (71) simplified as

$$\begin{aligned} \frac{\psi_{x,y-1,z}(a,b)}{\psi_{x,y-1,z-1}(a,b)} &= \frac{P_{z+b,z+b,a} P_{y+z+b-2,y+z+b-2,a}}{P_{z+b-1,z+b-1,a} P_{y+z+b-1,y+z+b-1,a}} \\ &\times \frac{V(2a+2b+3, y+z-2, y-1) V(2x+2a+2b+3, y+z-3, y-1)}{V(2a+2b+3, y+z-3, y-1) V(2x+2a+2b+3, y+z-2, y-1)} \\ &\times \frac{T(x+b+1, y+z+2a-1, y-1) T(b+1, y+z+2a-2, y-1)}{T(x+b+1, y+z+2a-2, y-1) T(b+1, y+z+2a-1, y-1)} \\ &\times \frac{T(z+1, y+b-2, b) T(x+z, y+b-2, b) Q(a, b, x, y+z-1)}{T(z, y+b-2, b) T(x+z+1, y+b-2, b) Q(a, b, x, y+z-2)} \quad (75) \\ &= \frac{P_{z+b,z+b,a}}{P_{z+b-1,z+b-1,a}} \\ &\times \frac{y+z+b-1}{y+z+a+b-1} \frac{(y+z+2a+b-1)!}{(2y+2z+2a+2b-3)!} \frac{(2y+2z+2b-3)!}{(y+z+b-1)!} \\ &\times \frac{[2z+2a+2b+1]_{y-1}}{[2x+2z+2a+2b+1]_{y-1}} \frac{(x+z+2a+b+1)_{y-1}}{(z+2a+b+1)_{y-1}} \frac{(y+z-1)_b}{(z)_b} \frac{(x+z)_b}{(x+y+z-1)_b} \\ &\times \frac{(x+y+z+a+b-1)(2x+2y+2z+2a+2b-3)!}{(2y+2z+2b-3)!} \\ &\times \frac{(y+z-2)!(x+y+z+b-2)!(x+y+z+2a+2b-1)!}{(x+y+z-2)!(x+y+z+2a+b-1)!(2x+y+z+2a+2b-1)!} \quad (76) \end{aligned}$$

Working the same for the fraction $\frac{\psi_{x,y,z-1}(a,b)}{\psi_{x,y,z}(a,b)}$ and multiplying up the two simplifications, we get

$$\frac{\psi_{x,y-1,z}(a,b)}{\psi_{x,y-1,z-1}(a,b)} \frac{\psi_{x,y,z-1}(a,b)}{\psi_{x,y,z}(a,b)} = \frac{(y+z+a+b)(2x+y+z+2a+2b)}{(x+y+z+a+b)(x+y+z+2a+2b)}. \quad (77)$$

Next, we work on the second term in (71). We get the following simplifications:

$$\begin{aligned}
\frac{\psi_{x-1,y,z}(a,b)}{\psi_{x,y,z}(a,b)} &= \frac{V(2x+2a+2b+3, y+z-1, y)}{V(2x+2a+2b+1, y+z-1, y)} \\
&\times \frac{T(x+b, y+z+2a, y)}{T(x+b+1, y+z+2a, y)} \frac{T(x+2a+b+2, y+b-1, b)}{T(x+2a+b+1, y+b-1, b)} \frac{T(x+z+1, y+b-1, b)}{T(x+z, y+b-1, b)} \\
&\times \frac{Q(a,b, x-1, y+z)}{Q(a,b, x, y+z)} \tag{78} \\
&= \frac{[2x+2z+2a+2b+1]_y}{[2x+2a+2b+1]_y} \frac{(x+b)_y}{(x+z+2a+b+1)_y} \frac{(x+y+2a+b+1)_b}{(x+2a+b+1)_b} \frac{(x+y+z)_b}{(x+z)_b} \\
&\times \frac{(x+a+b)}{(x+y+z+a+b)} \frac{1}{(2x+2y+2z+2a+2b-1)!(2x+2a+2b)!} \\
&\times \frac{(x+y+z-1)!(x+b-1)!}{(x+y+z+b-1)!(x-1)!} \\
&\times \frac{(x+y+z+2a+b)!(x+2a+2b)!(2x+y+z+2a+2b-1)!(2x+y+z+2a+2b)!}{(x+y+z+2a+2b)!(x+2a+b)!}, \tag{79}
\end{aligned}$$

and then

$$\frac{\psi_{x+1,y-1,z-1}(a,b)}{\psi_{x,y-1,z-1}(a,b)} \frac{\psi_{x-1,y,z}(a,b)}{\psi_{x,y,z}(a,b)} \frac{x(x+a+b)}{(x+y+z+a+b)(x+y+z+2a+2b)}. \tag{80}$$

It is easy to see that

$$\begin{aligned}
&\frac{\psi_{x,y-1,z}(a,b)}{\psi_{x,y-1,z-1}(a,b)} \frac{\psi_{x,y,z-1}(a,b)}{\psi_{x,y,z}(a,b)} + \frac{\psi_{x+1,y-1,z-1}(a,b)}{\psi_{x,y-1,z-1}(a,b)} \frac{\psi_{x-1,y,z}(a,b)}{\psi_{x,y,z}(a,b)} = \\
&\frac{(y+z+a+b)(2x+y+z+2a+2b)}{(x+y+z+a+b)(x+y+z+2a+2b)} + \frac{x(x+a+b)}{(x+y+z+a+b)(x+y+z+2a+2b)} \\
&= 1, \tag{81}
\end{aligned}$$

and (71) follows.

We also prove (23) by induction on $x+y+z$. The base cases are also the situations when at least one of the three parameters x, y, z is equal to 0. This cases follows from the equations when proving the base cases for (22).

The induction step here is exactly the same as that in the proof of (22). We only need to show that the formula on the right-hand side of (23) also satisfies the recurrence (69). Similar to the proof of Theorem 4, this verification follows from (71). \square

Theorems 9–11 can be treated similarly to Theorem 8 in a completely analogous manner, and we also omit the proofs of these theorems here.

We finally prove Theorem 12.




Figure 16: Applying Kuo condensation to a $H^{(5)}$ -type region.




Figure 17: Apply Ciucu's factorization theorem to a symmetric hexagon with an array of holes on the symmetric axis.

Proof of Theorem 12. Apply Ciucu's Factorization Theorem 15 to the dual graph G of $\mathcal{S} = \mathcal{S}_{x,y,z}(\mathbf{a})$, we obtain

$$M(\mathcal{S}) = M(G) = 2^{y+a_2+\dots+a_n} M(G^+) M(G^-). \quad (82)$$

We consider first the case when x and a_1 are both even. Note that we also have z even in this case (z and x always have the same parity). It turns out that the component graph G^- (defined by the cutting procedure in Theorem 15) is the dual graph of the region

$$H_{\frac{x}{2}+E(\mathbf{a}), y-\frac{z}{2}+E(\mathbf{a}), \frac{z}{2}-E(\mathbf{a})}^{(3)} \left(\frac{a_1}{2}, a_2, \dots, a_n \right)$$

(see Figure 17(a)). Thus,

$$M(G^-) = H_{\frac{x}{2}+E(\mathbf{a}), y-\frac{z}{2}+E(\mathbf{a}), \frac{z}{2}-E(\mathbf{a})}^{(3)} \left(\frac{a_1}{2}, a_2, \dots, a_n \right). \quad (83)$$

The graph G^+ after removed several forced edges (the edges corresponding to the forced lozenges in Figure 17(a)) is the dual graph of the region

$$H_{\frac{x}{2}+E(\mathbf{a}), y-\frac{z}{2}+E(\mathbf{a}), \frac{z}{2}-E(\mathbf{a})}^{(2)} \left(\frac{a_1}{2}, a_2, \dots, a_n \right).$$

Since the removal of these forced edges does not affect to the number of matchings, we have

$$M(G^+) = H_{\frac{x}{2}+E(\mathbf{a}), y-\frac{z}{2}+E(\mathbf{a}), \frac{z}{2}-E(\mathbf{a})}^{(2)} \left(\frac{a_1}{2}, a_2, \dots, a_n \right). \quad (84)$$

Then part (1) of the theorem follows from (82)–(84).

Parts (2), (3), and (4) can be similarly treated by using Ciucu’s Factorization Theorem, based on Figures 17(b), (c), and (d), respectively. \square

Remark 23. If the parameter z in the region $\mathcal{S} = \mathcal{S}_{x,y,z}(\mathbf{a})$ is too small or too large, then the region does not have a tiling. We can still apply Ciucu’s Factorization Theorem to the dual graph G of the region \mathcal{S} for any z , and still get the equality (82). However, if $z < 2E(\mathbf{a}) - 1$ or $z > 2y + 2E(\mathbf{a}) + 1$, then at least one of the component graphs G^+ and G^- has no matching. To see this, we consider the regions corresponding to them. Let us work in detail here for the case of even x and a_1 (the other cases can be treated in the same manner). If $z < 2E(\mathbf{a}) - 1$, then the region corresponding to G^- is still an $H^{(3)}$ -type region, denoted by R , however, it has a *negative* y -parameter. In this case, divide the region into two subregions, denoted R_1 and R_2 for the upper and lower ones, along the line containing the bases of the triangular holes. By the same arguments in the proof of the Region-Splitting Lemma 13, we can show that a tiling of R (if exists) can be written as union of two independent tilings of R_1 and R_2 . However, it is easy to see that R_2 has no tiling (since it is not balanced). Similar in the case of $z > 2y + 2E(\mathbf{a}) + 1$, the upper subregion R_1 has no tiling. In summary, R has no tiling, so does \mathcal{S} .

References

- [1] M. Ciucu. Enumeration of perfect matchings in graphs with reflective symmetry. *J. Combin. Theory Ser. A*, 77: 67–97, 1997.
- [2] M. Ciucu. Plane partition I: A generalization of MacMahon’s formula. *Memoirs of Amer. Math. Soc.*, 178(839):107–144, 2005.
- [3] M. Ciucu. Another dual of MacMahon’s theorem on plane partitions. *Adv. Math.*, 306:427–450, 2017.
- [4] M. Ciucu and C. Krattenthaler. Enumeration of lozenge tilings of hexagons with cut off corners. *J. Combin. Theory Ser. A*, 100:201–231, 2002.
- [5] M. Ciucu and T. Lai. Lozenge Tilings of Doubly-intruded Hexagons. [arXiv:1712.08024](https://arxiv.org/abs/1712.08024), 2017.
- [6] E. H. Kuo. Applications of Graphical Condensation for Enumerating Matchings and Tilings. *Theor. Comput. Sci.*, 319: 29–57, 2004.

- [7] T. Lai. Enumeration of tilings of quartered Aztec rectangles. *Electron. J. Combin.*, 21(4): #P4.46, 2014.
- [8] T. Lai. A new proof for the number of lozenge tilings of quartered hexagons. *Discrete Math.*, 338:1866–1872, 2015.
- [9] T. Lai. Lozenge tilings of a halved hexagon with an array of triangles removed from the boundary. *SIAM J. Discrete Math.*, 32(1):783–814, 2018.
- [10] T. Lai. A q -enumeration of a hexagon with four adjacent triangles removed from the boundary. *European J. Combin.*, 64:66–87, 2017.
- [11] P. A. MacMahon. *Combinatory Analysis*, vol. 2. Cambridge Univ. Press, 1916, reprinted by Chelsea, New York, 1960.
- [12] R. Proctor. Odd symplectic groups. *Inven. Math.*, 92(2):307–332, 1988.
- [13] R. Rohatgi. Enumeration of lozenge tilings of halved hexagons with a boundary defect. *Electron. J. Combin.*, 22(4):#P4.22, 2015.
- [14] R. Stanley. Symmetries of plane partitions. *J. Combin. Theory Ser. A*, 43: 103–113, 1986.