

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications, Department of
Mathematics

Mathematics, Department of

1-1992

A NOTE ON THE CONNECTEDNESS PROBLEM FOR NEST ALGEBRAS

David R. Pitts

Follow this and additional works at: <https://digitalcommons.unl.edu/mathfacpub>



Part of the [Applied Mathematics Commons](#), and the [Mathematics Commons](#)

This Article is brought to you for free and open access by the Mathematics, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Mathematics by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

A NOTE ON THE CONNECTEDNESS PROBLEM FOR NEST ALGEBRAS

DAVID R. PITTS

(Communicated by Paul S. Muhly)

ABSTRACT. It has been conjectured that a certain operator T belonging to the group \mathcal{G} of invertible elements of the algebra $\text{Alg } \mathbf{Z}$ of doubly infinite upper-triangular bounded matrices lies outside the connected component of the identity in \mathcal{G} . In this note we show that T actually lies inside the connected component of the identity of \mathcal{G} .

Let \mathbf{T} be the unit circle in the complex plane with normalized Lebesgue measure. For $1 \leq p \leq \infty$, let H^p be the usual Hardy space of all functions in $L^p(\mathbf{T})$ that have analytic extensions to the open unit disk \mathbf{D} . Let $\mathcal{H} = L^2(\mathbf{T})$ and let $\mathcal{B}(\mathcal{H})$ be set of all bounded linear operators on \mathcal{H} . Let $W \in \mathcal{B}(\mathcal{H})$ be the shift operator: $(Wf)(e^{i\theta}) = e^{i\theta} f(e^{i\theta})$. In this paper, we consider the nest $\{W^n H^2 : n \in \mathbf{Z}\}$ of subspaces of $L^2(\mathbf{T})$, and its associated nest algebra,

$$\text{Alg } \mathbf{Z} = \{T \in \mathcal{B}(\mathcal{H}) : TW^n H^2 \subseteq W^n H^2 \text{ for all } n \in \mathbf{Z}\}.$$

A question which has been unanswered for several years is the following:

Question. Is the group of invertible elements of the Banach algebra $\text{Alg } \mathbf{Z}$ connected in the norm topology?

It is frequently conjectured that the answer to this question is no. The reason for conjecturing a negative answer is because of a strong analogy between nest algebras and analytic function theory. We refer the reader to the book by Davidson [1] for details and more background on this question.

For each $f \in L^\infty(\mathbf{T})$, let $M_f \in \mathcal{B}(\mathcal{H})$ be the multiplication operator,

$$M_f \phi = f \phi, \quad \phi \in L^2(\mathbf{T}).$$

Note that for $f \in H^\infty$, we have $M_f \in \text{Alg } \mathbf{Z}$. Let a be a positive real number and set

$$h(z) = \frac{ai}{\pi} \log \left(\frac{1+z}{1-z} \right).$$

Received by the editors July 19, 1990.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 47D25; Secondary 46B35.

Key words and phrases. Nest, nest algebra.

Research partially supported by NSF grant DMS-8702982 and by an NSF Mathematical Sciences Postdoctoral Fellowship.

Then h is a conformal map of the open unit disk onto the unbounded vertical strip $\{z \in \mathbf{C} : -a < \operatorname{Re}(z) < a\}$.

If $f = \exp(h)$ then it is easy to see that both f and $1/f$ are H^∞ functions and moreover, that f is not the exponential of any H^∞ function. Therefore f cannot be connected to the constant function 1 via a norm continuous path within the group of invertible elements of the Banach algebra H^∞ . For this reason, the operator M_f has been suggested as a possible example of an operator which cannot be connected to the identity via a norm continuous path inside the group of invertibles in $\operatorname{Alg} \mathbf{Z}$.

The purpose of this note is to show that in fact, M_f may be connected to the identity via a norm continuous path of invertible elements in $\operatorname{Alg} \mathbf{Z}$.

Before giving the proof we pause for some terminology and to make a few simple remarks.

Let \mathcal{A} be a unital Banach algebra with unit I . Say that an invertible element a of \mathcal{A} may be *connected to the identity* if there exists a norm continuous function $f: [0, 1] \rightarrow \mathcal{A}$ such that $f(0) = a$, $f(1) = I$, and $f(t)$ is an invertible element of \mathcal{A} for each t . The algebra \mathcal{A} has the *connectedness property* if every invertible element of \mathcal{A} may be connected to the identity. We use the term *symmetry* to describe a square root of the identity in a unital Banach algebra \mathcal{A} . Such elements have spectrum contained in the set $\{-1, 1\}$ and hence are connected to the identity. In fact, if $\gamma(t)$ is an arc in the complex plane connecting -1 to 1 which does not pass through the origin, then

$$(1) \quad \sigma(t) = \frac{I + S}{2} + \gamma(t) \frac{I - S}{2}$$

is a norm continuous path of invertible elements of \mathcal{A} which connects the symmetry S to the identity I .

The algebra

$$\mathcal{D} = \operatorname{Alg} \mathbf{Z} \cap (\operatorname{Alg} \mathbf{Z})^*$$

is a von Neumann subalgebra of $\operatorname{Alg} \mathbf{Z}$ and since any von Neumann algebra has the connectedness property, we see that any invertible operator in \mathcal{D} can be connected to the identity in $\operatorname{Alg} \mathbf{Z}$.

Remark. Let α be a complex number of unit modulus and let $g \in L^\infty(\mathbf{T})$. Let

$$g_\alpha(z) = g(\alpha z), \quad z \in \mathbf{T},$$

and define a unitary operator $S_\alpha \in \mathcal{D}$ by

$$S_\alpha e_n = \alpha^n e_n,$$

where $e_n(e^{i\theta}) = e^{in\theta}$ is the usual orthonormal basis for $L^2(\mathbf{T})$.

We then have

$$(2) \quad S_\alpha M_g S_\alpha^* = M_{g_\alpha}.$$

Note that by the above remarks, M_g and M_{g_α} belong to the same connectedness class of invertibles in $\operatorname{Alg} \mathbf{Z}$.

We now show that M_f can be connected to the identity. Note that $h(z) = -h(-z)$. It follows that we have

$$f(z)f(-z) = 1 \quad \text{for all } z \in \overline{\mathbf{D}}.$$

If $S = S_{-1}$, equation (2) yields,

$$SM_f SM_f = I.$$

Hence both S and SM_f are symmetries in $\text{Alg } \mathbf{Z}$ and

$$M_f = S(SM_f).$$

Therefore M_f can be connected to the identity in $\text{Alg } \mathbf{Z}$. Moreover, equation (1) enables one to obtain an explicit path connecting M_f to the identity.

Question. Let m be a conformal mapping of the disk onto itself and set $g = f \circ m$. Is M_g connected to the identity in $\text{Alg } \mathbf{Z}$? Note that the remark above shows that if m is a rotation, then this is the case.

Remark. Let R be any proper open subset of the complex plane that is simply connected and satisfies $R = -R$. Then $0 \in R$ and if h is any conformal map from the disk onto R with $h(0) = 0$, we have $h(z) = -h(-z)$. (Indeed, the function $g(z) = -h(-z)$ is also a conformal map of the disk onto R . Since $h(0) = g(0)$ and $h'(0) = g'(0)$, the Riemann mapping theorem implies $g = h$.) The argument given above now shows that if we assume that $\{\text{Re}(z) : z \in R\}$ is bounded and set $f = \exp(h)$, then M_f is a product of two symmetries in $\text{Alg } \mathbf{Z}$ and hence is connected to the identity in $\text{Alg } \mathbf{Z}$.

REFERENCES

1. K. R. Davidson, *Nest Algebras*, Research Notes in Math., vol. 191, Pitman, Boston, London, and Melbourne, 1988.
2. Z. Nehari, *Conformal mapping*, 1st ed., Internat. Ser. Pure and Appl. Math., McGraw-Hill, New York, Toronto, and London, 1952.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEBRASKA, LINCOLN, NEBRASKA, 68588

Current address: Department of Mathematics, University of California, Los Angeles, California 90024