2016

Evidence for Pipecolate Oxidase in Mediating Protection Against Hydrogen Peroxide Stress

Sathish Kumar Natarajan

University of Nebraska - Lincoln, snatarajan2@unl.edu

Ezhumalai Muthukrishnan

University of Nebraska-Lincoln

Oleh Khalimonchuk

University of Nebraska-Lincoln, okhalimonchuk2@unl.edu

Justin L. Mott

University of Nebraska Medical Center

Donald F. Becker

University of Nebraska-Lincoln, dbecker3@unl.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/biochemfacpub>

Part of the [Biochemistry Commons](#), [Biotechnology Commons](#), and the [Other Biochemistry, Biophysics, and Structural Biology Commons](#)

Natarajan, Sathish Kumar; Muthukrishnan, Ezhumalai; Khalimonchuk, Oleh; Mott, Justin L.; and Becker, Donald F., "Evidence for Pipecolate Oxidase in Mediating Protection Against Hydrogen Peroxide Stress" (2016). *Biochemistry -- Faculty Publications*. 278.
<http://digitalcommons.unl.edu/biochemfacpub/278>

This Article is brought to you for free and open access by the Biochemistry, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Biochemistry -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Evidence for Pipecolate Oxidase in Mediating Protection Against Hydrogen Peroxide Stress

Sathish Kumar Natarajan,^{1,2} Ezhumalai Muthukrishnan,² Oleh Khalimonchuk,¹
Justin L. Mott,³ and Donald F. Becker¹

1 Department of Biochemistry and Redox Biology Center, University of Nebraska–Lincoln, Lincoln, Nebraska 68588

2 Department of Nutrition and Health Sciences, University of Nebraska–Lincoln, Lincoln, Nebraska 68583

3 Department of Biochemistry and Molecular Biology, University of Nebraska Medical Center, Omaha, Nebraska 68198

Corresponding author — Donald F. Becker, Department of Biochemistry and Redox Biology Center, University of Nebraska–Lincoln, N258 Beadle Center, Lincoln, NE 68588; email dbecker3@unl.edu

Abstract

Pipecolate, an intermediate of the lysine catabolic pathway, is oxidized to Δ^1 -piperideine-6-carboxylate (P6C) by the flavoenzyme L-pipecolate oxidase (PIPOX). P6C spontaneously hydrolyzes to generate α -amino adipate semialdehyde, which is then converted into α -amino adipate acid by α -amino adipate semialdehyde dehydrogenase. L-pipecolate was previously reported to protect mammalian cells against oxidative stress. Here, we examined whether PIPOX is involved in the mechanism of pipecolate stress protection. Knockdown of PIPOX by small interference RNA abolished pipecolate protection against hydrogen peroxide-induced cell death in HEK293 cells suggesting a critical role for PIPOX. Subcellular fractionation analysis showed that PIPOX is localized in the mitochondria of HEK293 cells consistent with its role in lysine catabolism. Signaling pathways potentially involved in pipecolate protection were explored by treating cells with small molecule inhibitors. Inhibition of both mTORC1 and mTORC2 kinase complexes or inhibition of Akt kinase alone blocked pipecolate protection suggesting the involvement of these signaling pathways. Phosphorylation of the Akt downstream target, forkhead transcription factor O3 (FoxO3), was also significantly increased in cells treated with pipecolate, further implicating Akt in the protective mechanism and revealing FoxO3 inhibition as a potentially key step. The results presented here demonstrate that pipecolate metabolism can influence cell signaling during oxidative stress to promote cell survival and suggest that the mechanism of pipecolate protection parallels that of proline, which is also metabolized in the mitochondria.

Keywords: pipecolate, pipecolate oxidase, amino acid metabolism, oxidative stress, mitochondria

Abbreviations: AAA, α -amino adipic acid; AAS, α -AAA semialdehyde; AASDH, α -AAS dehydrogenase; 6AN, 6-aminonicotinamide; DHEA, dehydroepiandrosterone; DMSO, dimethyl sulfoxide; FoxO, fork head family of transcription factor class O; G6PDH, glucose-6-phosphate dehydrogenase; GSA, glutamate γ -semialdehyde; HEK293, human embryonic kidney 293; mTORC, mammalian target of rapamycin complex; MTS, [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium]; PTS, peroxisomal targeting signal; PIPOX, pipecolate oxidase; P6C, Δ^1 -piperideine-6-carboxylate; PRODH, proline dehydrogenase; P5C, Δ^1 -pyrroline-5-carboxylate; P5CDH, P5C dehydrogenase; P5CR, P5C reductase; VDAC, voltage-dependent anion channel

L-pipecolic acid is a nonproteinogenic amino acid of lysine metabolism. In humans, pipecolate is derived from the saccharopine branch of lysine catabolism and by an alternative lysine degradation pathway initiated by α -deamination of lysine [Struys and Jakobs, 2010]. L-pipecolate is oxidized to Δ^1 -piperideine-6-carboxylate (P6C) by the flavoenzyme pipecolate oxidase (PIPOX) (Fig. 1A). Spontaneous hydrolysis of P6C then generates α -amino adipate semialdehyde (AAS), an intermediate that is also generated from saccharopine by AAS synthase of the main lysine degradation pathway [Struys and Jakobs, 2010]. AAS is subsequently oxidized by NAD⁺-dependent AAS dehydrogenase

(AASDH1, also known as aldehyde dehydrogenase 7A1) [Luo and Tanner, 2015] to form α -amino adipic acid (AAA), which is eventually converted to glutaryl-CoA via α -ketoadipic acid [Struys and Jakobs, 2010].

Human PIPOX is a polypeptide of 390 residues with a molecular weight of 44 kDa and contains a covalently bound flavin cofactor [Dodt et al., 2000]. Characterization of PIPOX from different mammalian sources shows that in addition to oxidizing pipecolate, PIPOX can utilize sarcosine and proline as alternative substrates, albeit with lower efficiency [Mihalik et al., 1991; Dodt et al., 2000]. Elevated pipecolic acid concentration in serum

Fig. 1. Reactions catalyzed by pipecolate oxidase and proline dehydrogenase. (A) Pipecolate oxidase (PIPOX) catalyzes the oxidation of L-pipecolate (Pipe) to Δ^1 -piperideine-6-carboxylate (P6C). P6C spontaneously hydrolyzes to form α -amino adipic semialdehyde (AAS). (B) In an analogous reaction, proline dehydrogenase (PRODH) catalyzes the oxidation of L-proline (Pro) to Δ^1 -pyrroline-5-carboxylate (P5C), which spontaneously hydrolyzes to glutamate γ -semialdehyde (GSA).

(hyperpipecolic acidemia) is used as a marker of pyridoxine dependent epilepsy and AASDH deficiency [Stockler et al., 2011], and is associated with peroxisome biogenesis disorders such as Zellweger syndrome [Wanders et al., 1988; Moser et al., 1989].

Our interest in L-pipecolic acid is from a previous study in which we tested the ability of proline and proline analogs to protect mammalian cells against oxidative stress. In addition to proline, we found that pipecolate, a six-member proline analog, significantly increased cell survival after exposure to H₂O₂ stress [Krishnan et al., 2008]. The mechanism by which proline improves cell survival was shown to be mediated by proline dehydrogenase (PRODH) [Natarajan et al., 2012]. PRODH is a mitochondrial enzyme that performs the first step of proline degradation by catalyzing the oxidation of proline to Δ^1 -pyrroline-5-carboxylate (P5C) (Fig. 1B), which is then converted into glutamate by P5C dehydrogenase. Knockdown of PRODH eliminated the protective effect of proline against H₂O₂ stress [Natarajan et al., 2012]. Additionally, it was shown that proline and PRODH increase activation of Akt signaling during stress suggesting a mechanism for how proline enhances cell survival [Natarajan et al., 2012].

Here we investigate the mechanism of pipecolate protection against H₂O₂ induced cell death. PIPOX catalyzes the oxidation of L-pipecolate in a reaction analogous to PRODH (Fig. 1A) suggesting that similar to proline, the protective effect of pipecolate may depend on this first oxidative step. The role of PIPOX in pipecolate protection was thus examined and the mechanistic target of rapamycin complex (mTOR) and Akt kinase were explored to test their potential involvement in pipecolate protection. Subcellular localization of PIPOX was also assessed to establish whether PIPOX is expressed in mitochondria. We found that in HEK293 cells a majority of PIPOX localizes to

mitochondria and our results suggest that the mechanism by which pipecolate provides cellular stress protection parallels that of proline.

Materials and Methods

Materials

Unless stated otherwise, all chemicals, enzymes, and buffers were purchased from Fisher Scientific and Sigma-Aldrich, Inc. For this study, we used human embryonic kidney 293 (HEK293) cells purchased from ATCC.

Antibodies

Antibodies recognizing VDAC/Porin (#V2139) and b-Actin (#A5441) were from Sigma. The monoclonal antibody against PIPOX (PIPOX F-9, #sc-166749) was from Santa-Cruz and anticalase antibody (#ab15834) was from Abcam. Antibodies against phospho-Akt T308 (#4056), phospho-Akt S473 (#4058), phospho-JNK (#4668), phospho-FoxO3 (Thr32) (#9464) and total forms of FoxO3 (#2497), Akt (#4685), and JNK (#9252) were from Cell Signaling Technology. Horseradish peroxidase-labeled anti-mouse (#NA931), and anti-rabbit (#RPN4301) secondary antibodies were from GE Healthcare.

Stress Tests and Cell Viability

HEK293 cells were grown in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% FBS and penicillin-streptomycin (100 U/ml). For stress treatments, HEK293 were grown to 80% confluence and then pretreated with and without L-pipecolate in serum containing medium for 12 h. After the pretreatment period, cells were exposed to H₂O₂ (50 and 500 μ M) for 3 h in serum free medium with and without L-pipecolate.

The protective effect of pipecolate was initially tested with different pipecolate concentrations (0.05–10 mM) and then 1, 5, and 10 mM L-pipecolate were used in subsequent experiments. After stress treatment, cell viability was quantified using the tetrazolium compound [3-(4, 5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium] (MTS) according to the recommendation of the manufacturers (Promega). The CellTiter-Glo Luminescent assay (Promega) was also used as an alternative method according to the recommendation of the manufacturers. Percent cell survival was determined relative to the appropriate controls. Stress tests with proline were performed similarly to pipecolate but using 5 mM L-proline. Stress tests were also performed in the presence of different inhibitor compounds. To inhibit the pentose phosphate pathway, HEK293 cells were incubated with 6-aminonicotinamide (6-AN) (50 μ M) or dehydroepiandrosterone (DHEA) (50 μ M) for 24 h before adding pipecolate (10 mM) to the cell culture. After 12 h incubation with pipecolate, cells were then treated with H₂O₂ as described above. Inhibition of the mechanistic target of rapamycin (mTOR) pathway was achieved by incubating HEK293 cells with rapamycin (Calbiochem) at 10 and 100 nM concentrations for 2 h prior to H₂O₂ stress. Inhibition of both mTORC1 and mTORC2 was achieved by incubating HEK293 cells with 10 nM of KU-0063794 for 2 h prior to H₂O₂ stress. To inhibit Akt activity, HEK293 were treated with 10 μ M Akt Inhibitor VIII (Akti-Calbiochem), which selectively inhibits Akt1/Akt2 activity, for 30 min prior to H₂O₂ stress. Stock solutions of the inhibitor compounds DHEA, 6-AN, rapamycin, and Akti were prepared in dimethyl sulfoxide (DMSO).

G6pdh Assay and NADPH Measurements

Glucose-6-phosphate dehydrogenase (G6PDH) activity was measured according to the recommendation of the manufacturer (BioVision). One unit of enzyme is defined as 1 μ mol of product formed per min at 37°C. Total protein concentration in the cell lysates was determined using the bicinchoninic acid (BCA) protein assay (Pierce) using bovine serum albumin as a standard. Cellular NADPH/NADP⁺ (MilliporeSigma) levels were estimated as previously described according to the manufacturer's recommendation [Natarajan et al., 2012].

Western Blot Analysis

Cells grown in 10-cm diameter dishes were rinsed once with ice cold PBS and cell lysates were prepared using M-PER according to the recommendation of the manufacturers along with the addition of mammalian protease inhibitor cocktail (Sigma) and HALT-phosphatase inhibitor (Pierce). Lysates containing 50–100 μ g of protein were resolved by SDS-PAGE. Proteins were transferred to a PVDF membrane and visualized by immunoblotting.

Mitochondrial Isolation

Cells were released with trypsin and washed twice via centrifugation with ice cold PBS. Cell pellets were then resuspended in Tris buffer (5 mM, pH 7.4) containing 0.25 M sucrose, 1 mM EDTA, and mammalian protease inhibitor cocktail (Sigma). The resuspended cells were next hand homogenized using a dounce homogenizer for 1 min on ice. Mitochondria and cytosolic fractions

were isolated by differential centrifugation as described [Natarajan et al., 2006a]. Briefly, cell homogenates were first centrifuged at 600g for 10 min to remove cell debris. Mitochondria were then pelleted at 17,000g for 15 min (4°C) and washed twice with mitochondrial suspension buffer (5 mM Tris, pH 7.4, 0.25 M sucrose). The post-mitochondrial supernatant was then centrifuged at 32,000g for 20 min to isolate the peroxisome fraction [Natarajan et al., 2006a]. The peroxisomal fraction was resuspended in the mitochondrial suspension buffer and then centrifuged at 17,000g for 15 min to remove mitochondria. The supernatant was then centrifuged again at 32,000g for 20 min to collect pure peroxisomes [Natarajan et al., 2006a,b]. The postperoxisomal supernatant was centrifuged at 105,000g for 1 h to isolate the cytosolic fraction [Natarajan et al., 2006a]. Density-purified mitochondria were isolated on a discontinuous Histodenz [5-(N-2,3-Dihydroxypropylacetamido)-2,4,6-triiodo-N,N'-bis(2,3-dihydroxypropyl) isophthalamide] gradient as described [Boldogh and Pon, 2007; Atkinson et al., 2011]. Briefly, the mitochondria-enriched fraction obtained from differential centrifugation was layered onto a discontinuous Histodenz gradient consisting of 14% and 22% layers dissolved in sorbitol buffer. Histodenz gradients were centrifuged at 303,000g for 1 h [Atkinson et al., 2011]. Mitochondrial fractions were then resuspended in mitochondrial suspension buffer and used for Western blot analysis. Total protein concentrations were quantified using the Bradford method (Bio-Rad).

siRNA Transfection

Cells were transfected with On-Target Plus Smart Pool siRNA of PIPOX and On-Target Plus Non-targeting Pool, referred to as control siRNA (Dharmacon RNA Technologies). siRNA transfection was performed for 48 h before doing the stress test or whole cell extract collection. Knockdown was confirmed by immunoblotting. Following transfection, stress tests were performed as described above.

Statistical Analysis

Data are expressed as mean $\pm$ standard deviation (SD). Statistical analysis was performed using student's *t*-test. A *P*-value of < 0.05 was considered to be statistically significant.

Results

Pipecolate Protects HEK293 Cells

Pipecolate was previously shown to protect HEK293 cells against H₂O₂-induced cell death [Krishnan et al., 2008]. Here we pre-treated HEK293 cells with different concentrations of pipecolate (0.05–10 mM) for 12 h prior to H₂O₂ stress treatment (0.5 mM, 3 h). Pipecolate protected HEK293 cells against oxidative stress with significant protection observed at ≥ 0.75 mM pipecolate relative to control cells without pipecolate supplementation (Fig. 2A and B) [Krishnan et al., 2008]. The protective effect of pipecolate was also investigated using luminescence quantification of ATP generation. In these assays, pipecolate increased cell survival in HEK293 cells (Fig. 2C) from 40% to 74%.

Pipecolate protection was also examined at a more pathophysiology-relevant level of oxidative stress by using a lower concentration of H₂O₂. At 50 μ M H₂O₂ (3 h), percent cell survival

Fig. 2. Oxidative stress protection of HEK293 cells by pipecolate. **(A)** HEK293 cells were treated without pipecolate (Con) and with different concentrations of pipecolate (Pipe) for 12 h (0.05–1 mM). Control and pipecolate treated cells were then incubated in serum free medium with and without 0.5 mM H₂O₂ for 3 h. Percent cell survival was estimated using the MTS cell viability assay. **(B)** HEK 293 cells were treated with or without pipecolate (0.75–10 mM) for 12 h. Control and pipecolate treated cells were then incubated in serum free medium with and without 500 μM H₂O₂ for 3 h. Percent cell survival was estimated using the MTS cell viability assay. **(C)** HEK293 cells treated with and without pipecolate (10 mM) for 12 h. Control and pipecolate treated cells were then incubated in serum free medium with and without 500 μM H₂O₂ for 3 h. Percent cell survival was determined using the Cell Titer-Glo Luminescent assay. **(D)** HEK293 cells were treated as above. Control and pipecolate (10 mM) treated cells were then incubated in serum free medium with and without 50 μM H₂O₂ for 3 h. Percent cell survival was determined using the Cell Titer-Glo Luminescent assay. Each value represents mean ± SD from five different experiments (n = 5) (* P < 0.05).

was significantly higher in HEK293 cells treated with pipecolate relative to control cells without pipecolate (Fig. 2D). Thus, pipecolate was able to protect cells against a H₂O₂ concentration that more closely mimics pathophysiological stress.

PIPOX Knockdown by siRNA

We next examined the effect of reducing endogenous PIPOX expression by transfecting HEK293 cells with PIPOX siRNA (siPIPOX). Knockdown of PIPOX expression was confirmed by Western blot analysis of whole cell lysates. PIPOX expression was observed to be significantly lower in HEK293 cells after 48 h of siPIPOX treatment relative to cells treated with control siRNA (Fig. 3A). In siPIPOX-transfected cells, pipecolate protection against H₂O₂-induced cell death was eliminated whereas in control siRNA treated cells pipecolate protection was still observed (Fig. 3B). Pipecolate (1–5 mM) was also unprotective in siPIPOX-transfected cells exposed to pathophysiological levels of H₂O₂ (50 μM) (Fig. 3C). Next, we tested whether proline supplementation could protect against H₂O₂ stress in cells treated with PIPOX siRNA. Proline significantly increased survival of HEK293 cells treated with PIPOX siRNA (Fig. 3D). These results show that although PIPOX can utilize proline as an alternative substrate, PIPOX is not essential for proline protection. Thus, it appears PIPOX is necessary for pipecolate protection.

PIPOX Subcellular Localization

Because PIPOX was observed to be essential for pipecolate protection, we sought to establish the subcellular localization of PIPOX in HEK293 cells. Human PIPOX terminates at the C-terminus with an Ala-His-Leu tripeptide peroxisome targeting signal [IJlst et al., 2000], however, PIPOX has been shown previously to localize to the peroxisomes and the mitochondria depending on the species [IJlst et al., 2000]. Western analysis of PIPOX expression in the different subcellular compartments was performed using highly purified mitochondria, peroxisomes, and cytosolic fractions from HEK293 cells. Figure 4 shows that PIPOX was detected predominantly in isolated mitochondria. To check whether the purified mitochondria were contaminated with peroxisomes, we tested for the presence of catalase. High expression levels of catalase were observed in the peroxisomal fraction whereas catalase was not detected in the mitochondria indicating that the purified mitochondria were not contaminated with peroxisomes (Fig. 4).

NADPH/NADP⁺ and Pentose Phosphate Pathway

The effect of pipecolate on the NADPH/NADP⁺ ratio was measured in cells before and after H₂O₂ stress. The cellular NADPH/NADP⁺ ratio in HEK293 cells decreased in control cells after 3 h of 500 μM H₂O₂ stress (Fig. 5A). In cells treated with pipecolate,

Fig. 3. Knockdown of PIPOX abolishes pipecolate protection. **(A)** Western blot analysis of PIPOX in HEK293 cells transfected with control siRNA (siCon, 100 nM) and PIPOX siRNA (siPIPOX, 100 nM) for 48 h. Actin was used as a control. **(B)** Percent cell survival of control siRNA (siCon) or PIPOX siRNA (siPIPOX) treated HEK293 cells with and without 0.5mM H₂O₂ (3 h) in the absence (Con) and presence of 10mM pipecolate (Pipe). **(C)** Percent cell survival of control siRNA or PIPOX siRNA treated HEK293 cells with and without 50 μM H₂O₂ (3 h) in the absence (Con) and presence of 1 or 5 mM pipecolate (Pipe). **(D)** Percent cell survival of control siRNA (siCon) or PIPOX siRNA (siPIPOX) treated HEK293 cells with and without 500μMH₂O₂ (3 h) in the absence (Con) and presence of 5mM proline. Percent cell survival in all experiments was estimated using the MTS cell viability assay. Each value represents mean ± SD of separate experiments (n = 5) (* P < 0.05).

however, the NADPH/NADP⁺ ratio increased during H₂O₂ stress resulting in a approximately fivefold higher ratio than in control cells (Fig. 5A). Next, we tested whether increased NADPH/NADP⁺ ratio with pipecolate treatment is via activation of the pentose phosphate pathway. Cells were treated with 6-aminonicotinamide (6-AN) and dehydroepiandrosterone (DHEA), which inhibit glucose 6-phosphate dehydrogenase (G6PDH) of the pentose phosphate pathway. Treatment of cells with 6AN and DHEA showed a significant decrease in G6PDH activity (Fig. 5B) while the pipecolate-induced increase in the NADPH/NADP⁺ ratio was abolished in cells treated with the inhibitors (Fig. 5A). Further, we also tested the effect of pipecolate protection during inhibition of pentose phosphate pathway. Interestingly, pipecolate protection of HEK293 cells was still observed in the presence of the

Fig. 4. Subcellular localization of PIPOX. Immunoblot analysis of PIPOX in mitochondria, peroxisomes, and cytosolic fractions isolated from HEK293 cells. Actin, VDAC and catalase visualized with respective antibodies were used as fraction-specific markers for cytosol (Cyto.), mitochondria (Mito.), and peroxisomes (Per.), respectively.

6-AN and DHEA indicating that the pentose phosphate pathway is not critical in these H₂O₂ stress experiments (Fig. 5C and D).

Inhibition of mTORC

The mechanistic target of rapamycin complex 1 (mTORC1) is known to be activated by amino acids [Lynch, 2001; Zoncu et al., 2011] and we recently discovered a role for mTORC in proline protection [Natarajan et al., 2012]. To test whether mTORC1 and mTORC2 are also involved in pipecolate protection, HEK293 cells were incubated with 10nM rapamycin, a concentration previously shown to inhibit mTORC1 in cells [Lasithiotakis et al., 2008]. Treatment of cells with rapamycin did not attenuate pipecolate protection against H₂O₂ stress in HEK293 cells (Fig. 6A) suggesting that mTORC1 is not involved. Treating cells with a higher concentration of rapamycin (100 nM), which inhibits mTORC1 and the Rictor-complex of mTOR (mTORC2) [Sarabassov et al., 2006], abrogated pipecolate protection (Fig. 6A) implying that mTORC2 may be involved. This was further tested by treating HEK293 cells with 10nM of KU-0063794, which also inhibits mTORC1 and mTORC2. Similar to using a high concentration of rapamycin, treatment of cells with KU-0063794 abolished pipecolate protection against H₂O₂ stress, suggesting a role for mTORC2 in pipecolate protection (Fig. 6B).

Role of Akt and FoxO3 during H₂O₂ Stress

Akt/PKB (Protein kinase B) can significantly impact cell survival and proliferation and, is a downstream target of mTORC2 [Berndt et al., 2010; Altomare and Khaled, 2012]. Cells were treated with the Akt inhibitor, Akti, to test the involvement of Akt in pipecolate protection [Calleja et al., 2009]. At different concentrations of pipecolate (1 and 10 mM), Akti was observed to block pipecolate protection of HEK293 cells against H₂O₂ stress (Fig. 7A). Akti inhibition of Akt was confirmed by loss of phosphorylation at Akt residues T308 and S473 as shown in Figure 7B. Exposure of control cells to H₂O₂ increases phospho-Akt slightly at T308 and more significantly at S473 (Fig. 7B, -Akti). In the absence of H₂O₂, pipecolate does not increase the levels of phospho-Akt at T308 and S473 relative to control cells (Fig. 7B, -Akti). However, in the presence of H₂O₂, pipecolate appears to significantly increase

Fig. 5. Effect of pipecolate on intracellular NADPH/NADP⁺ ratio. **(A)** The ratio of NADPH/NADP⁺ was determined for control (Con) and pipecolate-treated (Pipe, 10 mM) HEK293 cells with and without 500 μ M H₂O₂ (3 h). The NADPH/NADP⁺ ratio was also estimated in cells incubated in the presence of 6AN or DHEA. Each value represents mean $\pm$ SD of separate experiments ($n = 5$) (* $P < 0.05$). **(B)** Glucose 6-phosphate dehydrogenase activity in HEK293 cells under different treatments. Percent survival of HEK 293 cells first treated with 6-amino nicotinamide (6-AN) **(C)** and dehydroepiandrosterone (DHEA). **(D)** Percent cell survival was estimated using the MTS cell viability assay. Each value represents mean $\pm$ SD of separate experiments ($n = 5$) (* $P < 0.05$).

the levels of phospho-Akt at S473 (Fig. 7B–D, –Akti). Treatment of cells with Akti abolished phosphorylation of Akt under all conditions consistent with inhibition of the Akt signaling pathway (Fig. 7B–D, +Akti). In these experiments, no changes in the levels of total-Akt and the actin control were observed (Fig. 7B). The effects of Akti on other stress signaling pathways such as c-Jun N-terminal Kinase (JNK) were also examined. Akti did not perturb the phosphorylation status of JNK (Fig. 7E).

Activation of Akt can lead to phosphorylation of several transcription factors including the forkhead transcription factor O3 (FoxO3). To test the downstream effects of Akt in pipecolate protection, phosphorylation of FoxO3 (Thr32) was examined after 0.5 and 3 h of H₂O₂ stress in cells with or without pipecolate. In control cells, a slight increase in phospho-FoxO3 levels was observed with 0.5 h of H₂O₂ treatment (Fig. 8A and B). A much more dramatic increase in phospho-FoxO3, however,

Fig. 6. Inhibition of mTORC2 blocks pipecolate protection. **(A)** Percent cell survival of HEK293 cells incubated with rapamycin (10 and 100 nM) and treated with and without 0.5 mM H₂O₂ (3 h) in the absence (Con) and presence of 10 mM pipecolate (Pipe). **(B)** Percent cell survival of HEK293 cells incubated with Ku-0063794 (10 nM) and treated with and without 0.5 mM H₂O₂ (3 h) in the absence (Con) and presence of 10 mM pipecolate (Pipe). Cells treated with DMSO were used as controls. Percent cell survival was estimated using the MTS cell viability assay. Each value represents mean $\pm$ SD of separate experiments ($n = 5$) (* $P < 0.05$).

was observed with H₂O₂ (0.5 h) in the presence of pipecolate (Fig. 8A and B). Addition of Akti (+Akti) eliminated phosphorylation of FoxO3 indicating involvement of the Akt pathway (Fig. 8A and B). At 3 h of H₂O₂ treatment, phospho-FoxO3 levels remained slightly higher in pipecolate treated cells relative to control cells (Fig. 8C). These results suggest increased phosphorylation of FoxO3 is part of the mechanism by which pipecolate protects cells against oxidative stress.

Discussion

Here we show that PIPOX is required for the observed effects of pipecolate, a finding which is analogous to the role of PRODH in proline protection [Natarajan et al., 2012]. The specificity of proline and pipecolate protection is governed by PRODH and PIPOX, respectively. Although PIPOX can use proline as a substrate, knockdown of PIPOX does not affect proline protection consistent with PRODH being the key proline catabolic enzyme. Conversely, pipecolate is not expected to be a substrate for PRODH [Zhu et al., 2002]. The mutual protective effects of proline and pipecolate, both of which are transported into mammalian cells via the sodium dependent IMINO transporter and the system IMINO transporter 1 [Kowalczyk et al., 2005; Takanaga et al., 2005], are likely due to the analogous flavin-dependent catalyzed reactions of PRODH and PIPOX in mitochondria. The oxidation of these related amino acids involves reduction of a flavin cofactor and subsequent electron transfer from the reduced

Fig. 7. Role of Akt in pipecolate protection. **(A)** Percent cell survival of HEK293 cells incubated without Akti (-Akti) or with 10 μ M Akti (+Akti) and treated with and without 500 μ M H₂O₂ stress (3 h) in the absence (Con) and presence of 1 or 10 mM pipecolate (Pipe). Cells treated with DMSO were used as controls. Percent cell survival was estimated using the MTS cell viability assay. Each value represents mean $\pm$ SD of separate experiments ($n = 5$) (* $P < 0.05$). **(B)** Western analysis of P-Akt (T308 and S473), T-Akt and Actin in HEK293 cells treated with and without pipecolate (1 mM) for 12 h and then incubated with and without 0.5 mM H₂O₂ for 0.5 h in serum free medium. Western analysis of P-Akt and T-Akt were also determined in cells with similar treatment conditions but in the presence of 10 μ M Akt inhibitor (+Akti). Quantified levels of **(C)** P-Akt (T308) and **(D)** P-Akt (S473) relative to total Akt (P-Akt/Akt). **(E)** Western analysis of P-JNK, and T-JNK in HEK293 cells treated with and without pipecolate (1 mM) for 12 h and then incubated with and without 0.5 mM H₂O₂ for 0.5 h in serum free medium. Western analysis of P-JNK and T-JNK were also determined with similar treatment conditions but in the presence of 10 μ M Akt inhibitor (+Akti). Blots shown are representative results from three independent experiments. Each value represents the mean $\pm$ SD of separate experiments ($n = 3$) (* $P < 0.05$).

flavin to an electron acceptor. PRODH is a mitochondrial enzyme that couples the oxidation of proline to reduction of the ubiquinone pool in the mitochondrial membrane (Fig. 1) [Wanduragala et al., 2010; Liu et al., 2012]. PRODH activity has been reported to generate reactive oxygen species (ROS) leading to either cell survival or cell death [Liu et al., 2012; Natarajan and Becker, 2012; Zarse et al., 2012; Zhang et al., 2015]. Here, we

show that PIPOX is also predominantly localized in the mitochondria. Whether PIPOX transfers electrons directly to the mitochondrial electron transport chain is not clear but PIPOX has been shown to generate H₂O₂ during catalytic turnover [Wanders et al., 1989; Mihalik et al., 1991]. Energy metabolism in the mitochondria is a potent source of ROS, which has been shown to have a plethora of effects on cell proliferation, death, and

Fig. 8. Involvement of FoxO3 in pipecolate protection. **(A)** Western analysis of P-FoxO3 (P-Thr32) and T-FoxO3 (total FoxO3) in HEK293 cells treated with and without pipecolate (1 mM) for 12 h and then incubated with and without 0.5 mM H₂O₂ for 0.5 h in serum free medium. Blots are shown for experiments performed in the absence (-Akti) and presence (+Akti) of 10 μ M Akt inhibitor **(B)** Quantified levels of P-FoxO3 (T32) relative to T-FoxO3 from Western blots as described in panel A in the absence (-Akti) and presence (+Akti) of Akt inhibitor. **(C)** Same as in panel A but incubated with and without 0.5 mM H₂O₂ for 3 h in serum-free medium. Blots shown are representative results from three independent experiments. Each value represents the mean $\pm$ SD of separate experiments ($n = 3$) (* $P < 0.05$).

other growth processes [Natarajan and Becker, 2012; Figueira et al., 2013]. We found that pipecolate shifts the redox status of the NADPH/NADP⁺ and helps maintain cellular NADPH/NADP⁺ ratios during stress consistent with PIPOX having a role in redox homeostasis. However, the increased NADPH/NADP⁺ ratio and pentose phosphate pathway were found not be critical for pipecolate protection against H₂O₂ stress. Further studies are underway to elucidate the mechanism by which pipecolate influences the redox environment during oxidative stress.

Finding that PIPOX was mainly localized in mitochondria of HEK293 cells was unexpected. Previous studies of PIPOX in rat and human liver detected PIPOX activity in both mitochondria and peroxisomes [Wanders et al., 1989; Rao et al., 1993]. The subcellular localization of PIPOX has been shown to be species-dependent with localization in the peroxisomes suggested to predominate in humans [Mihalik and Rhead, 1989, 1991]. In human fibroblasts overexpression of PIPOX resulted in localization of PIPOX in the peroxisomes as detected by immunofluorescence staining [IJlst et al., 2000]. Currently known peroxisomal targeting signals (PTS) are the C-terminal PTS-1 and the N-terminal PTS-2 [Gould and Collins, 2002; Wolf et al., 2010]. PIPOX has a PTS-1 sequence of Ala-His-Leu and does not have a mitochondrial target sequence [IJlst et al., 2000]. Bioinformatics analysis

of PIPOX using MitoProt [Claros and Vincens, 1996], a program that does not rely on identifying a targeting peptide but instead evaluates up to 47 parameters, gives a low probability (MitoProt score of 15%) for PIPOX import into mitochondria despite PIPOX not having a mitochondrial targeting sequence. Another bioinformatics tool, WoLF PSORT [Horton et al., 2007], predicts that PIPOX is favored to be localized in the mitochondria rather than in the peroxisomes. It is established that peroxisomal targeting sequences do not necessarily limit protein localization to the peroxisomes [Wolf et al., 2010]. In addition, it is becoming more recognized that proteins can be located in more than one compartment [O'Byrne et al., 2003; Hogenboom et al., 2004; Taylor et al., 2004; Yogev and Pines, 2011; Islinger et al., 2012]. Dual localization of proteins can occur by various mechanisms involving two target signals in a single polypeptide and target signals that are inaccessible or ambiguous [O'Byrne et al., 2003; Hogenboom et al., 2004; Taylor et al., 2004; Yogev and Pines, 2011; Islinger et al., 2012].

Mitochondrial localization of PIPOX in HEK293 cells is consistent with its role in the lysine catabolic pathway, which involves mitochondrial enzymes such as AASDH (also known as ALDH7A1) [Brocker et al., 2010; Struys and Jakobs, 2010; Stockler et al., 2011; Luo and Tanner, 2015]. Recently, PIPOX was suggested

to function as a metabolic repair enzyme by minimizing pipecolate accumulation and increasing the overall conversion of lysine to glutaryl-CoA [Struys and Jakobs, 2010; Van Schaftingen et al., 2013]. Pyrroline-5-carboxylate reductase (P5CR) has been proposed to catalyze the NADPH-dependent reduction of P6C to pipecolate as a side reaction thereby diminishing lysine catabolic efficiency [Struys and Jakobs, 2010; Van Schaftingen et al., 2013]. The preferred substrate for P5CR is Δ^1 -pyrroline-5-carboxylate, but P5CR from *Escherichia coli* has been shown to convert P6C to pipecolate [Fujii et al., 2002]. Humans have three isoforms of P5CR known as PYCR1, PYCR2, and PYCRL [Reversade et al., 2009; De Ingeniis et al., 2012]. PYCR1 and PYCR2 were recently shown to localize in the mitochondria [Reversade et al., 2009; De Ingeniis et al., 2012]. Finding that PIPOX is also localized in the mitochondria further supports a repair role for PIPOX, which merits future investigation.

The results here with pipecolate and previously with proline [Natarajan et al., 2012], indicate that the protective effects of these amino acids involve the Akt and its downstream mediators. In the previous study with proline, PRODH was shown to increase Akt phosphorylation (Ser473) and phosphorylation of the Akt downstream target, FoxO3 [Natarajan et al., 2012]. Phosphorylation of FoxO3 by Akt prevents activation of pro-apoptotic genes thus promoting cell survival [Guo et al., 2011]. In this study, inhibition of Akt abolished phosphorylation of FoxO3 and the protective effect of pipecolate. Thus, the protective mechanism of pipecolate appears to be mediated by Akt and FoxO3. As previously observed with proline, dual inhibition of mTORC complexes, blocks pipecolate protection against oxidative stress. These observations indicate that mTORC2 may be involved in the mechanism of cell protection. mTORC2 has been shown to activate Akt via phosphorylation of Ser473 [Guo et al., 2011; Murata et al., 2011]. How pipecolate and proline influence mTORC2 and Akt signaling is not clear but amino acids have been shown to induce mTORC1 and mTORC2 in mammalian cells [Tato et al., 2011]. In addition, evidence has been reported for mTORC2 being activated by mitochondrial ROS suggesting that the intracellular redox environment modulates mTORC2 [Kumari Kanchan et al., 2012]. Whether ROS is a key mediator by which proline and pipecolate metabolism mediate mTORC2 and Akt pathway activity remains to be determined.

In summary, we show that pipecolate metabolism leads to protection against H_2O_2 stress in mammalian cells. Our findings expand the potential role of pipecolate metabolism in mediating cell signaling. Previous studies reporting that pipecolate impacts oxidative stress were focused on exploring the role of pipecolate in neurological function. Incubation of rat cerebral cortex supernatants with pipecolate for 1 h was shown to decrease the activity of antioxidant enzymes, an effect that could be blocked by lipoic acid [Dalazen et al., 2014]. Pipecolate was also shown to stimulate apoptotic cell death in neurons [Matsumoto et al., 2003]. Treatment of mouse neuroblastoma cells with D- or L-pipecolate over 72 h decreased cell viability with a racemic mixture of DL-pipecolate having the most toxic effect [Matsumoto et al., 2003]. Because D-pipecolate decreased cell viability, the mechanism appears to be independent of pipecolate oxidase activity, which is specific for L-pipecolate [Matsumoto et al., 2003].

Different roles for D- and L-pipecolate have also been found in other biological contexts such as L-pipecolate acting as an antihypertensive by competitively inhibiting the angiotensin I-converting enzyme, whereas, the D-enantiomer does not have an antihypertensive function [Vranova et al., 2013]. Future studies will investigate the protective role of pipecolate metabolism and downstream pathways involving mTORC2, Akt and FoxO3 using other cell types to broaden our understanding of how pipecolate impacts the cellular redox environment.

Acknowledgments — We thank Dr. Nishikant Wase for his help with the bioinformatics analysis of PIPOX. This research was supported in part by grants P20GM104320 to SKN and JLM, and GM079393 and P30GM103335 to DFB from the National Institutes of Health.

Authors' contributions — SKN designed and performed the experiments. EM performed the NADPH/NADP⁺ measurements. OK designed and guided the subcellular localization experiments. JLM helped with the signaling pathway analysis. DFB designed and coordinated the overall project and all authors contributed to the writing of the manuscript.

References

- Altomare DA, Khaled AR. 2012. Homeostasis and the importance for a balance between AKT/mTOR activity and intracellular signaling. *Curr Med Chem* 19:3748–3762.
- Atkinson A, Smith P, Fox JL, Cui TZ, Khalimonchuk O, Winge DR. 2011. The LYR protein Mzm1 functions in the insertion of the Rieske Fe/S protein in yeast mitochondria. *Mol Cell Biol* 31:3988–3996.
- Berndt N, Yang H, Trinczek B, Betzi S, Zhang Z, Wu B, Lawrence NJ, Pellecchia M, Schonbrunn E, Cheng JQ, Sebti SM. 2010. The Akt activation inhibitor TCN-P inhibits Akt phosphorylation by binding to the PH domain of Akt and blocking its recruitment to the plasma membrane. *Cell Death Differ* 17:1795–1804.
- Boldogh IR, Pon LA. 2007. Purification and subfractionation of mitochondria from the yeast *Saccharomyces cerevisiae*. *Methods Cell Biol* 80:45–64.
- Brocker C, Lassen N, Estey T, Pappa A, Cantore M, Orlova VV, Chavakis T, Kavanagh KL, Oppermann U, Vasiliou V. 2010. Aldehyde dehydrogenase 7A1 (ALDH7A1) is a novel enzyme involved in cellular defense against hyperosmotic stress. *J Biol Chem* 285:18452–18463.
- Calleja V, Laguerre M, Parker PJ, Larijani B. 2009. Role of a novel PH-kinase domain interface in PKB/Akt regulation: Structural mechanism for allosteric inhibition. *PLoS Biol* 7:e17.
- Claros MG, Vincens P. 1996. Computational method to predict mitochondrially imported proteins and their targeting sequences. *Eur J Biochem* 241:779–786.
- Dalazen GR, Terra M, Jacques CE, Coelho JG, Freitas R, Mazzola PN, Dutra-Filho CS. 2014. Pipecolic acid induces oxidative stress in vitro in cerebral cortex of young rats and the protective role of lipoic acid. *Metab Brain Dis* 29:175–183.
- De Ingeniis J, Ratnikov B, Richardson AD, Scott DA, Aza-Blanc P, De SK, Kazanov M, Pellecchia M, Ronai Z, Osterman AL, Smith JW. 2012. Functional specialization in proline biosynthesis of melanoma. *PLoS ONE* 7:e45190.

- Dodt G, Kim DG, Reimann SA, Reuber BE, McCabe K, Gould SJ, Mihalik SJ. 2000. L-Pipecolic acid oxidase, a human enzyme essential for the degradation of L-pipecolic acid, is most similar to the monomeric sarcosine oxidases. *Biochem J* 345(3):487–494.
- Figueira TR, Barros MH, Camargo AA, Castilho RF, Ferreira JC, Kowaltowski AJ, Sluse FE, Souza-Pinto NC, Vercesi AE. 2013. Mitochondria as a source of reactive oxygen and nitrogen species: From molecular mechanisms to human health. *Antioxid Redox Signal* 18:2029–2074.
- Fujii T, Mukaiharu M, Agematu H, Tsunekawa H. 2002. Biotransformation of L-lysine to L-pipecolic acid catalyzed by L-lysine 6-aminotransferase and pyrroline-5-carboxylate reductase. *Biosci Biotech Biochem* 66:622–627.
- Gould SJ, Collins CS. 2002. Opinion: Peroxisomal-protein import: Is it really that complex? *Nat Rev Mol Cell Biol* 3:382–389.
- Guo JP, Coppola D, Cheng JQ. 2011. IKBKE protein activates Akt independent of phosphatidylinositol 3-kinase/PDK1/mTORC2 and the pleckstrin homology domain to sustain malignant transformation. *J Biol Chem* 286:37389–37398.
- Hogenboom S, Tuyp JJM, Espeel M, Koster J, Wanders RJA, Waterham HR. 2004. Mevalonate kinase is a cytosolic enzyme in humans. *J Cell Sci* 117:631–639.
- Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai K. 2007. WoLF PSORT: Protein localization predictor. *Nucleic Acids Res* 35: W585–587.
- IJlst L, de Kromme I, Oostheim W, Wanders RJ. 2000. Molecular cloning and expression of human L-pipecolate oxidase. *Biochem Biophys Res Commun* 270:1101–1105.
- Islinger M, Grille S, Fahimi HD, Schrader M. 2012. The peroxisome: An update on mysteries. *Histochem Cell Biol* 137:547–544.
- Kowalczyk S, Broer A, Munzinger M, Tietze N, Klingel K, Broer S. 2005. Molecular cloning of the mouse IMINO system: An Na⁺- and Cl⁻-dependent proline transporter. *Biochem J* 386:417–422.
- Krishnan N, Dickman MB, Becker DF. 2008. Proline modulates the intracellular redox environment and protects mammalian cells against oxidative stress. *Free Radic Biol Med* 44:671–681.
- Kumari Kanchan R, Tripathi C, Singh Baghel K, Kumar Dwivedi S, Kumar B, Sanyal S, Sharma S, Mitra K, Garg V, Singh K, Sultana S, Kamal Tripathi R, Kumar Rath S, Bhadauria S. 2012. Estrogen receptor potentiates mTORC2 signaling in breast cancer cells by upregulating superoxide anions. *Free Radic Biol Med* 53:1929–1941.
- Lasithiotakis KG, Sinnberg TW, Schitteck B, Flaherty KT, Kulms D, Maczey E, Garbe C, Meier FE. 2008. Combined inhibition of MAPK and mTOR signaling inhibits growth, induces cell death, and abrogates invasive growth of melanoma cells. *J Invest Dermatol* 128:2013–2023.
- Liu W, Le A, Hancock C, Lane AN, Dang CV, Fan TWM, Phang JM. 2012. Reprogramming of proline and glutamine metabolism contributes to the proliferative and metabolic responses regulated by oncogenic transcription factor c-MYC. *Proc Natl Acad Sci USA* 109:8983–8988.
- Luo M, Tanner JJ. 2015. Structural basis of substrate recognition by aldehyde dehydrogenase 7A1. *Biochemistry* 54:5513–5522.
- Lynch CJ. 2001. Role of leucine in the regulation of mTOR by amino acids: Revelations from structure-activity studies. *J. Nutr* 131:861S–865S.
- Matsumoto S, Yamamoto S, Sai K, Maruo K, Adachi M, Saitoh M, Nishizaki T. 2003. Pipecolic acid induces apoptosis in neuronal cells. *Brain Res* 980:179–184.
- Mihalik SJ, McGuinness M, Watkins PA. 1991. Purification and characterization of peroxisomal L-pipecolic acid oxidase from monkey liver. *J Biol Chem* 266:4822–4830.
- Mihalik SJ, Rhead WJ. 1989. L-pipecolic acid oxidation in the rabbit and cynomolgus monkey. Evidence for differing organellar locations and cofactor requirements in each species. *J Biol Chem* 264:2509–2517.
- Mihalik SJ, Rhead WJ. 1991. Species variation in organellar location and activity of L-pipecolic acid oxidation in mammals. *J Comp Physiol B* 160:671–676.
- Moser HW, Mihalik SJ, Watkins PA. 1989. Adrenoleukodystrophy and other peroxisomal disorders that affect the nervous-system, including new observations on L-pipecolic acid oxidase in primates. *Brain Dev* 11:80–90.
- Murata H, Sakaguchi M, Jin Y, Sakaguchi Y, Futami J, Yamada H, Kataoka K, Huh NH. 2011. A new cytosolic pathway from a Parkinson disease-associated kinase, BRPK/PINK1: Activation of AKT via mTORC2. *J Biol Chem* 286:7182–7189.
- Natarajan SK, Becker DF. 2012. Role of apoptosis-inducing factor, proline dehydrogenase, and NADPH oxidase in apoptosis and oxidative stress. *Cell Health Cytoskeleton* 2012:11–27.
- Natarajan SK, Eapen CE, Pullimood AB, Balasubramanian KA. 2006a. Oxidative stress in experimental liver microvesicular steatosis: Role of mitochondria and peroxisomes. *J Gastroenterol Hepatol* 21:1240–1249.
- Natarajan SK, Eapen CE, Pullimood AB, Balasubramanian KA. 2006b. Oxidative stress in experimental liver microvesicular steatosis: Role of mitochondria and peroxisomes. *J Gastroenterol Hepatol* 21:1240–1249.
- Natarajan SK, Zhu W, Liang X, Zhang L, Demers AJ, Zimmerman MC, Simpson MA, Becker DF. 2012. Proline dehydrogenase is essential for proline protection against hydrogen peroxide-induced cell death. *Free Radic Biol Med* 53:1181–1191.
- O'Byrne J, Hunt MC, Rai DK, Saeki M, Alexson SE. 2003. The human bile acid-CoA: Amino acid N-acyltransferase functions in the conjugation of fatty acids to glycine. *J Biol Chem* 278:34237–34244.
- Rao VV, Tsai MJ, Pan XM, Chang YF. 1993. L-Pipecolic acid oxidation in rat—Subcellular-localization and developmental-study. *Biochim Biophys Acta* 1164:29–35.
- Reversade B, Escande-Beillard N, Dimopoulou A, Fischer B, Chng SC, Li Y, Shboul M, Tham PY, Kayserili H, Al-Gazali L, Shahwan M, Brancati F, Lee H, O'Connor BD, Schmidt-von Kegler M, Merriam B, Nelson SF, Masri A, Alkazaleh F, Guerra D, Ferrari P, Nanda A, Rajab A, Markie D, Gray M, Nelson J, Grix A, Sommer A, Savarirayan R, Janecke AR, Steichen E, Sillence D, Hausser I, Budde B, Nurnberg G, Nurnberg P, Seemann P, Kunkel D, Zambono G, Dallapiccola B, Schuelke M, Robertson S, Hamamy H, Wollnik B, Van Maldergem L, Mundlos S, Kornak U. 2009. Mutations in PYCR1 cause cutis laxa with progeroid features. *Nat Genet* 41:1016–1021.
- Sarbassov DD, Ali SM, Sengupta S, Sheen JH, Hsu PP, Bagley AF, Markhard AL, Sabatini DM. 2006. Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB. *Mol Cell* 22:159–168.

- Stockler S, Plecko B, Gospe SM, Jr., Coulter-Mackie M, Connolly M, van Karnebeek C, Mercimek-Mahmutoglu S, Hartmann H, Scharrer G, Struijs E, Tein I, Jakobs C, Clayton P, Van Hove JL. 2011. Pyridoxine dependent epilepsy and antiquitin deficiency: Clinical and molecular characteristics and recommendations for diagnosis, treatment and followup. *Mol Genet Metab* 104:48–60.
- Struys EA, Jakobs C. 2010. Metabolism of lysine in alpha-aminoacidipic semialdehyde dehydrogenase-deficient fibroblasts: Evidence for an alternative pathway of pipecolic acid formation. *FEBS Lett* 584:181–186.
- Takanaga H, Mackenzie B, Suzuki Y, Hediger MA. 2005. Identification of mammalian proline transporter SIT1 (SLC6A20) with characteristics of classical system imino. *J Biol Chem* 280:8974–8984.
- Tato I, Bartrons R, Ventura F, Rosa JL. 2011. Amino acids activate mammalian target of rapamycin complex 2 (mTORC2) via PI3 K/Akt signaling. *J Biol Chem* 286:6128–6142.
- Taylor NL, Heazlewood JL, Day DA, Millar AH. 2004. Lipoic acid-dependent oxidative catabolism of alpha-keto acids in mitochondria provides evidence for branched-chain amino acid catabolism in *Arabidopsis*. *Plant Physiol* 134:838–848.
- Van Schaftingen E, Rzem R, Marbaix A, Collard F, Veiga-da-Cunha M, Linster CL. 2013. Metabolite proofreading, a neglected aspect of intermediary metabolism. *J Inher Metab Dis* 36:427–434.
- Vranova V, Lojkova L, Rejsek K, Formanek P. 2013. Significance of the natural occurrence of L- versus D-pipecolic acid: A review. *Chirality* 25:823–831.
- Wanders RJ, Romeyn GJ, Schutgens RB, Tager JM. 1989. L-pipecolate oxidase: A distinct peroxisomal enzyme in man. *Biochem Biophys Res Commun* 164:550–555.
- Wanders RJ, Romeyn GJ, van Roermund CW, Schutgens RB, van den Bosch H, Tager JM. 1988. Identification of L-pipecolate oxidase in human liver and its deficiency in the Zellweger syndrome. *Biochem Biophys Res Commun* 154:33–38.
- Wanduragala S, Sanyal N, Liang XW, Becker DF. 2010. Purification and characterization of Put1p from *Saccharomyces cerevisiae*. *Arch Biochem Biophys* 498:136–142.
- Wolf J, Schliebs W, Erdmann R. 2010. Peroxisomes as dynamic organelles: Peroxisomal matrix protein import. *FEBS J* 277:3268–3278.
- Yogev O, Pines O. 2011. Dual targeting of mitochondrial proteins: Mechanism, regulation and function. *Biochim Biophys Acta* 1808:1012–1020.
- Zarse K, Schmeisser S, Groth M, Priebe S, Beuster G, Kuhlow D, Guthke R, Platzer M, Kahn CR, Ristow M. 2012. Impaired insulin/IGF1 signaling extends life span by promoting mitochondrial L-proline catabolism to induce a transient ROS signal. *Cell Metab* 15:451–465.
- Zhang L, Alfano JR, Becker DF. 2015. Proline metabolism increases katG expression and oxidative stress resistance in *Escherichia coli*. *J Bacteriol* 197:431–440.
- Zhu W, Gincherman Y, Docherty P, Spilling CD, Becker DF. 2002. Effects of proline analog binding on the spectroscopic and redox properties of PutA. *Arch Biochem Biophys* 408:131–136.
- Zoncu R, Bar-Peled L, Efeyan A, Wang S, Sancak Y, Sabatini DM. 2011. mTORC1 senses lysosomal amino acids through an inside-out mechanism that requires the vacuolar H(+)-ATPase. *Science* 334:678–683.