

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Nebraska Beef Cattle Reports

Animal Science Department

---

January 2001

## A Diagnostic Strategy to Classify Pens of Feedlot Cattle by the Prevalence of *Escherichia coli* O157:H7 Fecal Shedding

David R. Smith

*University of Nebraska-Lincoln*, [dsmith8@unl.edu](mailto:dsmith8@unl.edu)

Spring Younts

*University of Nebraska-Lincoln*

Mark Blackford

*University of Nebraska-Lincoln*

Todd Milton

*University of Nebraska-Lincoln*

Rodney A. Moxley

*University of Nebraska-Lincoln*, [rmoxley1@unl.edu](mailto:rmoxley1@unl.edu)

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/animalscibcr>



Part of the [Animal Sciences Commons](#)

---

Smith, David R.; Younts, Spring; Blackford, Mark; Milton, Todd; Moxley, Rodney A.; Gray, Jeff; Hungerford, Laura; and Klopfenstein, Terry J., "A Diagnostic Strategy to Classify Pens of Feedlot Cattle by the Prevalence of *Escherichia coli* O157:H7 Fecal Shedding" (2001). *Nebraska Beef Cattle Reports*. 320. <https://digitalcommons.unl.edu/animalscibcr/320>

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Beef Cattle Reports by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

David R. Smith, Spring Younts, Mark Blackford, Todd Milton, Rodney A. Moxley, Jeff Gray, Laura Hungerford, and Terry J. Klopfenstein

Pens with the greatest percentage of cattle shedding *E. coli* O157:H7 were more likely to be wet and muddy at the time of sample collection. Pens with dusty conditions were intermediate in the percentage of cattle shedding the organism. The association between the environmental condition of a pen and the percentage of cattle shedding *E. coli* O157:H7 seems plausible. Compared to the normal pen conditions, muddy or dusty conditions would be expected to facilitate fecal-oral transmission of enteric agents because of greater opportunity for the agent to move with water or dust particles.

It is unlikely that selection bias was introduced by voluntary recruitment of the feedyards or selection of cattle pens by convenience. There was no prior knowledge of the *E. coli* O157:H7 status of any of the feedyards. Pens were selected for inclusion in the study with consideration for the feedyards' re-implanting schedule and the workload of the laboratories. When there was an option, pens with fewer cattle were preferentially chosen for study to minimize costs. Pens were selected without knowledge of the results from previous pens in the feedyards.

The results of this study suggest that *E. coli* O157:H7 should be considered an ubiquitous organism in pens of feedlot cattle and that factors in the pen environment may help to explain the prevalence of cattle shedding the organism. The limited time-period of the study (summer months) and the cross-sectional nature of the study did not permit observing the effect of time dependent variables on the outcome of pen prevalence. It would be interesting to observe changes in pen prevalence over time as pen conditions change.

---

<sup>1</sup>David Smith, assistant professor, Veterinary and Biomedical Sciences, Lincoln; Mark Blackford, graduate student, Animal Science, Lincoln; Spring Younts, graduate student, Veterinary and Biomedical Sciences, Lincoln; Todd Milton, assistant professor, Animal Science, Lincoln; Rodney Moxley, professor, Jeff Gray, assistant professor, Veterinary and Biomedical Sciences, Lincoln; Laura Hungerford, associate professor, Great Plains Veterinary Educational Center, Clay Center; Terry Klopfenstein, professor, Animal Science, Lincoln.

# A Diagnostic Strategy to Classify Pens of Feedlot Cattle by the Prevalence of *Escherichia coli* O157:H7 Fecal Shedding

David Smith  
Spring Younts  
Mark Blackford  
Rodney Moxley  
Jeff Gray  
Laura Hungerford  
Todd Milton  
Terry Klopfenstein<sup>1</sup>

This diagnostic strategy can be used in food-safety research or as a monitoring tool in animal production food-safety programs to classify feedlot pens by the percentage of cattle shedding *Escherichia coli* O157:H7.

## Summary

*This study evaluated two pen testing strategies to predict the percentage of cattle in a pen shedding detectable E. coli O157:H7. Culture of a composite fecal sample most accurately detected pens with 37% or more cattle shedding E. coli O157:H7 in feces. A new pen test device most accurately detected pens with 16% or more individuals shedding. The likelihood of detecting E. coli O157:H7 with either method increased as pen prevalence increased. If both pen-level test methods were used together, pens could be classified as high, medium or low prevalence with less labor and expense than testing individual cattle.*

## Introduction

The principles of hazard-analysis-critical-control-points (HACCP) were developed to minimize the likelihood that food will be contaminated with potentially dangerous pathogens. Ideally

food-safety would be maximized if HACCP principles were applied at all levels of food production and processing. Unfortunately, there is insufficient knowledge of the epidemiology and ecology of *E. coli* O157:H7 to design and implement HACCP-based food safety programs in cattle feedyards.

Research or development of on-farm HACCP programs to control *E. coli* O157:H7 in feedlot production systems have been hampered by difficulty in determining the infection status of cattle at any point in time. The difficulty in diagnosis results because infection with *E. coli* O157:H7 in cattle occurs without clinical signs, except in calves, and because there is a lack of field-validated methods to monitor livestock for food safety pathogens.

Determining if individual live cattle are shedding *E. coli* O157:H7 is expensive and impractical. For example, culture of the feces from most, if not all, animals in a feedlot pen requires considerable labor and supplies. Handling finished cattle prior to shipping is not desirable because of the loss in value to cattle due to shrink, dark cutters and bruising. It may be possible to control *E. coli* O157:H7 in feedlots without knowing the infection status of individual cattle because control points or interventions for reducing human food-borne pathogens in feedlot cattle would most likely be directed towards pens of cattle. Therefore, the *E. coli* O157:H7 status of pens of feedlot cattle is an important outcome for feedlot production food safety research and HACCP monitoring.

Research and development of HACCP-based feedlot food safety programs could advance if pens of cattle, rather than individuals, could be accurately and economically classified by




Figure 1. The relationship between the *Escherichia coli* O157:H7 culture results of the pen-test device, a composite fecal sample collected from the pen and the percent of cattle in the pen shedding detectable levels of the organism in rectal feces.

the level of fecal shedding of *E. coli* O157:H7. Such a “pen-test” could serve as a monitoring tool in feedlot production food-safety programs, and would allow researchers to test potential interventions, or look for feedlot production methods related to the presence or absence of food-borne pathogens. The objective of this study was to evaluate diagnostic strategies to efficiently identify pens of feedlot cattle with a high prevalence of cattle shedding *E. coli* O157:H7.

### Procedure

Twenty-nine feedlot pens from five Midwestern feedlots, ranging in size from 36 to 231 (median 107) cattle, were each studied once between June and September, 1999 (Smith et al. 2001 Beef Report). Seven pen-test devices that cattle could rub, lick or chew were placed in the pens the evening prior to sample collection. The morning of sampling, feces were collected from the rectums of

all cattle in each pen and concurrent samples were collected of pen-test devices and a single composite sample of 20 fresh fecal pats from the pen surface. Culture methods were specific to the type of sample but included selective enrichment and immunomagnetic separation. Isolates were confirmed by standard methods including PCR. Non-parametric statistical methods were used to test either rank differences or rank correlations between pen-level classifications and the results of individual animal testing.

### Results

*Escherichia coli* O157:H7 was isolated from at least one animal in each of the 29 pens. The percentage of cattle shedding detectable levels of the organism within a pen ranged from 0.7% to 79.8% (median 17.1%). *E. coli* O157:H7 was recovered from at least one pen-test device from 15 pens and from the composite fecal samples of eight pens

(Figure 1). Recovery of *E. coli* O157:H7 from at least one pen test device or from the composite fecal sample was most likely to occur from the pens with higher prevalence (Wilcoxon rank sums  $P=0.001$ ).

The pen-test devices and composite feces were evaluated singly as diagnostic tools to differentiate high prevalence pens from low prevalence pens. The new pen-test device most accurately detected as positive (greatest percentage of pens classified correctly) pens with 16% or greater prevalence (pen-level sensitivity = 82%, pen-level specificity = 92%). Culture of composite feces most accurately detected as positive pens with a 37% percent or higher prevalence (pen-level sensitivity = 86%, pen-level specificity = 91%).

Information from culture of the pen-test devices and the composite feces was combined to classify pens by three levels of fecal shedding prevalence. Pens were classified as high prevalence if *E. coli*

(Continued on next page)

O157:H7 was recovered from the composite fecal sample; pens were classified as medium prevalence if the organism was recovered from the device, but not from the composite feces; pens were classified as low prevalence if the organism was not recovered from the device or the composite feces. This classification scheme correlated well (Spearman's  $r = 0.76$ ,  $P < 0.0001$ ) with the pen-prevalence determined by culturing the feces from individual cattle. Pens classified as high prevalence had significantly higher rankings in pen-prevalence than pens classified as medium prevalence

( $P = 0.05$ ) or low prevalence ( $P = 0.0006$ ), and pens classified as medium prevalence had significantly higher rankings in pen-prevalence than pens classified as low prevalence ( $P = 0.005$ ).

The premise of the pen-test was to culture a few devices from which many cattle in a pen could have contributed organisms. Culture of the pen-test devices alone or in parallel with culture of a composite fecal sample may be a diagnostically efficient strategy to characterize *E. coli* O157:H7 fecal shedding in feedlot pens. This diagnostic strategy may be useful as a research tool or as a

monitoring tool in the development of animal production food safety programs.

<sup>1</sup>David Smith, assistant professor, Veterinary and Biomedical Sciences, Lincoln; Spring Younts, graduate student, Veterinary and Biomedical Sciences, Lincoln; Mark Blackford, graduate student, Animal Science, Lincoln; Todd Milton, assistant professor, Animal Science, Lincoln; Rodney Moxley, professor, Veterinary and Biomedical Sciences, Lincoln; Jeff Gray, assistant professor, Veterinary and Biomedical Sciences, Lincoln; Laura Hungerford, associate professor, Great Plains Veterinary Educational Center, Clay Center; Terry Klopfenstein, professor, Animal Science, Lincoln.

---

## Influence of Restricted Intake and Reduced Dietary Starch on Colonic pH and *E. coli* Prevalence

Tim Loy  
Casey Wilson  
Doreen Bailey  
Mark Blackford  
Terry Klopfenstein  
Rod Moxley  
David Smith  
Spring Younts<sup>1</sup>

A finishing diet low in starch increased fecal pH, lowered VFA, and reduced numbers of acid-resistant *E. coli* shed in the feces.

### Summary

Ninety feedlot steers were used to test the effects of reducing dietary starch and intake on colonic pH, VFA, total and acid-resistant coliform and *E. coli* populations, and *E. coli* O157:H7 shedding. When corn bran and wet corn gluten feed (WCGF), or high moisture corn and WCGF were substituted for dry rolled corn, colonic pH increased while VFA concentration decreased. The corn bran and WCGF diet reduced acid-resistant *E. coli* shedding. Restricting intake increased colonic pH and decreased VFA con-

centration, but did not affect acid-resistant *E. coli* shedding. Prevalence of *E. coli* O157:H7 was not affected by diet or intake.

### Introduction

Enterohemorrhagic *Escherichia coli* O157:H7 is a bacterium found commonly in the intestinal tract of livestock that can cause severe illness and death in humans. More than 100 outbreaks of *E. coli* O157:H7 have been reported since 1982, 52% of which have been linked to foods derived from cattle.

Besides its ubiquitous distribution in livestock, other noteworthy characteristics of *E. coli* O157:H7 are its low infective dose for humans and its acid resistance. Because it can thrive under low pH conditions, undigested feed that is fermented in the colon may facilitate growth of *E. coli* O157:H7 and ultimately increase the numbers of the organism being shed in the feces.

High grain finishing diets may result in large amounts of undigested starch reaching the colon. Because of this, it has been suggested that feeding hay instead of grain would decrease the amount of starch reaching the colon,

increase colonic pH and decrease the numbers of acid-resistant *E. coli* being shed in the feces. In 1998, it was reported that switching cattle from a grain-based diet to hay four days prior to slaughter reduced the prevalence of both generic and acid resistant *E. coli*. A similar study conducted at the University of Nebraska confirmed these results. When steers consuming dry-rolled corn, high-moisture corn, or wet corn gluten feed were switched to alfalfa hay for five days, acid-resistant *E. coli* populations in the feces were reduced by 99% (2000 Nebraska Beef Report, pp. 39-41).

Feeding hay at the end of the feeding period may not be a feasible management practice for cattle feeders. However, if reducing the amount of fermentation in the colon is the key to reducing the numbers of acid-resistant *E. coli* being shed in the feces, more practical approaches may be available. High moisture corn, wet corn gluten feed (WCGF) and corn bran are feedstuffs used commonly in Nebraska feedlots, and each would be expected to result in less undigested starch reaching the colon. Therefore, the objectives of this trial were to determine the effect of