

1994

# FECUNDITY AND FERTILITY OF RHYNCHOPHORUS CRUENTATUS (COLEOPTERA: CURCULIONIDAE)

Thomas J. Weissling

*University of Nebraska-Lincoln*, [tweissling2@unl.edu](mailto:tweissling2@unl.edu)

Robin M. Giblin-Davis

*University of Florida*

Follow this and additional works at: <http://digitalcommons.unl.edu/entomologyfacpub>



Part of the [Entomology Commons](#)

---

Weissling, Thomas J. and Giblin-Davis, Robin M., "FECUNDITY AND FERTILITY OF RHYNCHOPHORUS CRUENTATUS (COLEOPTERA: CURCULIONIDAE)" (1994). *Faculty Publications: Department of Entomology*. 320.  
<http://digitalcommons.unl.edu/entomologyfacpub/320>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

FECUNDITY AND FERTILITY OF *RHYNCHOPHORUS*  
*CRUENTATUS*  
(COLEOPTERA: CURCULIONIDAE)

THOMAS J. WEISSLING<sup>1</sup> AND ROBIN M. GIBLIN-DAVIS

University of Florida, Institute of Food and  
Agricultural Sciences, Ft. Lauderdale Research and Education  
Center, 3205 College Ave., Ft. Lauderdale, FL 33314

The palmetto weevil (*R. cruentatus* F.) breeds in a variety of stressed or dying palms (Giblin-Davis & Howard 1988, 1989). These large (1.9 - 3.0 cm long) weevils are associated with the native cabbage palmetto, *Sabal palmetto* (Walter), in Florida (Woodruff 1967). Semiochemicals emanating from stressed or dying palms and male conspecifics (Weissling et al. 1992, 1993, 1994, Giblin-Davis et al. 1994) are attractive to *R. cruentatus* adults. Although not fully understood, mating apparently takes place on dying palms and females lay their eggs in the leaf bases or directly into the wounds of the host. Larvae develop primarily in the crown region but can occasionally be found in the stem tissue. Last instar larvae migrate to the fibrous stem periphery or petiolar bases and construct cocoons from fiber (Giblin-Davis & Howard 1988).

Research on an improved method to culture *R. cruentatus* has required the collection of a large number of eggs to produce neonate larvae for evaluation of diets. Using pineapple, *Anana comosus* (L.), as an ovipositional substrate, Giblin-Davis et al. (1989) reported the mean lifetime fecundity of field-collected females as  $26 \pm 15$  eggs per female. However, pineapple proved to be a difficult media to dissect for removal of eggs. More suitable ovipositional substrates were investigated and we found that apple (*Pyrus malus* L.) slices were easily dissected and were readily accepted by *R. cruentatus* females. Using apple slices, we reinvestigated the fecundity of *R. cruentatus* females and determined fertility.

Cocoons were harvested in the field from infested palms or in the laboratory from sugarcane (*Saccharum officinarum* L.) stem (Giblin-Davis et al. 1989). Cocoons were placed individually in covered 100-ml plastic cups with moistened tissue paper (Giblin-Davis et al. 1989) and were stored at 29°C until adult emergence. One male and one female were placed in a 500-ml covered plastic container with moistened tissue. After 48-72 h, males were removed and a slice of apple ('Red Delicious') was added. Slices were thin (5-15 mm; 10-18 g wet weight), convex segments covered by peel. Females oviposit through the apple pulp and most eggs are found along the peel. All containers were placed in an environmental chamber at 29°C with a 13:11 (L:D) photoperiod. Apple slices were usually replaced every 1-3 days until female death. Slices removed from containers were carefully dissected and eggs were removed. This test was repeated five times with four or five females per test (22 females total). Data were converted to the number of eggs laid per female per week. In addition, the total number of eggs laid per female was determined. During two of the tests, eggs removed from apple slices were placed in 15 x 100 mm plastic petri dishes lined with water-moistened filter paper, sealed with parafilm, and placed in the environmental chamber. Neonate larvae were removed from each dish at daily intervals and the dish resealed until all eggs hatched or decomposed. For the first test, fertility was determined every one to three days for 45 d. During the second test, eggs were col-

<sup>1</sup>Current address: Yakima Agricultural Research Laboratory, USDA-ARS, 3706 W. Nob Hill Blvd., Yakima, WA 98902

lected every one to three weeks for 14 weeks. Data on overlapping dates were combined and all data were converted to percent eclosion.

*R. cruentatus* produced an average of  $207 \pm 19$  (range 12-433) eggs per female. This number is considerably higher than the previous estimate of  $26 \pm 15$  eggs per female (Giblin-Davis et al. 1989) and is more consistent with fecundity reported for other *Rhynchophorus* species (Wattanapongsiri 1966). A reduced estimate of fecundity by Giblin-Davis et al. (1989) may have been caused by suboptimal conditions on pineapple slices, decreasing egg and larval survival. In addition, pineapple slices were difficult to dissect and eggs or larvae may have been overlooked (R.M.G.-D. pers. observ.). In this study, males were removed from containers after 48 - 72 h while Giblin-Davis et al. (1989) confined females with males throughout the course of the study. The presence of males in a small container may have interfered with oviposition or increased damage to larvae and eggs (Giblin-Davis et al. 1989). Rananavare et al. (1975) determined that *R. ferrugineus* Oliver females laid less eggs when confined with males than without. The mean number of eggs laid per *R. cruentatus* female per day declined until almost no eggs were laid 14 weeks after mating (Fig. 1). However, egg laying by surviving females increased over the subsequent 6 weeks. Reasons for this increase are unclear.

Fertility of eggs laid by *R. cruentatus* varied through time, however, eggs collected eight weeks after female mating did not eclose (Fig. 2). These results suggest that females had utilized all sperm indicating the need for multiple matings. Rananavare et al. (1975) reported the need for multiple matings in *R. ferrugineus* to maintain fertility.




Fig. 1. Mean ( $\pm$  SEM) weekly egg production by newly-emerged *R. cruentatus* females ( $n = 22$ ) confined individually on an apple slice at  $29^{\circ}\text{C}$  after 48 - 72h confinement with males.




Fig. 2. Percent eclosion of eggs produced by newly-emerged *R. cruentatus* females (n = 9).

We thank J. Cangiamila for technical assistance, and R.H. Scheffrahn and F.W. Howard for their suggestions. This research was supported by a USDA Special Grant in Tropical and Subtropical Agriculture CRSR-90-34135-5233. This manuscript is Florida Agricultural Research Stations Journal Series R-03674.

### Summary

The mean fecundity of *R. cruentatus* when provided apple slices as an oviposition media was almost eight times higher ( $207 \pm 19$  eggs per female) than previously estimated ( $26 \pm 15$  eggs per female). The rate of egg-laying decreased through time until 14 weeks after mating when there was a temporary increase. Fertility of *R. cruentatus* eggs remained between 40 and 100 percent eclosion until seven weeks after females were mated but dropped to zero by nine weeks.

### REFERENCES CITED

- GIBLIN-DAVIS, AND F. W. HOWARD. 1988. Notes on the palmetto weevil, *Rhynchophorus cruentatus* (Coleoptera: Curculionidae). Proc. Florida State Hort. Soc. 101: 101-107.
- GIBLIN-DAVIS, R. M., AND F. W. HOWARD. 1989. Vulnerability of stressed palms to attack by *Rhynchophorus cruentatus* (Coleoptera: Curculionidae) and insecticidal control of the pest. J. Econ. Entomol. 82: 1185-1190.
- GIBLIN-DAVIS, R. M., K. GERBER, AND R. GRIFFITH. 1989. Laboratory rearing of *Rhynchophorus cruentatus* and *R. palmarum* (Coleoptera: Curculionidae). Florida Entomol. 72: 480-488.

- GIBLIN-DAVIS, R. M., T. J. WEISSLING, A. C. OEHLISCHLAGER, AND L. M. GONZALEZ. 1994. Field response of *Rhynchophorus cruentatus* (F.) (Coleoptera: Curculionidae) to its aggregation pheromone and fermenting plant volatiles. Florida Entomol. 77: 164-177.
- RANANAVARE, H. D., K. SHANTARUM, M. R. HARWALKAR, AND G. W. RAHLKAR. 1975. Method of laboratory rearing of red palm weevil, *Rhynchophorus ferrugineus* Oliv. J. Plantation Crops 3: 65-67.
- WATTANAPONGSIRI, A. 1966. A revision of the genera *Rhynchophorus* and *Dynamis* (Coleoptera: Curculionidae). Dept. Agric. Science Bulletin, Bangkok 1: 1-328.
- WEISSLING, T. J., R. M. GIBLIN-DAVIS, R. H. SCHEFFRAHN, AND N. M. MENDOZA. 1992. Trap for capturing and retaining *Rhynchophorus cruentatus* (Coleoptera: Curculionidae) adults using *Sabal palmetto* as bait. Florida Entomol. 75: 212-221.
- WEISSLING, T. J., R. M. GIBLIN-DAVIS, AND R. H. SCHEFFRAHN. 1993. Laboratory and field evidence for male-produced aggregation pheromone in *Rhynchophorus cruentatus* (F.) (Coleoptera: Curculionidae). J. Chem. Ecol. 19: 1195-1203.
- WEISSLING, T. J., R. M. GIBLIN-DAVIS, G. GRIES, R. GRIES, A. L. PEREZ, H. D. PIERCE, JR., AND A. C. OEHLISCHLAGER. 1994. Aggregation pheromone of palmetto weevil *Rhynchophorus cruentatus* (Coleoptera: Curculionidae). J. Chem. Ecol. 20: 505-515.
- WOODRUFF, R. E. 1967. A giant palm weevil, *Rhynchophorus cruentatus* (Fab.), in Florida (Coleoptera: Curculionidae). Florida Dept. Agr. Div. Plant Industry. Entomology Circ. No. 63.

