University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications: Department of
Entomology

Entomology, Department of

8-2012

Effect of distribution and concentration of topically applied neonicotinoid insecticides in buffalograss, *Buchloe dactyloides*, leaf tissues on the differential mortality of *Blissus occiduus* under field conditions

Mitchell D. Stamm

University of Nebraska-Lincoln, mstamm3@unl.edu

Tiffany M. Heng-Moss

University of Nebraska-Lincoln, thengmoss2@unl.edu

Frederick P. Baxendale

University of Nebraska-Lincoln, fbaxendale1@unl.edu

Blair D. Siegfried

University of Nebraska-Lincoln, bsiegfried1@unl.edu

Roch E. Gaussoin

University of Nebraska-Lincoln, rgaussoin1@unl.edu

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/entomologyfacpub>

Part of the [Entomology Commons](#)

Stamm, Mitchell D.; Heng-Moss, Tiffany M.; Baxendale, Frederick P.; Siegfried, Blair D.; Gaussoin, Roch E.; Snow, Daniel D.; and Cassada, David A., "Effect of distribution and concentration of topically applied neonicotinoid insecticides in buffalograss, *Buchloe dactyloides*, leaf tissues on the differential mortality of *Blissus occiduus* under field conditions" (2012). *Faculty Publications: Department of Entomology*. 334. <https://digitalcommons.unl.edu/entomologyfacpub/334>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Mitchell D. Stamm, Tiffany M. Heng-Moss, Frederick P. Baxendale, Blair D. Siegfried, Roch E. Gaussoin, Daniel D. Snow, and David A. Cassada

Effect of distribution and concentration of topically applied neonicotinoid insecticides in buffalograss, *Buchloe dactyloides*, leaf tissues on the differential mortality of *Blissus occiduus* under field conditions

Mitchell D. Stamm,¹ Tiffany M. Heng-Moss,¹ Frederick P. Baxendale,¹ Blair D. Siegfried,¹ Roch E. Gaussoin,² Daniel D. Snow,³ and David A Cassada³

1. Department of Entomology, University of Nebraska-Lincoln, Lincoln, NE, USA

2. Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, NE, USA

3. School of Natural Resources, Water Center, University of Nebraska-Lincoln, Lincoln, NE, USA

Corresponding author — Mitchell D. Stamm, Department of Entomology, University of Nebraska-Lincoln, 103 Entomology Hall, Lincoln, NE 68583; email mstamm3@unl.edu

Abstract

BACKGROUND: Neonicotinoid insecticides are generally efficacious against many turfgrass pests, including several important phloem-feeding insects. However, inconsistencies in control of western chinch bugs, *Blissus occiduus*, have been documented in field efficacy studies. This research investigated the efficacy of three neonicotinoid insecticides (clothianidin, imidacloprid and thiamethoxam) against *B. occiduus* in buffalograss under field conditions and detected statistically significant differences in *B. occiduus* numbers among treatments. A subsequent study documented the relative quantity and degradation rate of these insecticides in buffalograss systemic leaf tissues, using HPLC.

RESULTS: Neonicotinoid insecticides initially provided significant reductions in *B. occiduus* numbers, but mortality diminished over the course of the field studies. Furthermore, while all three neonicotinoids were present in the assayed buffalograss leaf tissues, imidacloprid concentrations were significantly higher than those of clothianidin and thiamethoxam. Over the course of the 28 day study, thiamethoxam concentrations declined 700-fold, whereas imidacloprid and clothianidin declined only 70-fold and 60-fold respectively.

CONCLUSIONS: Field studies continued to verify inconsistencies in *B. occiduus* control with neonicotinoid insecticides. This is the first study to document the relative concentrations of topically applied neonicotinoid insecticides in buffalograss systemic leaf tissues.

Keywords: *Blissus occiduus*, neonicotinoid insecticides, insecticide degradation, buffalograss

1. Introduction

For more than a decade, neonicotinoid insecticides have provided effective control of a wide array of insect pests. Since the introduction of imidacloprid in the 1990s, these compounds have represented one of the fastest growing insecticide classes in recent history.^{1, 2} The success of this and other neonicotinoid compounds has been due in large part to their broad-spectrum insecticidal activity, low application rates, systemic uptake and translocation in plants and favorable toxicological profile.³ While crop protection remains the major use for neonicotinoid insecticides,⁴ new markets have emerged for urban, veterinary, turf and ornamental pests.⁵ Neonicotinoid insecticides have been shown effectively to control a variety of turfgrass pests, including the chinch bugs *Blissus leucopterus hirtus* Montandon and *B. insularis* Barber (Shetlar DJ, <http://golfdom.com>).^{6–8}

Western chinch bug, *B. occiduus* Barber, control has been documented in field studies at the University of Nebraska-Lincoln. In these field studies on buffalograss, significant though inconsistent reductions in *B. occiduus* numbers were observed with imidacloprid and clothianidin.^{9–11} Other chemical, environmental or plant physiological factors may have been responsible for the inconsistent results.

Blissus occiduus has emerged as a serious buffalograss pest over the past two decades. This chinch bug is widely distributed, from California, Colorado, Kansas, Montana, Nebraska and New Mexico in the United States to Alberta, British Columbia, Manitoba and Saskatchewan in Canada.^{12–14} In Nebraska, two generations of *B. occiduus* occur annually. Adults overwinter within the turfgrass stand. The first generation completes development by mid-June and is present until mid-August. Second-gen-

eration adults appear in late August and remain active until fall temperatures cool.¹⁴ Chinch bugs injure grasses by withdrawing sap from stolons and plant tissues in the crown area.¹⁴

Laboratory bioassays further explored the intrinsic toxicity of the neonicotinoid insecticides. Clothianidin, imidacloprid and thiamethoxam are all toxic to *B. occiduus*, with both topical and systemic applications of these insecticides able to provide significant reductions in chinch bug numbers.¹⁵ The mortality observed for the systemically applied insecticides could only have occurred if they had been translocated at lethal concentrations through the roots and into the stems and above-ground leaf tissues where chinch bugs feed.¹⁵

Clothianidin, imidacloprid and thiamethoxam are in the same insecticide class, but differences in chemical structure can influence arthropod toxicity. Water solubility, for example, can influence insecticidal activity. Thiamethoxam has higher water solubility (4.1 g L^{-1}) than clothianidin ($0.30\text{--}0.34 \text{ g L}^{-1}$) and imidacloprid (0.61 g L^{-1}), suggesting it should have greater systemic movement and potential to dissolve in the vascular (xylem) tissues.⁵ This may result in greater toxicity at lower chemical concentrations in the plant. Systemicity, however, is only one of many biotic and abiotic environmental factors and plant responses that can influence insecticide toxicity under field conditions.

Insecticide translocation in plants generally occurs more rapidly in younger stem and leaf tissues.¹⁶ Xylem flow is driven by water transpiration from the leaves, which in turn is regulated by the stomata and varies relative to environmental factors, including light, temperature, wind, etc.¹⁷ The systemic properties of imidacloprid have been studied in numerous plant species, exhibiting upward movement in the xylem.^{18–20} This systemic movement can be explained in terms of its physicochemical properties. Imidacloprid, a polar and water-soluble compound, typically forms weak bonds with soil particles, making it available in the soil water for uptake by plant roots and transport in the xylem to leaf margins and interveinal spaces.²¹ However, the rate of insecticide uptake is variable among plant species. Soil properties may also influence neonicotinoid uptake in plants.^{22–24}

Currently, little information is available on the movement of systemic insecticide in turfgrasses, especially in buffalograss. This study documented the distribution and concentration of topically applied neonicotinoid insecticides (clothianidin, imidacloprid and thiamethoxam) in buffalograss leaf tissues and correlated these parameters with *B. occiduus* differential mortality under field conditions.

2. Experimental Methods

2.1 Field evaluations

Insecticides were applied to buffalograss in research plots located at the John Seaton Anderson Turf and Ornamental Research Facility (JSA Research Facility), University of Nebraska Agricultural Research and Development Center, near Mead, Nebraska. Soil content at the JSA Facility plots was primarily Tomek silt loam (pH 6.8–7.2). As mentioned, clay content may influence insecticide uptake in plants. Soil content at these sites was unlikely significantly to inhibit insecticide uptake in the plant. Plots were mowed weekly at 7.6 cm (clippings returned with weed content of < 5%), and subsequent irrigation was applied at $2.5 \text{ cm month}^{-1}$ to maintain optimal growing conditions. Cumulative rainfall during the studies was 29.7 cm (field study 1) and 20.3 cm (field study 2).

The plots were treated with the highest labeled rates of Arena 0.5 G [clothianidin, (*E*)-1-(2-chloro-1,3-thiazol-5-ylmethyl)-3-methyl-2-nitroguanidine] at 89.7 kg ha^{-1} ($0.45 \text{ kg AI clothianidin ha}^{-1}$), Merit 75 WP [imidacloprid, 1-[(6-chloro-3-pyridinyl) methyl]-*N*-nitro-2-imidazolidinimine] at 9.6 kg ha^{-1} ($0.45 \text{ kg AI imidacloprid ha}^{-1}$) and Meridian 25 WG [thiamethoxam, (*E*, *Z*)-3-(2-chloro-1,3-thiazol-5-ylmethyl)-5-methyl-1,3,5-oxadiazinan-4-ylidene(nitro) amine] at 19.1 kg ha^{-1} ($0.30 \text{ kg AI thiamethoxam ha}^{-1}$). Talstar One [bifenthrin, 2-methylbiphenyl-3-ylmethyl (*Z*)-(1*RS*,3*RS*)-3-(2-chloro-3,3,3-trifluoropro-1-enyl)-2,2-dimethyl-cyclopropanecarboxylate], a synthetic pyrethroid industry standard, served as a positive control and was applied at 1.6 L ha^{-1} ($0.11 \text{ kg AI bifenthrin ha}^{-1}$).

Two independent field studies were designed to evaluate the efficacy of these insecticides against first- and second-generation *B. occiduus*. Treatments were applied to third- and fourth-instar chinch bugs in both field studies. Pretreatment estimates were taken prior to each study. Chinch bug numbers in a 0.09 m^2 area ranged from 10 to 20 (field study 1) and from 35 to 45 (field study 2). Plots were $1.5 \text{ m} \times 1.5 \text{ m}$ arranged in a randomized complete block (RCB) design, with five replications. Meridian, Merit and Talstar were applied using a CO_2 sprayer at 276 kPa, with 1627 L ha^{-1} finished spray. Arena 0.5 G, a granular formulation, was uniformly applied using a hand shaker. Lack of product availability necessitated the substitution of Arena 0.5 G for the preferred 50 WDG formulation. Fortunately, efficacy trials conducted at the University of Nebraska-Lincoln had shown few differences in *B. occiduus* control between the 0.5 G and 50 WDG Arena formulations (Baxendale FB, unpublished). All plots were irrigated with approximately 0.38 cm of water immediately following application to activate the Arena granules, wash the liquid formulations off leaf surfaces and facilitate root uptake of the neonicotinoid insecticides.

Applications to first-generation chinch bugs (field study 1) were evaluated for efficacy at 3, 8 and 106 days after treatment (DAT), while the second generation (field study 2) was evaluated for efficacy 3, 7, 28 and 55 DAT. Two 24 cm diameter areas were sampled (0.09 m^2 total area per plot) by vacuuming the soil surface with an ECHO Shred 'N' Vac (Model No. 2400; ECHO Inc., Lake Zurich, IL).²⁵ At the conclusion of each sample date, the vacuum samples were placed individually in Berlese funnels for 48 h. The surviving chinch bugs were collected in 70% ethyl alcohol, sorted by age class and counted using a Leica Zoom 2000 microscope (Leica, Wetzlar, Germany). Treatments were compared with an untreated control for reductions in chinch bug numbers.

2.2 Materials and sample calibration

Individual stock solutions of all neonicotinoid analytes, the surrogate and the internal standard were prepared at concentrations of $5 \mu\text{g } \mu\text{L}^{-1}$ in methanol from analytical-grade clothianidin (99.4% AI), imidacloprid (99.5% AI), thiamethoxam (99.5% AI) (Chem Service Inc., West Chester, PA), terbutylazine (surrogate; Sigma-Aldrich, Milwaukee, WI) and $^{13}\text{C}_3$ -labeled atrazine (internal standard; Merck & Co., Whitehouse Station, NJ). Analyte, surrogate and internal standard calibration spiking solutions were prepared from the stock solutions diluted to $50 \text{ ng } \mu\text{L}^{-1}$ in methanol. Calibration standard samples were prepared from the calibration spiking solutions in sample matrix obtained from the method extraction of untreated buffalograss. Analytes and surrogate were added to individual calibration samples in amounts of

250, 1000 and 2500 ng to create a three-point calibration curve. Internal standard (2500 ng) was added to all calibration standards and samples to quantify analyte concentrations on the instrument. Mean percentage recovery of the surrogate from the 47 samples was $109 \pm 4\%$, which has met the acceptance criteria of the United States Environmental Protection Agency.²⁶ Analyte detection limits were estimated from the instrument signal-to-noise to be 32 ng mL^{-1} in the final injection matrix, corresponding to an analyte concentration in buffalograss of $0.01 \mu\text{g g}^{-1}$.

2.3 Extraction procedures

Insecticides were applied to 'Prestige' buffalograss refuge plots at the JSA Research Facility, which were independent of the previously described field studies. Application procedures followed the methods used in the field studies. Plots ($1.1 \text{ m} \times 1.2 \text{ m}$ in size) were treated using previously described insecticide rates. The experimental design was a RCBD design with four replications. Plots received a weekly mowing at 6.4 cm (clippings returned with weed content of $< 2\%$) and $\sim 2.5 \text{ cm}$ of irrigation and 9.3 cm of rainfall during the course of the 28 day study.

At 3, 7, 14 and 28 DAT, above-ground leaf material was harvested and stored on ice until being transferred to a -80°C freezer for later processing. A quantity of 5 g fresh weight of chilled leaf blades (stolons removed) was then randomly selected from the sample, frozen in liquid nitrogen and ground to a fine powder with a mortar and pestle. The ground tissue was transferred to 50 mL tubes (VWR International, San Diego, CA), and HPLC-grade acetonitrile (30 mL) (Sigma-Aldrich, Milwaukee, WI) was added to the material, which was then stored overnight (ca 15 h) at 4°C to extract the insecticides.

After extraction, samples were shaken for 30 min at 4°C on a multipurpose rotator (Model No. 2314; Thermo Scientific, Waltham, MA) and centrifuged at 3500 rpm at 16°C (IEC Multi-RF; Thermo Electron, Milford, MA) for 20 min. A 10 mL aliquot of the supernatant was mixed with 100 mL of reagent-grade water, and 2500 ng ($50 \mu\text{L}$ of a $50 \text{ ng } \mu\text{L}^{-1}$ solution) of terbutylazine was added as a surrogate. Aqueous extracts were passed through a 200 mg solid-phase extraction (SPE) cartridge (Oasis HLB; Waters, Milford, MA) connected to a vacuum manifold. The Oasis HLB cartridge used for SPE was previously prepared by sequential washing with 5 mL of acetonitrile, methanol and reagent-grade water.

Insecticides were eluted from the SPE cartridge with 2 mL of methanol into a disposable culture tube (13 mm in width by 100 mm in depth) (Fisher Scientific, Pittsburgh, PA), and 2500 ng ($50 \mu\text{L}$ of a $50 \text{ ng } \mu\text{L}^{-1}$ solution) of $^{13}\text{C}_3$ -labeled atrazine was added as an internal standard. The eluant was then evaporated at room temperature under a nitrogen flow to approximately 200–300 μL . The concentrated solution was diluted to a final volume of 500 μL with double-distilled water and filtered with a Mini-UniPrep™ syringeless filter ($0.45 \mu\text{m}$ pore size) (Whatman, Florham Park, NJ).

2.4 Instrument conditions

The following methods were adapted from a previous study conducted at the University of Nebraska-Lincoln.²⁷ The prepared aliquots (containing analyte, terbutylazine and $^{13}\text{C}_3$ -labeled atrazine) were analyzed by reverse-phase HPLC/MS/MS utilizing a Waters 2695 HPLC autosampler/pump coupled to a Finnegan LCQ (Thermo Scientific, Waltham, MA) ion-trap mass spectrometer. HPLC separation utilized a Luna C8 ($5 \mu\text{m}$ particle

size) column ($250 \text{ mm} \times 2 \text{ mm i.d.}$) (Phenomenex, Torrance, CA). The mobile phase was a 90:10 ratio of 0.1% (v/v) ammonium formate in water and 0.1% (v/v) ammonium formate in methanol for 2 min, followed by a 8 min linear gradient to a 20:80 mobile phase ratio, held for 12 min, then returned to a 90:10 ratio and held for another 10 min to re-equilibrate the column for a total run time of 30 min. The flow rate was 0.3 mL min^{-1} , and the sample injection volume was 25 μL . The LCQ mass spectrometer was operated in atmospheric-pressure chemical ionization (APCI) mode with the vaporizer temperature at 350°C , the discharge current at 5.0 μA , the sheath gas at 80 (arbitrary units), the auxiliary gas at 1 (arbitrary units), the tube lens voltage at -5.0 V , the capillary voltage at 3.0 V, the capillary temperature at 150°C , the lens voltage at -36.0 V , multipole 1 offset at -3.0 V , multipole 2 offset at -5.0 V and the multipole RF amplitude at 500 $\text{V}_{\text{p-p}}$. The daughter ion transitions and percentage collision energies used in the analysis for each analyte were as follows: imidacloprid ($m/z = 256 \rightarrow 210$, 30%), clothianidin ($m/z = 250 \rightarrow 169$, 25%), thiamethoxam ($m/z = 292 \rightarrow 211$, 25%), terbutylazine ($m/z = 230 \rightarrow 174$, 35%) and $^{13}\text{C}_3$ -labeled atrazine ($m/z = 219 \rightarrow 177$, 35%). The isolation width was 3 amu, and the activation time was 30 ms for all analytes. The collision gas was helium.

2.5 Statistical analysis

A generalized linear mixed model with a negative binomial distribution and randomized blocks was implemented in PROC GLIMMIX²⁸ for the two field studies. Chinch bug numbers in the Talstar One-treated plots were not included in the statistical analysis. Data obtained from HPLC were analyzed using PROC MIXED.²⁸ In all studies, when significant interactions were detected, *t*-tests were used for mean separation using Fisher's procedure ($\alpha = 0.05$).

3. Results

3.1 Field evaluations

Mean numbers of collected chinch bugs for field studies 1 and 2 are reported in Tables 1 and 2 respectively. For field study 1, analysis of variance suggested a significant one-way interaction for insecticide treatment ($F = 9.72$; $\text{df} = 3, 48$; $P < 0.0001$) and evaluation date ($F = 12.37$; $\text{df} = 3, 48$; $P < 0.0001$). Two-way interactions were not significant ($F = 1.52$; $\text{df} = 9, 48$; $P = 0.1906$). As one-way interactions were significant, simple effects were used to determine whether differences existed among treatment means. At 3 DAT, statistically significant differences in the number of chinch bugs were detected for all treatments when compared with the untreated control (clothianidin: $t = 2.36$, $\text{df} = 48$, $P < 0.0222$; thiamethoxam: $t = 2.27$, $\text{df} = 48$, $P < 0.0275$; imidacloprid: $t = 2.19$, $\text{df} = 48$, $P < 0.0336$). The treatments were equally effective at reducing chinch bug numbers, with no significant differences detected (clothianidin–thiamethoxam: $t = 0.10$, $\text{df} = 48$; $P = 0.9239$; clothianidin–imidacloprid: $t = 0.19$, $\text{df} = 48$; $P = 0.8520$; thiamethoxam–imidacloprid: $t = 0.09$, $\text{df} = 48$; $P = 0.9274$) (Table 1).

At 8 DAT, significant differences in the number of chinch bugs were detected in all treatments when compared with the untreated control (clothianidin: $t = 3.21$, $\text{df} = 48$, $P < 0.0023$; thiamethoxam: $t = 3.06$, $\text{df} = 48$, $P = 0.0036$; imidacloprid: $t = 2.92$, $\text{df} = 48$, $P = 0.0053$). However, no significant differences were detected among insecticide treatments, which was consistent with 3 day evaluations (clothianidin–thiamethoxam: $t = 0.21$, $\text{df} = 48$; $P = 0.8361$; clothianidin–imidacloprid: $t = 0.39$;

Table 1. Field study 1. Mean number of *Blissus occiduus* recovered from Berlese funnels after insecticide treatment

Treatment (active ingredient)	Mean number of chinch bugs ^a ± SEM ^b		
	3 DAT ^c	8 DAT	106 DAT
Clothianidin (Arena)	3.4±1.1 a	1.2±0.5 a	3.0±0.8 a
Thiamethoxam (Meridian)	3.6±2.5 a	1.4±0.5 a	15.8±1.8 b
Imidacloprid (Merit)	3.8±1.2 a	1.6±0.6 a	13.4±1.6 b
Untreated control	12.8±3.2 b	9.8±1.4 b	18.2±1.9 b

a. Mean number of chinch bugs per 0.09 m².b. Treatment means within the same column followed by the same letter indicate no significant differences ($P \leq 0.05$), LSD test.

c. DAT = days after treatment.

df = 48; $P = 0.6949$; thiamethoxam–imidacloprid: $t = 0.19$; df = 48; $P = 0.8522$) (Table 1).

Evaluations conducted at 106 DAT for field study 1 assessed the residual activity of the neonicotinoid insecticides against second-generation *B. occiduus* nymphs and adults. In this case, clothianidin had significantly fewer chinch bugs than the untreated control and provided $\approx 84\%$ control (Arena: $t = 3.20$, df = 48, $P < 0.0024$). Chinch bug numbers in the thiamethoxam and imidacloprid plots were not significantly different from the untreated control (thiamethoxam: $t = 0.28$, df = 48, $P = 0.7842$; imidacloprid: $t = 0.59$, df = 48, $P < 0.5555$) (Table 1).

For field study 2, analysis of variance suggested a significant one-way interaction for insecticide treatment ($F = 17.16$; df = 3, 64; $P < 0.0001$) and evaluation date ($F = 10.41$; df = 3, 64; $P < 0.0001$). Two-way interactions were not significant ($F = 0.43$; df = 9, 64; $P = 0.9114$). As one-way interactions were significant, simple effects were used to determine whether significant differences existed. At 3 DAT, clothianidin was the only treatment to provide significant reductions in *B. occiduus* numbers when compared with the untreated control ($t = 3.28$, df = 64, $P < 0.0017$). Reductions in *B. occiduus* numbers were not significantly different among the insecticide treatments and untreated control (thiamethoxam: $t = 1.76$, df = 64, $P < 0.0832$; imidacloprid: $t = 1.49$; df = 64; $P < 0.1400$) (Table 2).

As in field study 1, at 7 DAT the clothianidin treatment had significantly fewer chinch bugs compared with the untreated control ($t = 4.75$, df = 64, $P < 0.0001$). The numbers of chinch bugs in the thiamethoxam and imidacloprid treatments de-

clined and were significantly less than in the untreated control (thiamethoxam: $t = 2.63$, df = 64, $P < 0.0106$; imidacloprid: $t = 2.11$, df = 80, $P = 0.0391$). Again, however, the clothianidin treatment had significantly fewer chinch bugs than the thiamethoxam treatment ($t = 2.50$, df = 64, $P = 0.0149$) and the imidacloprid treatment ($t = 2.97$, df = 64, $P < 0.0041$) (Table 2).

At 28 DAT, clothianidin had significantly fewer chinch bugs compared with the untreated control ($t = 3.24$, df = 64, $P < 0.0019$). Although clothianidin and thiamethoxam were not significantly different ($t = 1.75$, df = 64, $P = 0.0848$), thiamethoxam did not have significantly fewer chinch bugs than the untreated control ($t = 1.95$, df = 64, $P = 0.0555$). In addition, chinch bug numbers in the imidacloprid treatment were not significantly different from those in the untreated control ($t = 1.63$, df = 64, $P = 0.1073$) (Table 2).

Field study 2 evaluations were also conducted at 55 DAT to assess the residual activity of the three compounds against second-generation *B. occiduus* nymphs and adults. Clothianidin, with $\approx 88\%$ control, was the only treatment that was significantly different from the untreated control (clothianidin: $t = 3.12$, df = 64, $P < 0.0027$; thiamethoxam: $t = 0.90$, df = 64, $P = 0.3720$; imidacloprid: $t = 1.63$, df = 64, $P = 0.1073$) (Table 2). These results are similar to those of field study 1. Long-term, residual control with thiamethoxam and imidacloprid was low.

3.2 Relative quantity of neonicotinoids in treated buffalograss

Mean insecticide concentrations in the buffalograss leaf samples are reported in Table 3. The analysis of variance suggested a significant one-way interaction for insecticide treatment ($F = 4.31$; df = 2, 35; $P = 0.0212$) and sampling date ($F = 21.77$; df = 3, 35; $P < 0.0001$). Two-way interactions were not significant ($F = 1.95$; df = 6, 35; $P = 0.1000$). As one-way interactions were significant, simple effects were used to determine whether significant differences existed.

At 3 DAT, mean concentrations of imidacloprid, clothianidin and thiamethoxam were $14.24 \mu\text{g g}^{-1}$ (active ingredient plant material⁻¹), $6.48 \mu\text{g g}^{-1}$ and $6.20 \mu\text{g g}^{-1}$ respectively (Table 3). The imidacloprid concentration in the treated buffalograss was significantly greater than the clothianidin concentration ($t = 3.44$, df = 35, $P = 0.0015$) and the thiamethoxam concentration ($t = 3.85$, df = 35, $P = 0.0005$). These differences may be due to water solubility, formulation or application rates, but do not explain why clothianidin and thiamethoxam had similar concentrations ($t = 0.13$, df = 35, $P = 0.8998$) when clothianidin was applied at a much higher rate.

Table 2. Field study 2. Mean number of *Blissus occiduus* recovered from Berlese funnels after insecticide treatment

Treatment (active ingredient)	Mean number of chinch bugs ^a ± SEM ^b			
	3 DAT ^c	7 DAT	28 DAT	55 DAT
Clothianidin (Arena)	5.8 ± 1.1 a	1.0 ± 0.4 a	0.4 ± 0.3 a	2.2 ± 0.7 a
Thiamethoxam (Meridian)	15.6 ± 1.8 a	6.8 ± 1.2 b	2.2 ± 0.7 ab	10.4 ± 1.4 b
Imidacloprid (Merit)	18.4 ± 1.9 a	9.6 ± 1.4 b	2.8 ± 0.7 b	10.2 ± 1.4 b
Untreated control	45.8 ± 3.0 b	35.4 ± 2.7 c	8.4 ± 1.3 b	18.2 ± 1.9 b

a. Mean number of chinch bugs per 0.09 m².b. Treatment means within the same column followed by the same letter indicate no significant differences ($P \leq 0.05$), LSD test.

c. DAT = days after treatment.

Table 3. Relative quantity of neonicotinoids in treated buffalograss

Treatment	Mean concentration ^a ± SEM ^b			
	3 DAT	7 DAT	14 DAT	28 DAT
Imidacloprid	14.24 ± 3.68 a	3.25 ± 0.72 a	0.77 ± 0.17 a	0.20 ± 0.05 a
Clothianidin	6.48 ± 1.97 b	1.56 ± 0.61 a	0.66 ± 0.17 a	0.19 ± 0.06 a
Thiamethoxam	6.20 ± 0.70 b	0.60 ± 0.29 a	0.16 ± 0.05 a	ND ^c

a. Mean insecticide concentration ($\mu\text{g g}^{-1}$ fresh weight).

b. Treatment means within the same column followed by the same letter indicate no significant differences ($P \leq 0.05$), LSD test.

c. ND = not detected (i.e. less than detection limit).

By 7 DAT, the concentration of imidacloprid, clothianidin and thiamethoxam in leaf tissues had significantly declined (clothianidin: $t = 2.18$, $df = 35$, $P < 0.0359$; imidacloprid: $t = 5.26$, $df = 35$, $P < 0.0001$; thiamethoxam: $t = 2.68$, $df = 35$, $P = 0.0112$). Thiamethoxam had the greatest decline (≈ 10 -fold), with $0.60 \mu\text{g g}^{-1}$ of active ingredient remaining in the leaf tissues, as opposed to imidacloprid ($3.25 \mu\text{g g}^{-1}$) and clothianidin ($1.56 \mu\text{g g}^{-1}$) with a ≈ 4 -fold and ≈ 8 -fold decline respectively (Table 3). Mean concentrations of the three insecticides were not significantly different (clothianidin–imidacloprid: $t = 0.81$, $df = 35$, $P = 0.4243$; clothianidin–thiamethoxam: $t = 0.46$, $df = 35$, $P = 0.6497$; imidacloprid–thiamethoxam: $t = 1.27$, $df = 35$, $P = 0.2137$).

From 7 to 14 DAT there were no significant declines in mean insecticide concentrations (clothianidin: $t = 0.43$, $df = 35$, $P < 0.6702$; imidacloprid: $t = 1.19$, $df = 35$, $P < 0.2435$; thiamethoxam: $t = 0.21$, $df = 35$, $P = 0.8346$) (Table 3). Comparing the three insecticides, active ingredient concentrations in the treated buffalograss were not significantly different (clothianidin–imidacloprid: $t = 0.05$, $df = 35$, $P = 0.9590$; clothianidin–thiamethoxam: $t = 0.24$, $df = 35$, $P = 0.8125$; imidacloprid–thiamethoxam: $t = 0.29$, $df = 35$, $P = 0.7729$).

At 28 DAT, insecticides were still detectable and had not significantly declined since the previous sampling date (clothianidin: $t = 0.23$, $df = 35$, $P < 0.8226$; imidacloprid: $t = 0.27$, $df = 35$, $P < 0.7889$; thiamethoxam: $t = 0.07$, $df = 35$, $P = 0.9425$). The insecticide concentrations were not significantly different (clothianidin–imidacloprid: $t = 0.01$, $df = 35$, $P = 0.9937$; clothianidin–thiamethoxam: $t = 0.09$, $df = 35$, $P = 0.9322$; imidacloprid–thiamethoxam: $t = 0.09$, $df = 35$, $P = 0.9259$); however, thiamethoxam had declined ≈ 20 -fold and was less than its method detection limit. Imidacloprid and clothianidin had declined ≈ 4 -fold, with 0.20 and $0.19 \mu\text{g g}^{-1}$ respectively (Table 3).

Insecticide concentrations declined significantly over the 28 day study (clothianidin: $t = 2.79$, $df = 35$, $P = 0.0085$; imidacloprid: $t = 6.72$, $df = 35$, $P < 0.0001$; thiamethoxam: $t = 2.96$, $df = 35$, $P = 0.0055$). In this 25 day period (3–28 DAT), thiamethoxam concentrations declined 700-fold, whereas imidacloprid and clothianidin declined only 70-fold and 60-fold respectively.

Some metabolites of neonicotinoid insecticides are known to be toxic to arthropod pests.²⁹ Concentrations of imidacloprid and clothianidin metabolites were below the present estimated detection limit ($0.01 \mu\text{g g}^{-1}$) and unlikely to contribute to observed chinch bug control. However, clothianidin is a known metabolite of thiamethoxam³⁰ and was detected in samples from the thiamethoxam treatment at 3, 7 and 14 DAT ($0.36 \pm 0.15 \mu\text{g g}^{-1}$, $0.12 \pm 0.05 \mu\text{g g}^{-1}$ and $0.07 \pm 0.02 \mu\text{g g}^{-1}$ respectively).

4. Discussion and Conclusions

Field studies continue to support previous findings^{9–11} and continue to verify contradictions in chinch bug control with neonicotinoid insecticides. Initial reductions in chinch bug numbers were significant, which may be the result of contact toxicity. Mortality quickly declined, possibly owing to the translaminar and systemic movement of these compounds. Although clothianidin was applied as granular formulation without translaminar systemicity, it continued to be more effective for controlling *B. occiduus*, especially for season-long chinch bug control. As mentioned, it is unlikely that a WDG formulation would have shown differences in *B. occiduus* control (Baxendale FB, unpublished).

This is the first report documenting neonicotinoid insecticide concentrations in buffalograss leaf tissues. While statistical comparisons were not made between studies, interestingly, initial concentrations of imidacloprid (at 3 DAT) in the buffalograss leaf tissues were greater than those of clothianidin and thiamethoxam. However, field mortality appeared to be similar, possibly owing to water solubility or other chemical properties that influence the amount of residue on the leaf surface and in the systemic tissues of the plant. Similar findings were documented in laboratory bioassays, which indicated no differences in systemic toxicity among the three neonicotinoids to *B. occiduus* nymphs.¹⁵ Chemical differences became less apparent in only a matter of days, with the relative persistence of these compounds being similar. In final comparisons, all three insecticides were detected at similar concentrations 28 DAT, while *B. occiduus* field mortality differed at 106 and 55 DAT (field study 1 and field study 2 respectively). This suggests that the concentration of insecticide in leaf tissues may not be closely correlated with chinch bug mortality under field conditions.

The biochemical and metabolic pathways of the neonicotinoids, especially imidacloprid, have been well studied within treated plant tissues.^{31, 32} An olefin derivative has been identified as the predominant metabolite of imidacloprid, with moderate to high insecticidal activity.^{21, 33, 34} In the present study, using full-scan MS to characterize transformation products, metabolites of imidacloprid and clothianidin were not detected. However, clothianidin was detected as a metabolite of thiamethoxam, suggesting that buffalograss has the capacity to metabolize thiamethoxam to clothianidin. At the 3 DAT sampling period, the concentration of the clothianidin metabolite was a small percentage of the total active ingredient available in the plant tissues. However, by 14 DAT, clothianidin represented a much higher proportion of the total of both parent

and metabolite, and may have contributed to the overall effect of the thiamethoxam treatment, especially later in the growing season.

This research provides a better understanding of neonicotinoid toxicity and degradation under field conditions. Neonicotinoids have been shown to photodegrade,^{35, 36} and soil bacteria are capable of metabolizing these compounds.^{37, 38} These factors could limit *B. occiduus* control with certain neonicotinoid insecticides, and may have important implications for management of other turfgrass insect pests. Additional research is also needed to understand *B. occiduus* behavior in response to neonicotinoid exposure. Chinch bug probing location has been documented,³⁹ but no studies have explored the specific feeding patterns on neonicotinoid-treated plant tissues. Electrical penetration graphs (EPGs)^{40, 41} would be a valuable tool to document the effects of neonicotinoid insecticides on chinch bug feeding and behavior.

Acknowledgments

The authors gratefully acknowledge Leonardo Magalhaes for his assistance in the development of HPLC methods and their application, and Erin Blankenship for her assistance with the statistical analysis. They also thank Nicholas Aliano, Lannie Wit, Travis Prochaska and Rachael Fithian for technical help.

References

- 1 Elbert A, Becker B, Hartwig J and Erdelen C, Imidacloprid – A new systemic insecticide. *Pflschutz Nachr Bayer* 44:113–136 (1991).
- 2 Jeschke P and Nauen R, Neonicotinoids – From zero to hero in insecticide chemistry. *Pest Manag Sci* 64:1084–1098 (2008).
- 3 Maienfisch P, Huerlimann H, Rindlisbacher A, Gsell L, Dettwiler H, Haettenschwiler J, et al, The discovery of thiamethoxam: A second-generation neonicotinoid. *Pest Manag Sci* 57:165–176 (2001).
- 4 Elbert A, Haas M, Springer B, Thielert W and Nauen R, Mini-review. Applied aspects of neonicotinoid uses in crop protection. *Pest Manag Sci* 64:1099–1105 (2008).
- 5 Tomizawa M and Casida JE, Neonicotinoid insecticide toxicology: Mechanisms of selective action. *Annu Rev Pharmacol Toxicol* 45:247–268 (2005).
- 6 Heller PR and Walker R, *Curative Management of Hairy Chinch Bug with Experimental and Registered Formulations on Fine Fescue*. Online: <http://www.entsoc.org/>
- 7 Swier SR, Rollins A and Duncan J, Comparison of three pyrethroids, Meridian, and Merit against second generation hairy chinch bug. (2001). Online: <http://www.entsoc.org/>
- 8 Buss EA and Ruppert P, *Southern Chinch Bug Control with Clothianidin*. Online: <http://www.entsoc.org/>
- 9 Eickhoff TE, Carstens JD and Baxendale FP, Evaluation of Selected Insecticides for Chinch Bug Control on Buffalograss Lawn. (2001). Online: <http://www.entsoc.org/>
- 10 Anderson WG, Eickhoff TE and Heng-Moss TM, *Evaluation of Arena for Control of First Generation Chinch Bugs*. (2004). Online: <http://www.entsoc.org/>
- 11 Eickhoff TE, Baxendale RW, Toda L and Baxendale FP, *Residual Control of Chinch Bugs with Allectus Insecticide*. (2005). Online: <http://www.entsoc.org/>
- 12 Bird RD and Mitchener AV, Insects of the season 1949 in Manitoba. *Can Insect Pest Rev* 28:41 (1950).
- 13 Slater JA, *A Catalogue of the Lygaeidae of the World*. Vol. 2. University of Connecticut, Storrs, CT (1964).
- 14 Baxendale FP, Heng-Moss TM and Riordan TP, *Blissus occiduus* (Hemiptera: Lygaeidae): a chinch bug pest new to buffalograss turf. *J Econ Entomol* 92:1172–1176 (1999).
- 15 Stamm MD, Baxendale FP, Heng-Moss TM, Siegfried BD, Blankenship EE and Gaussoin RE, Dose–response relationships of clothianidin, imidacloprid, and thiamethoxam to *Blissus occiduus* (Hemiptera: Blissidae). *J Econ Entomol* 104:205–210 (2011).
- 16 Bennett SH, The behaviour of systemic insecticides applied to plants. *Annu Rev Entomol* 2:279–296 (1957).
- 17 Baur P, Arnold R, Giessler S, Mansour P and Vermeer R, Bio-availability of insecticides from O-TEQ® formulations: overcoming barriers for systemic active ingredients. *Pflschutz Nachr Bayer* 60:27–42 (2007).
- 18 Castle SJ, Byrne FJ, Bi JL and Toscano NC, Spatial and temporal distribution of imidacloprid and thiamethoxam in citrus and impact on *Homalodisca coagulata* populations. *Pest Manag Sci* 61:75–84 (2005).
- 19 Cowles RS, Montgomery ME and Cheah CAS-J, Activity and residues of imidacloprid applied to soil and tree trunks to control hemlock woolly adelgid (Hemiptera: Adelgidae) in forests. *J Econ Entomol* 99:1258–1267 (2006).
- 20 Buchholz A and Nauen R, Translocation and translaminar bio-availability of two neonicotinoid insecticides after foliar application to cabbage and cotton. *Pest Manag Sci* 58:10–16 (2001).
- 21 Westwood F, Bean KM, Dewar AM, Bromilow RH and Chamberlain K, Movement and persistence of [¹⁴C] imidacloprid in sugar-beet plants following application to pelleted sugar-beet seed. *Pestic Sci* 52:97–103 (1998).
- 22 Cox L, Koskinen W and Yen P, Sorption–desorption of imidacloprid and its metabolites in soils. *J Agric Food Chem* 45:1468–1472 (1997).
- 23 Cox L, Koskinen W, Celis R, Yen P, Hermosin M and Cornejo J, Sorption of imidacloprid on soil clay mineral and organic components. *Soil Sci Soc Am J* 62:911–915 (1998).
- 24 Cox L, Koskinen W and Yen P, Influence of soil properties on sorption–desorption of imidacloprid. *J Environ Sci Health B* 33:123–134 (1998).
- 25 Eickhoff TE, Heng-Moss TM, Baxendale FP and Foster JE, Levels of tolerance, antibiosis, and antixenosis among resistant buffalograsses and zoysiagrasses. *J Econ Entomol* 101:533–540 (2008).
- 26 Munch JW, Method 525.2 – Determination of organic compounds in drinking water by liquid–solid extraction and capillary column gas chromatography/mass spectrometry, revision 2.0. National Exposure Research Laboratory, Office of Research and Development, US Environmental Protection Agency, pp. 1–60 (1995).
- 27 Magalhaes LC, Hunt TE and Siegfried BD, Efficacy of neonicotinoid seed treatments to reduce soybean aphid populations under field and controlled conditions in Nebraska. *J Econ Entomol* 102:187–195 (2009).

- 28 SAS/STAT User's Guide, Version 9.1. SAS Institute, Cary, NC (2006).
- 29 Tomizawa M, Zhang N, Durkin KA, Olmstead MM and Casida JE, The neonicotinoid electronegative pharmacophore plays the crucial role in the high affinity and selectivity for the *Drosophila* nicotinic receptor: an anomaly for the nicotinoid cation- π interaction model. *Biochemistry* 42:7819–7827 (2003).
- 30 Nauen R, Ebbinghaus-Kintscher U, Salgado VL and Kaussmann M, Thiamethoxam is a neonicotinoid precursor converted to clothianidin in insects and plants. *Pestic Biochem Physiol* 76:55–69 (2003).
- 31 Sur R and Stork A, Uptake, translocation and metabolism of imidacloprid in plants. *Bull Insectol* 56:35–40 (2003).
- 32 Ford KA and Casida JE, Comparative metabolism and pharmacokinetics of seven neonicotinoid insecticides in spinach. *J Agric Food Chem* 56:10 168–10 175 (2008).
- 33 Nauen R, Tietjen K, Wagner K and Elbert A, Efficacy of plant metabolites of imidacloprid against *Myzus persicae* and *Aphis gossypii* (Homoptera: Aphididae). *Pestic Sci* 52:53–57 (1998).
- 34 Nauen R, Reckmann U, Armbrorst S, Stupp HP and Elbert A, Whitefly active metabolites of imidacloprid: biological efficacy and translocation in cotton plants. *Pestic Sci* 55:265–271 (1999).
- 35 Kagabu S and Medej S, Stability comparison of imidacloprid and related compounds under simulated sunlight, hydrolysis conditions, and to oxygen. *Biosci Biotech Biochem* 59:980–985 (1995).
- 36 Wamhoff H and Schneider V, Photodegradation of imidacloprid. *J Agric Food Chem* 47:1730–1734 (1999).
- 37 Pandey G, Dorriana SJ, Russella RJ and Oakeshott JG, Bio-transformation of the neonicotinoid insecticides imidacloprid and thiamethoxam by *Pseudomonas* sp. 1G. *Biochem Biophys Res Commun* 380:710–714 (2009).
- 38 Liu Z, Dai Y, Huang G, Gu Y, Ni J, Wei H, *et al*, Soil microbial degradation of neonicotinoid insecticides imidacloprid, acetamiprid, thiacloprid and imidaclothiz and its effect on the persistence of bioefficacy against horsebean aphid *Aphis craccivora* Koch after soil application. *Pest Manag Sci* 67:1245–1252 (2011).
- 39 Anderson WG, Heng-Moss TM, Baxendale FP, Baird LM, Sarath G and Higley L, Chinchbug (Hemiptera: Blissidae) mouthpart morphology, probing frequencies, and locations on resistant and susceptible germplasm. *J Econ Entomol* 99:212–221 (2006).
- 40 McLean DL and Kinsey MG, A technique for electronically recording aphid feeding and salivation. *Nature* 202:1358–1359 (1964).
- 41 Tjallingii WF, Electronic recording of penetration behaviour by aphids. *Entomol Exp Appl* 24:521–530 (1978).