

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and Plant
Health Inspection Service

October 2004

AVIAN USE OF ROADSIDE HABITAT IN THE SOUTHERN DRIFT PLAINS OF NORTH DAKOTA AND IMPLICATIONS FOR CATTAIL (*TYPHA* SPP.) MANAGEMENT

Bryan D. Safratowich

North Dakota State University, Fargo

George M. Linz

USDA/APHIS/WS National Wildlife Research Center, george.m.linz@aphis.usda.gov

William J. Bleier

North Dakota State University, Fargo

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Environmental Sciences Commons](#)

Safratowich, Bryan D.; Linz, George M.; and Bleier, William J., "AVIAN USE OF ROADSIDE HABITAT IN THE SOUTHERN DRIFT PLAINS OF NORTH DAKOTA AND IMPLICATIONS FOR CATTAIL (*TYPHA* SPP.) MANAGEMENT" (2004). *USDA National Wildlife Research Center - Staff Publications*. 353.
https://digitalcommons.unl.edu/icwdm_usdanwrc/353

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

AVIAN USE OF ROADSIDE HABITAT IN THE SOUTHERN DRIFT
PLAINS OF NORTH DAKOTA AND IMPLICATIONS FOR CATTAIL
(*TYPHA* SPP.) MANAGEMENT

Bryan D. Safratowich, Department of Biological Sciences, Stevens Hall, North
Dakota State University, Fargo, ND 58105 USA
E-mail: Bryan.Safratowich@ndsu.nodak.edu

George M. Linz, USDA/APHIS, National Wildlife Research Center, Great Plains
Field Station, 2110 Miriam Circle, Suite B., Bismarck, ND 58501 USA
E-mail: George.M.Linz@aphis.usda.gov

William J. Bleier, Department of Biological Sciences, Stevens Hall, North Dakota
State University, Fargo, ND 58105 USA
E-mail: William.Bleier@ndsu.nodak.edu

Abstract

We determined avian use of roadside rights-of-way to develop proper management strategies for the manipulation of roadside cattail. Cattail management is a technique used to reduce nesting and roosting habitat for problematic blackbird species, which might feed on sunflower crops in the vicinity of cattail-dominated wetlands. Thirty quarter sections (1 quarter section \approx 64.75 ha) located in the Southern Drift Plains of North Dakota served as our study units. Roadside habitat along two 0.5 mile (\approx 0.8 km) transects bordering these quarter sections was surveyed to assess avian use. Additionally, nest surveys were conducted to provide an index of breeding bird use of this roadside habitat. Finally, roadside habitat was surveyed for a number of different habitat variables. We found 49 different species during the surveys. Of the 2,529 birds found in this habitat, 1,479 (41.5%) were blackbirds. Blackbirds were also the primary nesting birds, contributing to 89% of the active nests found in roadside habitat. In terms of avian use and nesting, blackbirds, especially redwings, were the dominant bird species using roadside cattail. With proper management of roadside habitat, potential effects on non-blackbirds can be minimized, and nesting habitat for blackbirds can be reduced. Our data indicate that a loss in nesting habitat will likely result in fewer blackbirds and a subsequent reduction in sunflower damage.

Introduction

Experimental field studies on reducing dense cattail (*Typha* spp.) stands in wetlands results in a decrease in blackbird (red-winged blackbirds, yellow-headed blackbirds, and common grackles) use and, in some cases, an increase in use by waterfowl and other birds (Solberg and Higgins, 1989; Linz et al., 1994; Linz et al., 1996). With this reduction in blackbird use, we see a concomitant drop in damage to nearby sunflower fields. If not protected, sunflower fields

adjacent to wetland habitat can suffer severe damage (Hothem et al., 1988; Otis and Kilburn, 1988). One area of possible high productivity for blackbirds, especially redwings, is roadside wetland habitat. Roadside rights-of-way provide a unique management option because there is no need to contact multiple landowners for permission to manage wetland vegetation. Prior to manipulation of roadside habitat, data on both blackbird and non-blackbird avian use of this habitat is necessary in order to develop proper management strategies. Thus, the major objective of this study was to determine avian use of roadside habitat. It is also the goal of this study to assess the relationships between bird species, number, and habitat variables.

Materials and Methods

Study Area. The study area consisted of thirty randomly selected quarter sections (1 quarter section \approx 64.75 ha) containing at least 200 linear meters of cattail. These quarter sections were located in the Southern Drift Plains of North Dakota (Figure 1). Counties in this region containing surveyed quarter sections were Stutsman, Ransom, Barnes, and Sargent. This area was selected because previous studies have shown a high density of breeding red-winged blackbirds in this region (Nelms et al., 1994; Wimberly et al., 2002). As a result of glaciation, this area of the Prairie Pothole Region contains a high density of wetlands (Euliss et al., 1999). Land in the region is used primarily for agriculture, including high yields of sunflower, which are damaged by blackbirds. Surveys of these quarter sections were conducted during June 2001 and 2002.




Figure 1. Map of the Southern Drift Plains of North Dakota.

Avian Use Surveys. Roadside habitat along two 0.5 mile (\approx 0.8 km) transects bordering these quarter sections was surveyed in order to assess avian use. Avian use surveys were initiated at sunrise and were terminated four hours thereafter. The species, number, and location of birds within these transects were recorded. In addition to morning avian use surveys, nest surveys were conducted to provide an index of breeding bird use of this roadside habitat.

Habitat Assessment. Roadside grass density was surveyed using a visual obstruction method (Camp and Best, 1994); cattail density was determined using a stem count method. Ancillary data were collected on cattail stand length, water depth, and inter-cattail stand distances. Habitats adjacent to the roadside were also categorized.

Statistical Analysis. Abundance and nest data were tested for normality using normal probability plots; homogeneity of variances were tested using a Hartley's test. Logarithmic transformations were performed on nest data to correct for deviations from normality and homogeneity of variance. Paired *t*-tests were conducted on abundance and nest data between blackbirds and non-blackbirds. Bird abundance was further analyzed using analysis of variance and Duncan's multiple comparison tests. All statistical tests were conducted using an alpha level of 0.05.

Results and Discussion

We found 2,529 birds representing 49 different species during the two study years. In 2001, 1,206 birds representing 38 species were observed, with 805 representing blackbird species (red-winged blackbirds, yellow-headed blackbirds, and common grackles), while the remaining 401 birds represented non-blackbirds. In 2002, we saw an increase in the number of birds, though blackbird numbers decreased; 1,323 birds representing 42 species were observed, with 674 representing blackbird species, while the remaining 649 birds represented non-blackbirds. In 2001, we observed significantly more blackbirds than non-blackbirds per transect. The numbers of observed blackbirds and non-blackbirds per transect was not significantly different in 2002 (Figure 2). We also compared our numbers of individual blackbird species with non-blackbird numbers (Figure 3). In 2001, we located 496 nests, and 181 of these nests, represented by 10 species, contained eggs and young and were designated as active. Of the active nests, 164 were blackbird nests, and 158 of these nests were from red-winged blackbirds. In 2002, we located 329 nests; 126 of these nests were active. In 2002, active nests were represented by 11 species. Of the active nests, 109 were blackbird nests, and 87 of these nests were from red-winged blackbirds. In 2001, we found significantly more blackbird nests than non-blackbird nests per transect. In 2002, we did not find a significant difference in blackbird versus non-blackbird nest numbers per transect (Figure 4). In the second year of our study, cattail management was observed on 15 of our 30 quarter sections. Management practices included burning, mowing, and plowing. These practices resulted in fewer cattail stands observed as well as decreased cattail density within cattail stands.




Figure 2. Comparison of mean blackbird and non-blackbird avian use numbers.




Figure 3. Comparison of mean avian use numbers using a Duncan's multiple comparison test. COGR=common grackle, RWBL=red-winged blackbird, and YHBL=yellow headed blackbird. Species with different letters are significant at $\alpha=0.5$.




Figure 4. Mean number of blackbird and non-blackbird nests occurring in roadside habitat.

Conclusions

Based on avian use and nesting data, blackbirds, especially red-winged blackbirds, are the dominant bird species using roadside cattail. With proper management of roadside habitat, potential effects on non-blackbirds can be minimized, and nesting habitat for blackbirds can be reduced. Though preliminary, our data indicate that a loss in nesting habitat will likely result in

fewer blackbirds and subsequently might reduce sunflower damage. Also, management of cattail habitat will likely result in an increase in the abundance of non-blackbirds, especially waterfowl.

Acknowledgments

We thank Carina Lee and Alegra Galle for assistance with data collection in 2001 and 2002, respectively. Greg Forcey provided suggestions regarding data analysis. We also thank Gary Clambey, Don Kirby, and Gary Nuechterlein for their insight and suggestions with our study. This research was funded by the National Wildlife Research Center, a unit within the Wildlife Services program of the United States Department of Agriculture, Animal and Plant Health Inspection Service.

References

- Camp M, and L.B. Best. 1994. Nest density and nesting success of birds in roadsides adjacent to rowcrop fields. *Am. Midl. Nat.* 131:347-358.
- Euliss, N.H., Jr., D.M. Mushet, and D.A. Wrubleski. 1999. Wetlands of the Prairie Pothole Region: Invertebrate Species Composition, Ecology, and Management. p. 471-514. *In:* D. P. Batzer, R. B. Rader and S. A. Wissinger, eds. *Invertebrates in Freshwater Wetlands of North America: Ecology and Management*, Chapter 21. John Wiley & Sons, New York. Jamestown, ND: Northern Prairie Wildlife Research Center Home Page. <http://www.npwrc.usgs.gov/resource/1999/pothole/pothole.htm> (Version 02SEP99).
- Hothem, R.L., R.W. DeHaven, and S.D. Fairaizl. 1988. Bird damage to sunflower in North Dakota, South Dakota, and Minnesota, 1979-1981. *U.S. Fish Wildl. Serv. Tech. Rep.* 15. 11pp.
- Linz, G.M., D.C. Blixt, D.L. Bergman, and W.J. Bleier. 1994. Response of black terns (*Chlidonias niger*) to glyphosate-induced habitat alterations on wetlands. *Colonial Waterbirds.* 17(2):160-167.
- Linz, G.M., D.C. Blixt, D.L. Bergman, and W.J. Bleier. 1996. Responses of red-winged blackbirds, yellow-headed blackbirds and marsh wrens to glyphosate-induced alterations in cattail density. *Journal of Field Ornithology.* 67(1):167-176.
- Nelms, C.O., W.J. Bleier, D.L. Otis, and G.M. Linz. 1994. Population estimates of breeding blackbirds in North Dakota, 1967, 1981-1982 and 1990. *Am. Midl. Nat.* 132:256-263.
- Otis, D.L., and C.M. Kilburn. 1988. Influence of environmental factors on blackbird damage to sunflower. *U.S. Fish and Wildl. Tech. Rep.* 16. 11 pp.
- Solberg, K.L., and K. Higgins. 1992. Chemical treatment of monodominant cattail stands in semipermanent wetlands: duck, invertebrate and vegetation response. *Cattail Management Symposium*. U.S. Dept. of Agriculture, Animal and Plant Health Inspection Service, Animal Damage Control, Denver Wildlife Research Center, U.S. Dept. of Interior, Fish and Wildlife Service. Jamestown, ND: Northern Prairie Wildlife Research Center Home Page. <http://www.npwrc.usgs.gov/resource/othrdata/cattail/cattail.htm> (Version 15AUG97).
- Wimberly, R.L., G.M. Linz, W.J. Bleier, and H.J. Homan. 2002. Landscape effects on breeding blackbird abundance and sunflower damage in the Southern Drift Plains of North Dakota. pp 152-154. *In:* Proc. 24th Sunflower Research Workshop, Fargo, ND. Natl. Sunflower Assoc., Bismarck, ND. USA.

Proceedings of the
**16th International
Sunflower Conference**



Fargo, North Dakota, USA

August 29-September 2, 2004

Sponsored by

The International Sunflower Association, Paris, France

In cooperation with

**The National Sunflower Association, Bismarck, North Dakota,
USA**

Vol. I