

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Roman L. Hruska U.S. Meat Animal Research
Center

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2004

Characterization of biological types of cattle (Cycle VI): Carcass, yield, and longissimus palatability traits

T. L. Wheeler

U.S. Meat Animal Research Center, tommy.wheeler@ars.usda.gov

L. V. Cundiff

U.S. Meat Animal Research Center, Larry.Cundiff@ars.usda.gov

S. D. Shackelford

ARS, USDA, Roman L. Hruska U.S. Meat Animal Research Center

M. Koohmaraie

ARS, USDA, Roman L. Hruska U.S. Meat Animal Research Center

Follow this and additional works at: <http://digitalcommons.unl.edu/hruskareports>

Wheeler, T. L.; Cundiff, L. V.; Shackelford, S. D.; and Koohmaraie, M., "Characterization of biological types of cattle (Cycle VI): Carcass, yield, and longissimus palatability traits" (2004). *Roman L. Hruska U.S. Meat Animal Research Center*. 369.
<http://digitalcommons.unl.edu/hruskareports/369>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Roman L. Hruska U.S. Meat Animal Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Characterization of biological types of cattle (Cycle VI): Carcass, yield, and longissimus palatability traits^{1,2}

T. L. Wheeler³, L. V. Cundiff, S. D. Shackelford, and M. Koohmaraie

ARS, USDA, Roman L. Hruska U.S. Meat Animal Research Center, Clay Center, NE 68933-0166

ABSTRACT: Carcass (n = 568) and longissimus thoracis palatability (n = 460) traits from F₁ steers obtained from mating Hereford (H), Angus (A), and U.S. Meat Animal Research Center (MARC) III cows to H, A, Norwegian Red (NR), Swedish Red and White (RW), Friesian (F), or Wagyu (W) sires were compared. Data were adjusted to constant age (471 d), carcass weight (356 kg), fat thickness (1.0 cm), percentage of fat trim (24%), and marbling (Small³⁵) end points. For Warner-Bratzler shear force and trained sensory panel traits, data were obtained on longissimus thoracis steaks stored at 2°C for 14 d postmortem. The following comparisons were from the age-constant end point. Carcasses from H- and A-sired steers (377 and 374 kg, respectively) were the heaviest ($P < 0.05$) and carcasses from W-sired steers (334 kg) were the lightest ($P < 0.05$). A greater ($P < 0.05$) percentage of carcasses from A- and W-sired steers graded USDA Choice (88 and 85%, respectively) than carcasses from other sire breeds (52 to 71%). Adjusted fat thickness for carcasses from A-sired steers (1.3 cm) was highest ($P < 0.05$), followed by H-sired steers (1.1 cm) and W- and F-sired steers (0.9 cm); NR- and RW-sired steers (0.8 cm) had the

lowest ($P < 0.05$) adjusted fat thickness. Longissimus thoracis area was not different ($P > 0.05$) among sire breeds (mean = 80.6 cm²). Carcass yield of boneless, totally trimmed retail product was least ($P < 0.05$) for A-sired steers (60.1%), intermediate for H-sired steers (61.5%), and similar ($P > 0.05$) for all other sire breeds (62.5 to 62.8%). Longissimus thoracis steaks from carcasses of A- (3.7 kg) and W-sired (3.7 kg) steers had lower ($P < 0.05$) shear force values than longissimus thoracis steaks from other sire breeds (4.1 to 4.2 kg). Trained sensory panel tenderness, juiciness, or beef flavor intensity ratings for longissimus thoracis steaks did not differ ($P > 0.05$) among the sire breeds. Sire breed comparisons were affected by adjusting data to other end points. Heritability estimates for various carcass, yield, and palatability traits ranged from very low ($h^2 = 0.06$ for percentage of kidney, pelvic, and heart fat) to relatively high ($h^2 = 0.71$ for percentage of retail product yield). Relative to the other sire breeds, W-sired steers had the highest percentage of USDA Choice, Yield grade 1 and 2 carcasses, but their carcasses were the lightest.

Key Words: Beef, Breeds, Carcass, Palatability, Quality, Tenderness

©2004 American Society of Animal Science. All rights reserved.

J. Anim. Sci. 2004. 82:1177–1189

Introduction

The first five cycles of the Germplasm Evaluation (GPE) program at the Roman L. Hruska U.S. Meat Animal Research Center (MARC) characterized 22

breeds representing several biological types of cattle. Carcass and longissimus thoracis (LM) palatability traits from these studies have been reported by Koch et al. (1976, 1979, 1982b) and Wheeler et al. (1996, 2001). Breed differences in production traits are important genetic resources for improving beef production efficiency, and meat composition and quality. No one breed excels in all traits that are important to beef production. Diverse breeds are required to exploit heterosis and complementarity through crossbreeding, and to match genetic potential with diverse markets, feed resources, and climates. Evaluation of carcass traits and meat palatability from different breeds, or breed crosses, is important in determining the potential value of alternative germplasm resources for profitable beef production. This paper reports on Cycle VI (which includes two Scandinavian breeds and the Japanese Wagyu) of the GPE program that characterizes cattle

¹Names are necessary to report factually on available data; however, the USDA neither guarantees nor warrants the standard of the product, and the use of the name by USDA implies no approval of the product to the exclusion of others that may also be suitable.

²The authors wish to acknowledge the technical assistance of the cattle operations staff, P. Beska, R. Harris, D. Kohmetscher, D. Light, K. Mihm, K. Ostdiek, P. Tammen, and J. Waechter, and the secretarial assistance of M. Bierman.

³Correspondence: P.O. Box 166 (phone: 402-762-4229; fax: 402-762-4149; e-mail: wheeler@email.marc.usda.gov).

Received August 8, 2003.

Accepted November 13, 2003.

breeds representing diverse biological types for carcass and LM palatability traits that affect the quantity, quality, and value of production.

Materials and Methods

Animals

Hereford, Angus, and MARC III ($\frac{1}{4}$ Angus, $\frac{1}{4}$ Hereford, $\frac{1}{4}$ Pinzgauer, and $\frac{1}{4}$ Red Poll) dams of mature age (4 to 11 yr) were mated by AI to 30 Angus, 32 Hereford, 14 Norwegian Red, 16 Swedish Red and White, 24 Friesian, and 19 Wagyu bulls to produce 568 steer calves. To avoid confounding sire breed effects with heterosis effects, no purebred Hereford or purebred Angus matings were made. Semen from 11 of the Hereford and 10 of the Angus sires (born 1982 to 1984) was used for the first time in Cycle IV. Semen from 10 of the Hereford and 9 of the Angus sires (born in 1989 or 1990) was used for the first time in Cycle V. Semen from 11 of the Hereford and 11 of the Angus sires (born since 1994) was used for the first time in Cycle VI. At the time of semen purchase, the Hereford and Angus bulls used were unproven by progeny test, but were considered to be excellent young herd sire prospects in the Angus and Hereford breeds. In cooperation with seed-stock breeders and commercial AI organizations, young sires (2 to 3 yr old) identified as herd sire prospects, based on EPD for growth, were selected to represent the Hereford, and Angus breeds. For comparison to previous GPE cycles, the simple mean of the Hereford and Angus sire least squares means has been included in the tables labeled H \times , A \times .

Semen from Norwegian Red bulls was imported from the Norwegian Cattle AI and Breeding Cooperative (Hamar, Norway), whereas semen from Swedish Red and White bulls was imported from the Svensk Avel Ornsro (Skara, Sweden). The Norwegian Red and the Swedish Red and White breeds could be considered the same breed because their herd books have been open to one another; however, we have presented them separately. Over the past 20 yr, each breed has introduced germplasm from North American Holsteins; however, only bulls with non-Holstein influenced pedigrees were sampled for this experiment.

Semen from Friesian bulls was obtained through the American Beef Friesian breeders. Fourteen of the bulls were full bloods originally imported into the United States from Ireland, England, The Netherlands, and Germany. Ten of the bulls were upgrades or full-blood sons of imported parents born in the United States from 1979 to 1990. Only bulls with non-Holstein-influenced pedigrees were sampled for this experiment.

Semen from Wagyu bulls was obtained through the American Wagyu Association. Four bulls were full bloods born in 1973 and 1974 and imported into the United States in the late 1970s. Eleven bulls were full bloods born in 1989 to 1994 and imported from Japan into the United States in the 1990s. Four of the bulls

were upgraded purebreds ($\geq 15/16$ Wagyu) born in 1985 to 1989 in the United States.

Calves were born in the spring, beginning in March each year (1997 and 1998). Male calves were castrated within 24 h of birth. Calves were creep fed whole oats from mid-July or early August until weaning in early October. Calves averaged approximately 216 d of age at weaning. For 21 d following weaning, a diet comprising about 2.55 Mcal of ME/kg and 14.25% CP was fed. Following this postweaning adjustment period, steers were assigned to replicated pens within sire breed and fed separately by sire breed for an average of 255 d (range from 225 to 293 d). A growing diet (2.7 Mcal of ME/kg of DM and 11.8% CP) containing 66% corn silage, 22% corn, and 12% supplement (DM basis) was fed until steers weighed approximately 320 kg. A finishing diet (3.05 Mcal of ME/kg of DM and 13.1% CP) containing 25% corn silage, 70% corn, and 5% supplement (DM basis) was fed from approximately 320 kg to slaughter. Steers were implanted with Synovex S (200 mg of progesterone and 20 mg of estradiol benzoate) in mid-December and again in early March of each year. Steers were slaughtered serially each year, in five slaughter groups spaced about 14 d apart (spanning 56 d) in 1998 and in four slaughter groups spaced 14 to 21 d apart (spanning 56 d) in 1999.

Final unshrunk live weights were obtained 1 wk before slaughter. Steers were slaughtered in a commercial beef-processing facility. Carcass sides were electrically stimulated within 45 min postmortem with the following sequence: 68 (3 s on, 3 s off), 70 (2 s on, 3 s off), 70 (2 s on, 3 s off), and 70 V (2 s on, 3 s off). However, observation of the electrical stimulation equipment while the experimental carcasses were being processed indicated that the system was working sporadically and sometimes ineffectively, particularly in the second year. Carcasses were spray-chilled with a mist of 2°C water for 30 s every 5 min during the first 12 h of chilling. After a 36-h chill at 0°C, USDA yield and quality grade data were obtained by trained MARC personnel (USDA, 1997).

Samples

The wholesale rib (#103; NAMP, 1997) from the right side of each carcass was returned to the meat laboratory at MARC. At 3 d postmortem, the wholesale rib was dissected into the ribeye roll (NAMP #112), lean trim, fat trim, and short ribs for prediction of retail product yield as described by Shackelford et al. (1995). The ribeye roll was vacuum-packaged and stored at 2°C. In the first year, ribeye rolls were frozen at 14 d postmortem and steaks were cut on a band saw. In the second year, steaks were cut fresh at 7 d postmortem with a Biro slicer (model 109 PC; Marblehead, OH), vacuum packaged, stored at 2°C until 14 d postmortem, and then frozen at -30°C. The posterior end of the ribeye roll was squared off by removing a wedge-shaped slice that was trimmed of all fat, epimysium, and non-LM

muscles, and then frozen immediately at -30°C for later proximate analysis of the raw LM. Then, four 2.54-cm-thick LM steaks were cut from the posterior end of the ribeye roll. The first steak was not used in this experiment. The second and third steaks were used for trained sensory panel evaluation. The fourth steak was used for determination of Warner-Bratzler shear force and for proximate composition of the cooked LM. Steaks were stored frozen for 3 to 5 mo before thawing for evaluation.

Warner-Bratzler Shear Force

Frozen steaks were thawed at 5°C for 24 h and then cooked on a conveyerized belt grill to a final internal temperature of 71°C , as described by Wheeler et al. (1998). Warner-Bratzler shear force was determined as described by Wheeler et al. (1998).

Trained Sensory Evaluation

A subsample of 460 LM steaks (approximately 80% of each sire breed) was evaluated by a trained descriptive attribute sensory panel. Immediately after cooking, the LM was cut into $1\text{ cm} \times 1\text{ cm} \times$ steak thickness cubes. Three cubes were served warm to each panel member. An eight-member sensory panel, trained according to procedures described by Cross et al. (1978), evaluated cooked steaks for tenderness, juiciness, and beef flavor intensity on an eight-point scale (8 = extremely tender, juicy, or intense to 1 = extremely tough, dry, or bland). A warm-up sample was served first, and then four experimental steaks were served in each of two sessions (5 min between sessions) per day, three days per week. In addition, a duplicate sample to one of the experimental samples was served daily for monitoring panelist and panel performance.

Proximate Composition Analyses

Raw and cooked LM proximate composition (wet-weight basis) was determined according to AOAC (1985) methods as described by Wheeler et al. (2001).

Statistical Analyses

Data were analyzed by least squares, mixed model procedures (Harvey, 1985) using a model that included a random effect for sires nested within sire breed, and fixed effects for sire breed, dam breed, age of dam (4 to 5, 6 to 7, 8 to 9, and ≥ 10 yr), birth year, interaction of sire breed \times dam breed, and covariates for age at weaning (mean = 216 d) and days fed postweaning (mean = 255 d). Sire nested within sire breed was used to test sire breed and residual variance was used to test other fixed effects. Estimates of heritability and genetic and phenotypic correlations were derived following procedures outlined by Harvey (1985).

Regressions of traits on days fed provides a method of adjusting the age-constant sire breed means to alter-

native end points. The regressions were used for estimating values that would have been obtained if all animals in a sire breed had been fed fewer or more days until the breed group average reached a given end point (the mean for this experiment) with regard to age (471 d), carcass weight (356 kg), fat thickness (1.0 cm), fat trim percentage (24%; for cuts trimmed to 0 cm of fat cover), or marbling (Small³⁵) following procedures used in previous cycles of GPE (Koch et al., 1979, 1982b; Wheeler et al., 1996, 2001).

Consistent with previous reports (Koch et al., 1979, 1982b; Wheeler et al., 1996, 2001), the average regression over all sire breeds was modified by a proportionate adjustment of the sire breed mean to the general mean as described by Wheeler et al. (1996). Sire breeds were compared using the average LSD for $\alpha = 0.05$ computed for all possible pairwise contrasts using the sire within sire breed mean square as the error term in the linear contrast procedure of Harvey (1985).

Results and Discussion

The ANOVA indicated that sire breed, dam breed, and year were significant ($P < 0.05$) sources of variation for most traits (Table 2). The sire breed \times dam breed interaction was a significant ($P < 0.05$) source of variation for marbling score only. Linear regression of weaning age was significant ($P < 0.05$) for live and carcass weights and linear regression of days fed was significant ($P < 0.05$) for most traits.

Carcass Traits

Sire breeds differed ($P < 0.05$) in growth rate. Final live and carcass weights at a constant age of 471 d were heaviest ($P < 0.05$) for Hereford- and Angus-sired steers and lightest ($P < 0.05$) for Wagyu-sired steers (Table 3). At a constant fat thickness, Swedish Red and White- and Norwegian Red-sired steers were heaviest ($P < 0.05$), then Friesian- and Hereford-sired steers, and Wagyu- and Angus-sired steers were the lightest ($P < 0.05$). At a constant marbling degree, Hereford-sired steers were heaviest ($P < 0.05$), followed by H \times , A \times crosses, Swedish Red and White-, and Friesian-sired steers, then Angus- and Norwegian Red-sired steers, and Wagyu-sired steers were the lightest ($P < 0.05$). At a constant fat trim end point, Norwegian Red-sired steers were heaviest ($P < 0.05$), followed by Hereford-, Friesian-, and Swedish Red and White-sired steers, and then Angus-sired steers; Wagyu-sired steers were lightest ($P < 0.05$). Sire breed differences for live weight and carcass weight were similar ($P > 0.05$).

Dressing percentage was higher ($P < 0.05$) at 471 d of age, and tended to be higher at the other end points for carcasses from Wagyu-, Hereford-, and Angus-sired steers. Dressing percentage for carcasses from Friesian-, Swedish Red and White- and Norwegian Red-sired steers was lower ($P < 0.05$) at 471 d of age and tended to be lower at other end points.

Table 1. Number of sires used per breed and number of steers in each sire breed × dam breed subclass

Sire breed	No. of sires	Dam breed and number of steer progeny			
		Hereford	Angus	MARC III ^a	Total
Hereford	32	—	65	21	86
Angus	30	29	—	59	88
Norwegian Red	14	11	34	18	63
Swedish Red and White	16	10	27	37	74
Friesian	24	25	49	58	132
Wagyu	19	24	56	45	125
Total	145	99	231	238	568

^aComposite consisting of ¼ each Hereford, Angus, Pinzgauer, and Red Poll.

Adjusted fat thickness was highest ($P < 0.05$) for carcasses from Hereford- and Angus-sired steers and lowest ($P < 0.05$) for carcasses from Swedish Red and White- and Norwegian Red-sired steers at constant age. At constant weight, carcasses from Wagyu- and Angus-sired steers had the highest ($P < 0.05$) and carcasses from Norwegian Red- and Swedish Red and White-sired steers had the lowest ($P < 0.05$) adjusted fat thickness. At constant marbling, carcasses from Hereford-sired steers had the highest ($P < 0.05$) adjusted fat thickness followed by carcasses from Friesian-, Angus-, and Swedish Red and White-sired steers. Carcasses from Norwegian Red- and Wagyu-sired steers had the lowest ($P < 0.05$) adjusted fat thickness at constant marbling. At constant fat trim percentage, sire breed differences in adjusted fat thickness were greatly reduced with car-

casses from Angus- and Hereford-sired steers tending to have the greatest and carcasses from Swedish Red and White-sired steers tending to have the least adjusted fat thickness.

Sire breed differences in LM area were relatively small. Carcasses from Hereford-sired steers had larger ($P < 0.05$) LM areas than did carcasses from Swedish Red and White- and Norwegian Red-sired steers at all end points except constant marbling, where Hereford was different ($P < 0.05$) only from Swedish Red and White. Carcasses from Hereford- and Angus-sired steers had similar ($P > 0.05$) LM areas at all endpoints and had similar ($P < 0.05$) LM areas as carcasses from Wagyu-sired steers at all end points except constant weight. It should be noted that the days on feed regression coefficient for LM area has a sign opposite (–) from

Table 2. Analysis of variance

Source	df ^a	Mean squares						
		Live weight, kg	Hot carcass weight, kg	Dressing percent	Adjusted fat thickness, cm	Longissimus area, cm	Marbling score	USDA Choice, %
Sire breed (SB)	5	291,101*	109,182*	11.2*	0.56*	2.0	61,238*	1.93*
Sire (Sire breed)	126	11,100*	4,201*	2.5	0.03*	1.5*	6,967*	0.24*
Dam breed (DB)	2	39,092*	23,311*	9.8*	0.26*	0.2	32,600*	0.92*
Dam age	3	44,072*	18,083*	2.1	0.03	1.7	3,822	0.08
Year (Y)	1	186,462*	78,175*	1.3	0.13*	0.2	47,045*	1.42*
SB × DB	8	9,368	2,660	1.3	0.01	0.8	9,646*	0.26
b1 (weaning age)	1	56,444*	21,331*	0.1	0.01	3.0	8,350	0.01
b2 (days fed)	1	617,850*	252,670*	2.5	1.04*	1.2	115,764*	2.08*
Residual	420	7,917	3,341	2.2	0.02	1.1	4,680	0.16

Source	Mean squares						
	USDA yield grade	Boneless retail product yield, %	df ^b	Warner-Bratzler shear force, kg	Tenderness rating	Juiciness rating	Beef flavor intensity rating
Sire breed (SB)	6.9*	88.2*	5	2.54*	1.45*	0.21	0.06
Sire (Sire breed)	0.4*	12.4*	119	0.48*	0.56	0.13	0.09*
Dam breed (DB)	4.0*	23.0*	2	2.27*	2.11*	0.00	0.05
Dam age	0.3	6.6	3	0.15	0.65	0.00	0.05
Year (Y)	2.0*	122.7*	1	13.47*	66.21*	15.82*	2.47*
SB × DB	0.3	8.9	8	0.11	0.43	0.23	0.08
b1 (weaning age)	0.1	24.0	1	0.03	0.25	0.03	0.02
b2 (days fed)	27.9*	483.2*	1	1.50*	0.93	0.47*	0.20
Residual	0.3	6.6	319	0.39	0.46	0.10	0.07

^aDegrees of freedom for carcass traits.

^bDegrees of freedom for palatability traits.

* $P < 0.05$.

Table 3. Least squares means for carcass traits adjusted to a common age, carcass weight, fat thickness, marbling, or fat trim percent^a

Trait, $\mu \pm$ SEM, b1, b2 ^b	Sire breed, LSD ^c	Endpoint				
		Age (471 d)	Carcass wt (356 kg)	Fat thickness (1.0 cm)	Marbling (Small ³⁵)	Fat trim (24%)
Days on feed $\mu = 255$ SD = 24	Hereford	—	216	232	292	237
	Angus	—	226	214	200	213
	H \times , A \times	—	221	223	246	225
	Norwegian Red	—	258	301	245	285
	Swedish Red and White	—	267	305	279	272
	Friesian	—	269	284	284	277
	Wagyu	—	304	272	224	265
Live weight, kg $\mu = 585 \pm 2.7$ b1 = 0.8037 \pm 0.301 b2 = 0.8223 \pm 0.093	Hereford	618	—	599	648	603
	Angus	609	—	575	564	574
	H \times , A \times	612	—	587	606	589
	Norwegian Red	588	—	626	580	613
	Swedish Red and White	580	—	622	600	594
	Friesian	576	—	599	599	594
	Wagyu	543	—	557	517	551
LSD	15	—	19	18	17	
Hot carcass weight, kg $\mu = 358 \pm 1.5$ b1 = 0.4781 \pm 0.198 b2 = 0.5215 \pm 0.061	Hereford	380	—	368	399	370
	Angus	374	—	352	345	352
	H \times , A \times	377	—	360	372	361
	Norwegian Red	357	—	381	351	372
	Swedish Red & White	352	—	378	365	361
	Friesian	351	—	366	366	362
	Wagyu	334	—	343	318	339
LSD	9	—	12	11	10	
Dressing percent $\mu = 61.1 \pm 0.09$ b1 = -0.0017 \pm 0.011 b2 = 0.0037 \pm 0.0034	Hereford	61.4	61.3	61.4	61.5	61.4
	Angus	61.4	61.3	61.3	61.3	61.3
	H \times , A \times	61.4	61.3	61.3	61.4	61.3
	Norwegian Red	60.6	60.6	60.7	60.6	60.7
	Swedish Red & White	60.6	60.6	60.7	60.7	60.7
	Friesian	60.9	61.0	61.0	61.0	61.0
	Wagyu	61.5	61.7	61.6	61.4	61.5
LSD	0.5	0.6	0.7	0.6	0.6	
Adj. fat thickness, cm $\mu = 0.96 \pm 0.024$ b1 = 0.0018 \pm 0.0024 b2 = 0.0060 \pm 0.0008	Hereford	1.16	0.93	—	1.38	1.06
	Angus	1.33	1.16	—	1.01	1.08
	H \times , A \times	1.25	1.05	—	1.19	1.07
	Norwegian Red	0.78	0.80	—	0.72	0.96
	Swedish Red & White	0.76	0.83	—	0.91	0.86
	Friesian	0.85	0.93	—	1.02	0.98
	Wagyu	0.91	1.20	—	0.72	0.97
LSD	0.13	0.16	—	0.16	0.15	
Longissimus area, cm ² $\mu = 80.8 \pm 0.4$ b1 = 0.0830 \pm 0.0495 b2 = -0.0165 \pm 0.0153	Hereford	82.7	83.6	83.3	81.9	83.1
	Angus	81.2	81.8	82.1	82.4	82.1
	H \times , A \times	81.9	82.7	82.7	82.1	82.6
	Norwegian Red	80.0	79.9	78.9	80.2	79.3
	Swedish Red & White	79.2	79.0	78.1	78.7	78.9
	Friesian	80.3	79.9	79.6	79.6	79.8
	Wagyu	80.8	79.7	80.4	81.5	80.6
LSD	2.5	3.0	3.2	3.0	2.9	
KPH fat, % ^d $\mu = 2.55 \pm 0.03$ b1 = -0.0011 \pm 0.0036 b2 = 0.0053 \pm 0.0011	Hereford	2.29	2.08	2.17	2.49	2.20
	Angus	2.67	2.51	2.45	2.37	2.44
	H \times , A \times	2.48	2.30	2.31	2.43	2.32
	Norwegian Red	2.59	2.61	2.84	2.54	2.75
	Swedish Red and White	2.58	2.64	2.85	2.71	2.67
	Friesian	2.57	2.64	2.72	2.72	2.68
	Wagyu	2.59	2.85	2.68	2.42	2.64
LSD	0.16	0.19	0.21	0.19	0.19	

(Continued)

Table 3 continued. Least squares means for carcass traits adjusted to a common age, carcass weight, fat thickness, marbling, or fat trim percent^a

Trait, $\mu \pm$ SEM, b1, b2 ^b	Sire breed, LSD ^c	Endpoint				
		Age (471 d)	Carcass wt (356 kg)	Fat thickness (1.0 cm)	Marbling (Small ³⁵)	Fat trim (24%)
Yield grade	Hereford	3.2	2.7	2.9	3.6	3.0
$\mu = 2.95 \pm 0.04$	Angus	3.5	3.1	2.9	2.8	2.9
b1 = 0.0023 \pm 0.0038	H \times , A \times	3.3	2.9	2.9	3.2	2.9
b2 = 0.0122 \pm 0.0012	Norwegian Red	2.8	2.8	3.4	2.7	3.2
	Swedish Red & White	2.8	2.9	3.4	3.1	3.0
	Friesian	2.8	3.0	3.2	3.2	3.1
	Wagyu	2.7	3.3	2.9	2.3	2.8
	LSD	0.2	0.2	0.3	0.2	0.2
Marbling ^e	Hereford	509	480	492	—	495
$\mu = 536.8 \pm 4.7$	Angus	579	557	548	—	547
b1 = 0.681 \pm 0.510	H \times , A \times	544	518	520	—	521
b2 = 0.785 \pm 0.158	Norwegian Red	543	545	577	—	565
	Swedish Red & White	518	527	555	—	530
	Friesian	514	525	536	—	531
	Wagyu	559	595	572	—	566
	LSD	25	30	33	—	29
USDA Choice, % ^f	Hereford	60	47	53	—	54
$\mu = 68.9 \pm 0.03$	Angus	89	79	75	—	75
b1 = 0.00065 \pm 0.003	H \times , A \times	74	63	64	—	65
b2 = 0.00332 \pm 0.001	Norwegian Red	71	72	86	—	81
	Swedish Red & White	61	65	77	—	66
	Friesian	52	57	62	—	59
	Wagyu	85	100	91	—	89
	LSD	15	18	20	—	18

^aEndpoints represent the overall mean for that trait in this experiment.

^bb1 = regression coefficient for weaning age; b2 = regression coefficient for days on feed.

^cLSD = least significant difference among means, $P < 0.05$.

^dEstimated percentage of hot carcass weight as kidney, pelvic, and heart fat.

^e400 = Slight⁰⁰, 500 = Small⁰⁰ (USDA, 1997).

^fPercentage of carcasses grading USDA Choice or higher.

that expected; thus, the results of adjustments to different end points should be interpreted with caution. This likely resulted from random chance that more animals with larger LM areas were assigned to earlier harvest groups than would be expected.

Carcasses from Hereford-sired steers had the lowest ($P < 0.05$) percentage of kidney, pelvic, and heart (KPH) fat at all end points except constant marbling. At constant weight, the percentage of KPH fat was greatest ($P < 0.05$) in carcasses from Wagyu-sired steers. Carcasses from Norwegian Red-, Swedish Red and White-, and Friesian-sired steers had similar ($P > 0.05$) percentages of KPH fat.

At constant age, numerical USDA yield grade was higher ($P < 0.05$) for carcasses from Angus- and Hereford-sired steers than for all other sire breeds. At constant age, the mean yield grade of 3.5 for Angus-sired steers resulted in a relatively high percentage (20.5%) of carcasses with a yield grade 4.0 or greater. Hereford-, Norwegian Red-, Swedish Red and White-, Wagyu-, and Friesian-sired steers at constant age had 10.0, 3.2, 2.7, 2.3, and 1.5% of carcasses with a yield grade 4.0, or higher, respectively. At constant weight, carcasses from Wagyu-sired steers had higher ($P < 0.05$) yield grades than all other sire breeds except Angus. At constant

fat thickness, Norwegian Red- and Swedish Red and White-sired steers had higher ($P < 0.05$) yield grades than all other sire breeds except Friesian. At constant marbling, carcasses from Hereford-sired steers had the highest ($P < 0.05$) yield grades, followed by carcasses from Friesian- and Swedish Red and White-sired steers, and then Norwegian Red- and Angus-sired steers; Wagyu-sired steers had the lowest ($P < 0.05$) yield grades. At constant fat thickness, differences in yield grades among sire breeds were small.

At constant age, marbling score was higher ($P < 0.05$) in carcasses from Angus-sired steers than for all other sire breeds except Wagyu. At constant age, carcasses from Hereford-, Friesian-, and Swedish Red and White-sired steers had lower ($P < 0.05$) marbling scores than did all other sire breeds. At constant weight, carcasses from Wagyu-sired steers had the highest ($P < 0.05$) and carcasses from Hereford-sired steers had the lowest ($P < 0.05$) marbling scores. At constant fat thickness and constant fat trim level, carcasses from Norwegian Red- and Wagyu-sired steers tended to have the highest marbling scores, whereas carcasses from Hereford-sired steers tended to have the lowest. Sire breed differences for the percentage of carcasses grading USDA Choice at each end point were similar to marbling differences.

Only one carcass in the experiment graded USDA Standard.

Hereford- and Angus-sired steers were the heaviest ($P < 0.05$) at constant age and thus were the fastest growing sire breeds. Wagyu-sired steers had the slowest growth rate. Angus-sired steers were the earliest maturing, followed by Hereford-sired steers. The Angus-sired steers required the fewest days on feed to reach the 24% fat trim end point.

Most of the published data comparing the Wagyu breed to other breeds is based on a number of observations too small for valid conclusions. In a comparison of 12 calf-fed and 12 2-yr-old cattle in each of two breed crosses, Wertz et al. (2002) found that $\frac{1}{2}$ Wagyu \times $\frac{1}{2}$ Angus tended to have more marbling than $\frac{3}{4}$ blood Angus, and that the 2-yr-old Wagyu crosses had smaller carcass weights and lower fat thickness. However, the calf-fed Wagyu crosses had larger carcass weights and greater fat thickness than Angus crosses. Mir et al. (1999) reported that 50 and 75% Wagyu crosses had lighter carcasses with more marbling and tended to have decreased fat thickness compared with Continental crosses. Lunt et al. (1993) compared 10 Angus and 10 American Wagyu fed according to Japanese custom for 552 d, and found no differences in any carcass traits except that Angus had heavier carcasses. Mears et al. (2001) reported marbling levels were not different between Hereford or Angus compared with Wagyu crosses with Angus or Hereford. They also found conflicting results across years for differences in fat thickness. In an experiment that was more comprehensive than most involving Wagyu, Pitchford et al. (2002) reported that carcasses from Angus-sired progeny were heavier than carcasses from Hereford-sired progeny, which were heavier than carcasses from Wagyu-sired progeny. Pitchford et al. (2002) also found that LM from carcasses of Wagyu- and Angus-sired cattle had a greater percentage of i.m. fat than LM from carcasses of Hereford-sired cattle.

A comparison of cattle selected for twinning that were 16% Swedish Friesian, 7% Norwegian Red, and 2% Swedish Red and White to a reference population that was $\frac{1}{2}$ Simmental crosses indicated that twinner cattle had higher marbling scores, decreased adjusted fat thickness, and increased percentage grading USDA Choice or higher (Gregory et al., 1996). Kempster et al. (1982) compared breeds at 16 mo of age after slaughter at a constant fat level, and reported that Friesian and Hereford steers had heavier carcass weights than Angus steers (but lighter than Charolais and Simmental), and that Friesian had greater LM areas than Angus or Hereford. A serial slaughter at different fat levels resulted in similar findings (Kempster et al., 1988).

Carcass Yield

At a constant age of 471 d, carcasses from Norwegian Red-, Swedish Red and White-, and Friesian-sired steers tended to have the highest and carcasses from

Angus-sired steers had the lowest ($P < 0.05$) percentage retail product yield (Table 4). At constant weight, carcasses from Hereford-sired steers had a higher ($P < 0.05$) percentage of retail product yield than all other sire breeds except Norwegian Red, and carcasses from Wagyu-sired steers had the lowest ($P < 0.05$) percentage of retail product yield. At constant fat thickness, carcasses from Hereford- and Angus-sired steers tended to have the higher percentage of retail product yield. At constant marbling, carcasses from Wagyu- and Norwegian Red-sired steers tended to have the highest and carcasses from Hereford-sired steers had the lowest ($P < 0.05$) percentage of retail product yield. There were no differences ($P > 0.05$) among sire breeds for percentage of retail product yield adjusted to constant percentage of fat trim.

At constant age, carcasses from Angus-sired steers had the highest ($P < 0.05$) percentage of fat trim yield, and carcasses from Norwegian Red-, Friesian-, Swedish Red and White-, and Wagyu-sired steers had the lowest ($P < 0.05$) percentage fat yield. At constant weight, carcasses from Wagyu- and Angus-sired steers tended to have the highest and carcasses from Norwegian Red- and Hereford-sired steers tended to have the lowest percentage fat yield. At constant fat thickness, sire breed differences in fat yield were small, but carcasses from Swedish Red and White-sired steers tended to have the highest and carcasses from Hereford-sired steers tended to have the lowest percentage fat yield. At constant marbling, carcasses from Hereford-sired steers had the highest ($P < 0.05$) and carcasses from Wagyu-sired steers had the lowest ($P < 0.05$) percentage of fat yield.

Carcasses from Norwegian Red- and Friesian-sired steers tended to have the highest percentage of bone, regardless of end point. Carcasses from Angus-sired steers tended to have the lowest percentage of bone, regardless of end point.

Carcasses from Hereford-sired steers had the heaviest ($P < 0.05$) and carcasses from Wagyu-sired steers had the lightest ($P < 0.05$) weight of retail product for most end points. At constant age, carcasses from Hereford- and Angus-sired steers had heavier ($P < 0.05$) fat weight than other sire breeds. At constant weight, carcasses from Wagyu-sired steers had heavier ($P < 0.05$) fat weight than all other sire breeds except Angus. At constant fat thickness, carcasses from Swedish Red and White- and Norwegian Red-sired steers had heavier ($P < 0.05$) fat weight than carcasses from Angus- and Wagyu-sired steers. At constant marbling, carcasses from Hereford-sired steers had the heaviest ($P < 0.05$) fat weight, followed by Friesian- and Swedish Red and White-sired steers, and then Angus- and Norwegian Red-sired steers; Wagyu-sired steers had the lightest ($P < 0.05$) fat weight. At constant fat trim, carcasses from Hereford- and Norwegian Red-sired steers had heavier ($P < 0.05$) fat weights than carcasses from Wagyu-sired steers. Regardless of end point, carcasses from Hereford- and Norwegian Red-sired steers

Table 4. Least squares means for carcass yield traits adjusted to a common age, carcass weight, fat thickness, fat trim percent, or marbling endpoint^a

Trait, $\mu \pm$ SEM, b1, b2 ^b	Sire breed, LSD ^c	Endpoint				
		Age (471 d)	Carcass wt (356 kg)	Fat thickness (1.0 cm)	Marbling (Small ³⁵)	Fat trim (24%)
Retail product yield, % ^d $\mu = 62.1 \pm 0.21$ b1 = -0.0368 \pm 0.0193 b2 = -0.0515 \pm 0.0060	Hereford	61.5	63.6	62.7	59.6	62.5
	Angus	60.1	61.6	62.2	62.9	62.3
	H \times , A \times	60.8	62.6	62.5	61.3	62.4
	Norwegian Red	62.8	62.6	60.4	63.3	61.2
	Swedish Red and White	62.8	62.2	60.2	61.6	62.0
	Friesian	62.8	62.1	61.3	61.3	61.7
	Wagyu	62.5	59.9	61.6	64.1	62.0
	LSD	1.1	1.3	1.4	1.3	1.3
Fat yield, % ^d $\mu = 23.9 \pm 0.25$ b1 = 0.0405 \pm 0.0236 b2 = 0.0506 \pm 0.0074	Hereford	24.9	22.9	23.8	26.8	—
	Angus	26.4	24.9	24.3	23.6	—
	H \times , A \times	25.7	23.9	24.0	25.2	—
	Norwegian Red	22.5	22.7	24.9	22.0	—
	Swedish Red and White	23.1	23.8	25.7	24.4	—
	Friesian	22.9	23.6	24.4	24.4	—
	Wagyu	23.5	26.0	24.4	21.9	—
	LSD	1.3	1.6	1.6	1.6	—
Bone yield, % ^d $\mu = 14.6 \pm 0.06$ b1 = -0.0060 \pm 0.0068 b2 = -0.0046 \pm 0.0021	Hereford	14.5	14.6	14.6	14.3	14.5
	Angus	14.0	14.1	14.2	14.2	14.2
	H \times , A \times	14.2	14.4	14.4	14.3	14.4
	Norwegian Red	15.0	15.0	14.8	15.0	14.8
	Swedish Red and White	14.7	14.7	14.5	14.6	14.6
	Friesian	14.9	14.8	14.8	14.8	14.8
	Wagyu	14.4	14.2	14.4	14.6	14.4
	LSD	0.3	0.4	0.6	0.4	0.4
Retail product wt, kg ^e $\mu = 221 \pm 1.2$ b1 = 0.1646 \pm 0.1214 b2 = 0.1425 \pm 0.0379	Hereford	233	228	230	238	231
	Angus	224	220	219	217	218
	H \times , A \times	229	224	224	228	225
	Norwegian Red	224	224	230	222	228
	Swedish Red and White	221	222	228	224	223
	Friesian	220	223	224	224	223
	Wagyu	208	215	211	204	210
	LSD	6	8	8	8	7
Fat weight, kg ^e $\mu = 86 \pm 1.0$ b1 = 0.2527 \pm 0.1114 b2 = 0.3046 \pm 0.0348	Hereford	95	83	88	107	90
	Angus	99	90	86	82	86
	H \times , A \times	97	87	87	94	88
	Norwegian Red	81	82	95	78	90
	Swedish Red and White	82	86	97	89	87
	Friesian	81	85	90	90	88
	Wagyu	79	94	84	70	82
	LSD	6	7	7	7	7
Bone weight, kg ^e $\mu = 52 \pm 0.3$ b1 = 0.0489 \pm 0.0309 b2 = 0.0598 \pm 0.0096	Hereford	55	52	53	57	54
	Angus	52	51	50	49	50
	H \times , A \times	53	51	52	53	52
	Norwegian Red	53	54	56	53	55
	Swedish Red and White	52	52	55	53	53
	Friesian	52	53	54	54	53
	Wagyu	48	51	49	46	49
	LSD	2	2	2	2	2

^aEnd points represent the overall mean for that trait in this experiment.

^bb1 = regression coefficient for weaning age; b2 = regression coefficient for days on feed. W-B = Warner-Bratzler.

^cLSD = least significant difference among means, $P < 0.05$.

^dPredicted from wholesale rib dissection.

^eCalculated from hot carcass weight and predicted yields.

tended to have heavier bone weight, and carcasses from Wagyu-sired steers tended to have the lightest bone weights.

Kempster et al. (1982) compared breeds at 16 mo of age after slaughter at a constant fat level, and reported

that carcasses from Friesian steers had a lower saleable meat yield than Angus, Charolais, or Simmental steers. Similar results were obtained when cattle were serially slaughtered at different fat levels (Kempster et al., 1988).

Table 5. Least squares means for longissimus thoracis steak palatability traits adjusted to a common age, carcass weight, fat thickness, fat trim percent, or marbling endpoint^a

Trait, $\mu \pm \text{SEM}$, b1, b2 ^b	Sire breed, LSD ^c	Endpoint				
		Age (471 d)	Carcass wt (356 kg)	Fat thickness (1.0 cm)	Marbling (Small ³⁵)	Fat trim (24%)
W-B shear force, kg $\mu = 3.75 \pm 0.04$ b1 = 0.0011 \pm 0.0047 b2 = -0.0027 \pm 0.0015	Hereford	3.79	3.90	3.85	3.69	3.84
	Angus	3.55	3.63	3.67	3.70	3.67
	H \times , A \times	3.67	3.76	3.76	3.70	3.75
	Norwegian Red	3.77	3.76	3.65	3.80	3.69
	Swedish Red and White	3.93	3.90	3.80	3.87	3.89
	Friesian	3.94	3.90	3.86	3.86	3.88
	Wagyu	3.53	3.40	3.49	3.62	3.51
	LSD	0.21	0.25	0.27	0.25	0.25
Tenderness ^d $\mu = 6.40 \pm 0.04$ b1 = -0.0043 \pm 0.0058 b2 = 0.0026 \pm 0.0018	Hereford	6.25	6.15	6.19	6.35	6.20
	Angus	6.48	6.40	6.37	6.33	6.37
	H \times , A \times	6.36	6.27	6.28	6.34	6.28
	Norwegian Red	6.48	6.49	6.60	6.45	6.56
	Swedish Red and White	6.32	6.35	6.45	6.39	6.37
	Friesian	6.25	6.29	6.33	6.33	6.31
	Wagyu	6.60	6.73	6.64	6.52	6.63
	LSD	0.26	0.31	0.34	0.31	0.30
Juiciness ^e $\mu = 5.31 \pm 0.02$ b1 = 0.0015 \pm 0.0028 b2 = 0.0018 \pm 0.0009	Hereford	5.29	5.21	5.24	5.35	5.25
	Angus	5.29	5.24	5.21	5.19	5.21
	H \times , A \times	5.29	5.22	5.23	5.27	5.23
	Norwegian Red	5.34	5.34	5.42	5.32	5.39
	Swedish Red and White	5.32	5.34	5.41	5.36	5.35
	Friesian	5.24	5.27	5.30	5.30	5.28
	Wagyu	5.39	5.48	5.42	5.33	5.41
	LSD	0.12	0.15	0.16	0.15	0.14
Beef flavor intensity ^f $\mu = 4.71 \pm 0.02$ b1 = -0.0012 \pm 0.0022 b2 = 0.0012 \pm 0.0007	Hereford	4.70	4.65	4.67	4.74	4.68
	Angus	4.73	4.69	4.68	4.66	4.67
	H \times , A \times	4.71	4.67	4.67	4.70	4.68
	Norwegian Red	4.71	4.71	4.76	4.70	4.75
	Swedish Red and White	4.70	4.72	4.76	4.73	4.72
	Friesian	4.68	4.70	4.71	4.71	4.71
	Wagyu	4.76	4.81	4.76	4.72	4.77
	LSD	0.10	0.13	0.14	0.13	0.12

^aEndpoints represent the overall mean for that trait in this experiment.

^bb1 = regression coefficient for weaning age; b2 = regression coefficient for days on feed. WB = Warner-Bratzler.

^cLSD = least significant difference among means, $P < 0.05$.

^d1 = extremely tough, 4 = slightly tough, 5 = slightly tender, 8 = extremely tender.

^e1 = extremely dry, 4 = slightly dry, 5 = slightly juicy, 8 = extremely juicy.

^f1 = extremely bland, 4 = slightly bland, 5 = slightly intense, 8 = extremely intense.

Palatability Traits

Differences among sire breeds for LM palatability traits were generally small. At constant age, LM steaks from carcasses of Wagyu- and Angus-sired steers had the lowest ($P < 0.05$) 14-d postmortem Warner-Bratzler shear force values, followed by Norwegian Red- and Hereford-sired steers; Friesian- and Swedish Red and White-sired steers had the highest ($P < 0.05$) 14-d shear force (Table 5). At constant weight, shear force differences were similar to those for constant age. At constant fat thickness and constant fat trim percentage, LM steaks from carcasses of Wagyu-, Norwegian Red-, and Angus-sired steers tended to have lower shear force than other sire breeds. At constant marbling, 14-d LM Warner-Bratzler shear force values did not vary ($P > 0.05$) among sire breeds.

At constant age and constant weight, LM steaks of carcasses from Wagyu-sired steers had higher ($P < 0.05$)

trained sensory panel tenderness ratings than did other sire breeds except Norwegian Red (constant age and weight) and Angus (constant age). At constant fat thickness, LM steaks from Wagyu-sired steers had higher ($P < 0.05$) tenderness ratings than LM steaks from Hereford-sired steers. At constant percentage of fat trim, LM steaks from Wagyu-sired steers had higher ($P < 0.05$) tenderness ratings than did longissimus from Hereford- and Friesian-sired steers. At constant marbling, 14-d LM steak tenderness ratings did not vary ($P > 0.05$) among sire breeds.

Longissimus thoracis steaks from Wagyu-sired steers were juicier ($P < 0.05$) than those from Friesian-sired steers at constant age (Table 5). At constant weight, LM steaks from Wagyu-sired steers had higher ($P < 0.05$) juiciness ratings than those from all other sire breeds except Norwegian Red and Swedish Red and White. At constant fat thickness, LM steaks from Wagyu-, Norwegian Red-, and Swedish Red and White-

Table 6. Effect of sire breed on least squares means for chemical composition (wet-weight basis) of raw and cooked longissimus thoracis at 471 d of age

Sire breed	Raw			Cooked		
	Lipid, %	Moisture, %	Protein, % ^a	Lipid, %	Moisture, %	Protein, % ^a
$\mu \pm$ SEM	5.4 \pm 0.09	71.8 \pm 0.08	22.8 \pm 0.03	6.5 \pm 0.10	64.2 \pm 0.10	29.3 \pm 0.07
Hereford	4.9	72.4	22.7	6.0	64.6	29.4
Angus	6.2	71.1	22.7	7.4	63.5	29.1
H \times , A \times	5.5	71.8	22.7	6.7	64.1	29.2
Norwegian Red	5.5	71.6	22.9	6.5	64.2	29.3
Swedish Red and White	5.1	71.8	23.1	6.2	64.3	29.5
Friesian	4.8	72.3	22.9	5.9	64.5	29.6
Wagyu	5.8	71.4	22.8	7.0	63.7	29.3
LSD ^b	0.5	0.5	0.4	0.6	0.5	0.4

^aCalculated by difference.

^bLSD = least significant difference among means, $P < 0.05$.

sired steers were juicier ($P < 0.05$) than LM steaks from Hereford- and Angus-sired steers. At constant marbling, LM steaks from Swedish Red and White- and Hereford-sired steers had higher ($P < 0.05$) juiciness ratings than LM steaks from Angus-sired steers. At constant fat trim percentage, LM steaks from Wagyu-sired steers had higher ($P < 0.05$) juiciness ratings than LM steaks from Hereford- and Angus-sired steers, whereas LM steaks from Norwegian Red-sired steers was juicier ($P < 0.05$) than LM steaks from Angus-sired steers.

There were no differences ($P > 0.05$) detected among sire breeds in LM beef flavor intensity ratings at any end point except constant carcass weight (Table 5). At constant weight, LM steaks from Wagyu-sired steers had more ($P < 0.05$) intense beef flavor than did LM steaks from Hereford-sired steers; however, the magnitude of the difference indicated it was of little practical importance.

With only three or four observations per breed type, Busboom et al. (1993) reported that after feeding for 524 d, loin steaks from Japanese Wagyu received higher tenderness ratings from a trained sensory panel than did steaks from Angus or Longhorn, but received tenderness ratings similar to those for American Wagyu and USDA Choice steaks. No differences were detected in other sensory traits. Consumer evaluation of loin steaks found Japanese Wagyu to be more tender than Angus or USDA Choice, but not different from American Wagyu (Busboom et al., 1993). May et al. (1993) compared 10 Angus and 10 Wagyu crosses fed a Japanese diet for 552 d, and reported no difference in any trained sensory panel traits; however, consumer triangle tests detected differences between the breeds that were attributed to tenderness, flavor, juiciness, fatty taste, or some combination of these. Mir et al. (1999) reported that shear force values were not different among LM steaks from Continental crosses or 50 or 75% Wagyu steers.

Results from previous cycles of GPE (Koch et al., 1976, 1979, 1982b; Wheeler et al., 1996, 2001) have indicated a similar mean LM steak tenderness among

most breeds. Perhaps more important than breed averages is to consider that after 14 d postmortem, the range in breed mean differences was about equal to the range in breeding value within breed, indicating that among-breed variation in LM steak tenderness is about the same as variation within breeds (Wheeler et al., 1996, 2001).

Longissimus Thoracis Chemical Composition

Chemical composition of raw LM adjusted to 471 d of age indicated that LM from carcasses of Angus- and Wagyu-sired steers had the highest ($P < 0.05$) percentages of lipid, followed by Norwegian Red- and Swedish Red and White-sired steers, and that the LM from Friesian- and Hereford-sired steers tended to have the lowest percentages of lipid (Table 6). Differences among sire breeds in the percentage of cooked LM lipid were similar to those for raw LM. Sire breeds differences in LM moisture percentage were inversely related to lipid differences. Raw and cooked LM from carcasses of Angus- and Wagyu-sired steers tended to have the lowest percentages of moisture. No sire breed differences ($P > 0.05$) in protein content were detected in raw LM. Differences among sire breeds for percentage moisture in the cooked LM were statistically significant ($P < 0.05$) but were small in magnitude and appeared to be of little practical importance.

Lunt et al. (1993) reported raw LM from American Wagyu was higher in ether extract (18.9%) than was the LM from Angus (14.5%). Mir et al. (1999) reported that across 2 yr, inconsistent differences in LM fat percentage were detected among Continental crosses and 50 and 75% Wagyu steers and heifers.

Heritabilities and Correlation Coefficients

The range of differences among sire breed means (R) from topcross progeny estimates half of the breed differences (Table 7). Thus, R was doubled to assess purebred genetic variation relative to within-sire breed genetic (σ_g) and phenotypic (σ_p) variation. However, phenotypic

Table 7. Variation among sire breeds for carcass and palatability traits at 471 d of age

Trait	R ^a	h ² ± SE ^b	σ _g ^c	2R/σ _g	σ _p ^d	R/σ _p
Live weight, kg	76	0.31 ± 0.16	23.34	6.51	42.03	1.88
Hot carcass weight, kg	46	0.17 ± 0.15	11.10	8.29	26.88	1.71
Dressing percent	0.9	0.15 ± 0.15	0.57	3.16	1.51	0.60
Adjusted fat thickness, cm	0.57	0.54 ± 0.17	0.26	4.38	0.35	1.63
Longissimus area, cm ²	3.5	0.37 ± 0.16	4.26	1.64	6.99	0.50
Kidney, pelvic, heart fat, %	0.38	0.06 ± 0.14	0.13	5.85	0.49	0.78
Yield grade	0.8	0.49 ± 0.17	0.38	4.21	0.54	1.48
Marbling	65	0.35 ± 0.16	42.30	3.07	71.08	0.91
Retail product yield, %	2.3	0.71 ± 0.17	2.38	1.93	2.83	0.81
Retail product weight, kg	25	0.49 ± 0.17	11.98	4.17	17.21	1.45
Raw longissimus lipid, %	1.4	0.27 ± 0.16	0.75	3.73	1.43	0.98
Cooked longissimus lipid, %	1.5	0.32 ± 0.16	0.95	3.16	1.67	0.90
Warner-Bratzler shear force, kg	0.41	0.16 ± 0.15	0.25	3.28	0.63	0.65
Tenderness	0.35	0.25 ± 0.19	0.35	2.00	0.70	0.50
Juiciness	0.15	0.24 ± 0.18	0.16	1.88	0.33	0.45
Beef flavor intensity	0.08	0.40 ± 0.19	0.17	0.94	0.27	0.30

^aR = Range in sire breed means.

^bh² = Heritability.

^cσ_g = Genetic standard deviation.

^dσ_p = Phenotypic standard deviation.

variation was expressed without doubling R, thus representing F₁ progeny phenotypic variation. Heritability estimates for various carcass, yield, and palatability traits ranged from very low (h² = 0.06 for KPH percentage) to relatively high (h² = 0.71; percentage of retail product yield). Heritabilities of carcass traits ranged from low to high and were similar to those reported by Wheeler et al. (1996) and Koch et al. (1982a), but lower for many carcass traits than reported by Wheeler et al. (2001). Heritabilities of marbling and measures of LM chemical lipid were moderate and similar to one another. Tenderness, as measured by Warner-Bratzler shear force and trained sensory tenderness rating, had low to moderate heritability estimates.

These values are consistent with the average of heritabilities reported in the literature (reviewed by Koch et al., 1982a). Some estimates of the heritability of tenderness (or shear force) have been higher (h² = 0.53, Shackelford et al., 1994; h² = 0.50, Wheeler et al., 1996) and others lower (h² = 0.12, Gregory et al., 1994). Heritability estimates for juiciness and beef flavor intensity ratings were higher than those reported previously (Wheeler et al., 1996, 2001).

Estimates of the amount of variation between the two extreme breeds for a given trait in standard deviation units (2R/σ_g) from the present experiment were about the same or lower for most traits compared to values reported by Wheeler et al. (1996, 2001). Hot carcass weight had more variation among breeds than within breeds. Live weight and percentage of KPH fat had about the same amount of variation within as among breeds. All other traits had more variation within breeds than among breeds. These results are consistent with previous data indicating there is as much, or more, variation in LM steak tenderness within breeds as among the most extreme breeds for that trait. Phenotypic variation in carcass and palatability traits

was similar to, or slightly less than, that reported by Wheeler et al. (1996, 2001).

As was observed in Cycles I to V of GPE, little inherent genetic variation in juiciness and beef flavor intensity was detected in Cycle VI. Phenotypic variation in tenderness rating was about twice that of variation in ratings of juiciness and beef flavor intensity (CV 14.8, 8.3, and 6.4%, respectively). This occurred despite a wide range in marbling scores. Thus, when variation in juiciness and beef flavor intensity occurs at the consumer level, it may be mostly induced by cooking practices and the level and kind of flavor enhancers added.

The genetic correlation between fat thickness and marbling was moderately high, suggesting that it would be difficult, but not impossible, to decrease s.c. fat thickness without lowering marbling level (Table 8). Marbling had relatively high genetic correlations to all carcass traits, but not to all palatability traits. Tenderness traits had high genetic correlations to several carcass traits, but not to marbling. Retail product yield had high genetic correlations to most fatness traits. Juiciness rating had high genetic correlations to raw LM lipid percentage, but not marbling or cooked LM lipid percentage. Shear force had moderate to low genetic correlations to carcass and LM lipid traits and high genetic correlations to live and carcass weight. Tenderness rating was more strongly related to most carcass traits than was shear force. Beef flavor intensity rating had moderate to high genetic correlations to weight, fatness, and palatability traits.

Phenotypic correlations were not as high as genetic correlations (Table 8). Moderate phenotypic correlations were detected between hot carcass weight and fat thickness, LM area, and yield grade. Phenotypically, marbling was strongly related to measures of LM lipid and retail product yield. Yield grade had high phenotypic correlations to fat thickness and retail product

Table 8. Genetic and phenotypic correlation coefficients among carcass and palatability traits at 471 d of age^a

Trait	Trait						
	LWT	HCWT	AFT	LA	YG	MARB	RPY
Live weight, LWT		0.95	0.27	0.44	0.30	0.06	-0.26
Hot carcass weight, HCWT	0.99 ± 0.06		0.31	0.51	0.30	0.05	-0.26
Adj. fat thickness, AFT	0.08 ± 0.30	0.05 ± 0.40		-0.04	0.81	0.28	-0.65
Longissimus area, LA	0.12 ± 0.35	0.37 ± 0.38	-0.45 ± 0.30		-0.46	-0.09	0.20
USDA yield grade, YG	0.07 ± 0.31	-0.12 ± 0.40	0.90 ± 0.07	-0.83 ± 0.46		0.28	-0.69
Marbling score, MARB	-0.45 ± 0.36	-0.98 ± 0.59	0.53 ± 0.25	-0.82 ± 0.40	0.58 ± 0.25		-0.49
Retail product yield, RPY	0.11 ± 0.29	0.33 ± 0.43	-0.86 ± 0.37	0.34 ± 0.24	-0.66 ± 0.36	-0.77 ± 0.40	
Raw lipid, RLIPID ^b	-0.66 ± 0.40	1.00 ^c ± 0.65	0.27 ± 0.30	-0.40 ± 0.38	0.14 ± 0.32	0.98 ± 0.17	-0.51 ± 0.40
Cooked lipid, CLIPID ^b	-0.12 ± 0.37	-0.40 ± 0.49	0.54 ± 0.26	-0.12 ± 0.35	0.34 ± 0.28	0.81 ± 0.19	-0.81 ± 0.43
Warner-Bratzler shear force, WBS	0.85 ± 0.64	1.00 ^c ± 1.00	0.04 ± 0.41	0.57 ± 0.51	0.15 ± 0.43	-0.03 ± 0.49	0.03 ± 0.37
Tenderness, TEND	1.00 ^c ± 1.00	NE ^d	-0.63 ± 0.50	-0.49 ± 0.54	-0.61 ± 0.49	-0.08 ± 0.42	0.30 ± 0.34
Juiciness, JUICY	0.17 ± 0.93	NE ^d	0.28 ± 0.42	-0.16 ± 0.49	0.19 ± 0.42	0.21 ± 0.41	-0.41 ± 0.40
Beef flavor intensity, FLAV	-0.73 ± 0.98	NE ^d	0.29 ± 0.33	-0.43 ± 0.39	0.35 ± 0.34	0.15 ± 0.34	-0.36 ± 0.30

Trait	Trait					
	RLIPID	CLIPID	WBS	TEND	JUICY	FLAV
Live weight, LWT	0.11	0.06	-0.01	-0.14	-0.09	-0.02
Hot carcass weight, HCWT	0.11	0.06	-0.01	-0.14	-0.09	-0.02
Adj. fat thickness, AFT	0.27	0.26	0.00	-0.11	0.04	0.03
Longissimus area, LA	0.00	-0.07	0.05	-0.13	-0.16	-0.02
USDA yield grade, YG	0.23	0.26	-0.03	-0.05	0.08	0.03
Marbling score, MARB	0.64	0.59	-0.15	0.11	0.23	0.14
Retail product yield, RPY	-0.40	-0.46	0.01	0.08	-0.12	-0.09
Raw lipid, RLIPID ^b		0.57	-0.12	0.10	0.22	0.17
Cooked lipid, CLIPID ^b	1.00 ^c ± 0.22		-0.17	0.10	0.21	0.18
Warner-Bratzler shear force, WBS	-0.04 ± 0.55	0.07 ± 0.52		-0.62	-0.26	-0.10
Tenderness, TEND	-0.23 ± 0.50	-0.67 ± 0.51	-1.00 ^c ± 1.00		0.21	0.08
Juiciness, JUICY	0.92 ± 0.50	0.13 ± 0.49	-0.90 ± 1.00	0.68 ± 0.47		0.38
Beef flavor intensity, FLAV	0.41 ± 0.39	0.53 ± 0.38	-0.69 ± 1.00	0.09 ± 0.45	0.77 ± 0.36	

^aGenetic correlation coefficients and their standard errors are below the diagonal; phenotypic correlation coefficients are above the diagonal.

^bChemical analysis of the longissimus thoracis (wet-weight basis).

^cEstimate exceeded 1.00 and thus was set at 1.00.

^dNE = not estimable.

yield. Phenotypically, retail product yield was more highly correlated to fat thickness and yield grade than was marbling score. Raw LM lipid percentage had a slightly higher phenotypic correlation to marbling score than to cooked LM lipid percentage. Phenotypically, shear force and tenderness rating were strongly correlated only to each other, although measures of tenderness were not strongly related to beef flavor intensity and juiciness ratings.

Implications

Differences exist among and within cattle sire breeds in carcass and meat palatability traits. Selection of sire breed and end point of production are critical in order for producers to successfully target carcass and longissimus characteristics. Among the sire breeds evaluated in this cycle of the Germplasm Evaluation Project, Wagyu-sired steers had the highest percentage of USDA Choice, Yield grade 1 and 2 carcasses, were among the highest in longissimus thoracis steak tenderness, had greater retail product yield than the British sire breeds, but were the smallest of all the sire breeds. The Scandinavian sire breeds had greater retail product yields,

similar longissimus thoracis steak tenderness, a higher percentage of USDA Choice, Yield grade 1 and 2 carcasses, but lighter carcasses than the British sire breeds. These results provide producers with more information when deciding which sire breeds will maximize profit potential in their production situation.

Literature Cited

- AOAC. 1985. Official Methods of Analysis. 14th ed. Assoc. Offic. Anal. Chem., Washington, DC.
- Busboom, J. R., L. E. Jeremiah, L. L. Gibson, K. A. Johnson, C. T. Gaskins, J. J. Reeves, and R. W. Wright. 1993. Effects of biological source on cooking and palatability attributes of beef produced for the Japanese market. *Meat Sci.* 35:241–258.
- Cross, H. R., R. Moen, and M. S. Stanfield. 1978. Training and testing of judges for sensory analysis of meat quality. *Food Technol.* 32:48–54.
- Gregory, K. E., L. V. Cundiff, R. M. Koch, M. E. Dikeman, and M. Koohmaraie. 1994. Breed effects, retained heterosis, and estimates of genetic and phenotypic parameters for carcass and meat traits of beef cattle. *J. Anim. Sci.* 72:1174–1183.
- Gregory, K. E., S. E. Echterkamp, and L. V. Cundiff. 1996. Effects of twinning on Dystocia, calf survival, calf growth, carcass traits, and cow productivity. *J. Anim. Sci.* 74:1223–1233.
- Harvey, W. R. 1985. User's guide for LSML76. (Mimeo) The Ohio State Univ., Columbus.

- Kempster, A. J., G. L. Cook, and J. R. Southgate. 1982. A comparison of the progeny of British Friesian dams and different sire breeds in 16- and 24-month beef production systems. *Anim. Prod.* 34:167–178.
- Kempster, A. J., G. L. Cook, and J. R. Southgate. 1988. Evaluation of British Friesian, Canadian Holstein and beef breed \times British Friesian steers slaughtered over a commercial range of fatness from 16-month and 24-month beef production systems. *Anim. Prod.* 46:365–378.
- Koch, R. M., M. E. Dikeman, D. M. Allen, M. May, J. D. Crouse, and D. R. Campion. 1976. Characterization of biological types of cattle III. Carcass composition, quality and palatability. *J. Anim. Sci.* 43:48–62.
- Koch, R. M., M. E. Dikeman, R. J. Lipsey, D. M. Allen, and J. D. Crouse. 1979. Characterization of biological types of cattle—cycle II: III. Carcass composition, quality and palatability. *J. Anim. Sci.* 49:448–460.
- Koch, R. M., L. V. Cundiff, and K. E. Gregory. 1982a. Heritabilities and genetic, environmental and phenotypic correlations of carcass traits in a population of diverse biological types and their implications in selection programs. *J. Anim. Sci.* 55:1319–1329.
- Koch, R. M., M. E. Dikeman, and J. D. Crouse. 1982b. Characterization of biological types of cattle (cycle III). III. Carcass composition, quality and palatability. *J. Anim. Sci.* 54:35–45.
- Lunt, D. K., R. R. Riley, and S. B. Smith. 1993. Growth and carcass characteristics of Angus and American Wagyu steers. *Meat Sci.* 34:327–334.
- May, S. G., C. A. Sturdivant, D. K. Lunt, R. K. Miller, and S. B. Smith. 1993. Comparison of sensory characteristics and fatty acid composition between Wagyu crossbred and Angus steers. *Meat Sci.* 35:289–298.
- Mears, G. J., P. S. Mir, D. R. C. Bailey, and S. D. M. Jones. 2001. Effect of Wagyu genetics on marbling, backfat and circulating hormones in cattle. *Can. J. Anim. Sci.* 81:65–73.
- Mir, P. S., D. R. C. Bailey, Z. Mir, T. Entz, S. D. M. Jones, W. M. Robertson, R. J. Wseselake, and F. J. Lozeman. 1999. Growth, carcass, and meat quality characteristics of beef cattle with 0, 50 and 75 percent Wagyu genetic influence. *Can. J. Anim. Sci.* 79:129–137.
- NAMP. 1997. *The Meat Buyers Guide*. N. Am. Meat Processors Assoc., Reston, VA.
- Pitchford, W. S., M. P. B. Deland, B. D. Siebert, A. E. O. Malau-Aduli, and C. D. K. Bottema. 2002. Genetic variation in fatness and fatty acid composition of crossbred cattle. *J. Anim. Sci.* 80:2825–2832.
- Shackelford, S. D., L. V. Cundiff, K. E. Gregory, and M. Koohmariae. 1995. Predicting beef carcass cutability. *J. Anim. Sci.* 73:406–413.
- Shackelford, S. D., M. Koohmariae, L. V. Cundiff, K. E. Gregory, G. A. Rohrer, and J. W. Savell. 1994. Heritabilities and phenotypic and genetic correlations for bovine postrigor calpastatin activity, intramuscular fat content, Warner-Bratzler shear force, retail product yield, and growth rate. *J. Anim. Sci.* 72:857–863.
- USDA. 1997. *Official United States standards for grades of carcass beef*. Agric. Marketing Serv., USDA, Washington, DC.
- Wertz, A. E., L. L. Berger, P. M. Walker, D. B. Faulkner, F. K. McKeith, and S. L. Rodriguez-Zas. 2002. Early-weaning and postweaning nutritional management affect feedlot performance, carcass merit, and the relationship of 12th-rib fat, marbling score, and feed efficiency among Angus and Wagyu heifers. *J. Anim. Sci.* 80:28–37.
- Wheeler, T. L., L. V. Cundiff, R. M. Koch, and J. D. Crouse. 1996. Characterization of biological types of cattle (Cycle IV): Carcass traits and longissimus palatability. *J. Anim. Sci.* 74:1023–1035.
- Wheeler, T. L., L. V. Cundiff, S. D. Shackelford, and M. Koohmariae. 2001. Characterization of biological types of cattle (Cycle V): Carcass traits and longissimus palatability. *J. Anim. Sci.* 79:1209–1222.
- Wheeler, T. L., S. D. Shackelford, and M. Koohmariae. 1998. Cooking and palatability traits of beef longissimus steaks cooked with a belt grill or an open hearth electric broiler. *J. Anim. Sci.* 76:2805–2810.