

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Papers and Publications in Animal  
Science

Animal Science Department

---

January 1978

## Estimation of Genetic Trends from Cow and Sire Evaluations

R. L. Hintz  
*Cornell University*

R. W. Everett  
*Cornell University*

L. Dale Van Vleck  
*University of Nebraska-Lincoln*, [dvan-vleck1@unl.edu](mailto:dvan-vleck1@unl.edu)

Follow this and additional works at: <https://digitalcommons.unl.edu/animalscifacpub>



Part of the [Animal Sciences Commons](#)

---

Hintz, R. L.; Everett, R. W.; and Van Vleck, L. Dale, "Estimation of Genetic Trends from Cow and Sire Evaluations" (1978). *Faculty Papers and Publications in Animal Science*. 360.  
<https://digitalcommons.unl.edu/animalscifacpub/360>

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Papers and Publications in Animal Science by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## OUR INDUSTRY TODAY

### Estimation of Genetic Trends from Cow and Sire Evaluations

R. L. HINTZ<sup>1</sup>, R. W. EVERETT, and L. D. VAN VLECK

Department of Animal Science  
Cornell University  
Ithaca, NY 14853

#### ABSTRACT

Milk production records of Ayrshire, Guernsey, Holstein, Jersey, and Brown Swiss in the Northeast were used to estimate trends in transmitting ability of sires and genetic merit of cows. Genetic trends in the transmitting ability of Artificial Insemination sires for the Ayrshire, Guernsey, Holstein, Jersey, and Brown Swiss breeds averaged 23.7, 14.6, 17.9, 18.3, and 34.7 kg per year.

Genetic trends in the AI cow populations averaged 36.1, 25.4, 26.1, 25.0 and 38.1 kg. The genetic trends of cow populations were less than twice the contribution of sires to genetic trends, indicating that estimating genetic trends in cow populations by doubling the trend in transmitting ability of sires is biased upwards.

#### INTRODUCTION

A goal of dairy cattle breeders is to increase genetic merit for milk production. To determine the effectiveness of breeding programs, genetic trends in dairy cattle populations must be monitored. The purpose of this study was to investigate genetic trends in milk production among Holstein, Ayrshire, Guernsey, Jersey, and Brown Swiss cows and sires in the Northeast.

#### MATERIALS AND METHODS

Records of Ayrshire, Guernsey, Holstein, Jersey, and Brown Swiss cows which were on Dairy Herd Improvement test in 1975 in the Northeast were used. Only cows calving first between the ages of 18 and 35 mo and between 1961 and 1974 were included. All available

lactation records of a cow were used to estimate genetic merit of cows and sires. The cows were divided into two groups: artificially-sired cows (AI) and naturally-sired cows (non-AI). The AI cows were cows with an identified AI sire, but non-AI cows were cows either with an identified natural-service sire or cows without an identified sire. Numbers of cows per year of freshening are summarized in Tables 1 and 2.

Unbiased predictors of the additive genetic value of cows and sires were obtained by best linear unbiased prediction (BLUP) as described by Henderson (10) and Slinger et al. (21). The model was

$$y = X\beta + Za + Zp + e$$

where  $y$  is the vector of age-adjusted milk records;  $\beta$  is a vector of fixed herd-year-season effects and the population mean;  $X$  is a known design matrix;  $a$  is a vector of additive genetic values of the cows and sires;  $Z$  is a known matrix relating elements of  $y$  to elements of  $a$ ;  $p$  is a vector with each element being the remainder of the total genetic value plus the permanent environmental effect specific to a particular cow; and  $e$  is a vector of temporary environmental effects. All elements of  $a$ ,  $p$ , and  $e$  have zero expected values;  $\text{Var}(a) = Ab^2\sigma_y^2$  where  $A$  is the numerator relationship matrix of the animals in the herd,  $b^2$  is the heritability, and  $\sigma_y^2$  is the within herd-year-season variance of records in a noninbred population;  $\text{Var}(p) = I(r-b^2)\sigma_y^2$  where  $r$  is repeatability;  $\text{Var}(e) = I(l-r)\sigma_y^2$ . Further,  $a$ ,  $p$ , and  $e$  are uncorrelated. The covariance between records on different animals is due to additive genetic effects; thus,  $Ab^2\sigma_y^2$  is the matrix of covariances between the cows and sires used in the herd.

The predictions of additive genetic value or genetic merits were used to estimate genetic trends. For estimating genetic trends, a cow's estimated genetic merit was used only in the

Received June 3, 1977.

<sup>1</sup>Department of Animal Science, Oklahoma State University, Stillwater, OK 74074.

TABLE 1. Number of artificially-sired cows that initiated first lactation within 1961 to 1974 in herds which were on test in 1975.

Year of freshening	Ayrshire	Guernsey	Holstein	Jersey	Brown Swiss
61	225	250	5817	338	62
62	393	557	10761	605	130
63	390	590	13677	740	152
64	513	643	16851	864	184
65	670	778	19647	1028	193
66	784	869	22748	1143	291
67	903	950	25358	1369	267
68	832	1019	28431	1382	310
69	1054	1122	32914	1699	324
70	1124	1202	35722	1830	352
71	1271	1203	39787	2016	366
72	1207	1242	44167	2239	459
73	1241	1257	46173	2397	411
74	1171	1259	49681	2714	450

year of first calving and a sire's estimated transmitting ability was used only in the year of a daughter's first calving. Transmitting ability was defined as one-half of additive genetic value.

Trends in transmitting ability of sires were obtained from the regression of the weighted average of sires' transmitting abilities for each year on year. For the  $p$ th year, the weighted average of sires' transmitting abilities was

$$\sum_i n_{ip} \hat{s}_i / n_{.p}$$

where  $n_{ip}$  is the number of daughters of sire

$i$  first freshening in the  $p$ th year, and  $\hat{s}_i$  is the predicted transmitting ability of the  $i$ th sire.

Genetic trends in the cow population were estimated from the regression of yearly average cow merit on year. For the  $p$ th year, the average cow merit was

$$\sum_j \hat{c}_j / m_p,$$

where  $m_p$  is the number of cows first freshening in the  $p$ th year, and  $\hat{c}_j$  is the predicted genetic merit of the  $j$ th cows.

TABLE 2. Number of naturally-sired cows that initiated first lactation within 1961 to 1974 in herds which were on test in 1975.

Year of freshening	Ayrshire	Guernsey	Holstein	Jersey	Brown Swiss
61	219	231	3628	368	54
62	356	484	6166	595	117
63	419	541	7331	707	122
64	396	574	8199	828	109
65	362	615	7911	922	105
66	333	666	8827	1018	144
67	388	701	10112	1107	137
68	369	741	11497	1133	143
69	441	710	14144	1310	163
70	431	742	16951	1230	203
71	369	745	21010	1244	191
72	617	725	26673	1309	219
73	659	604	32331	1165	208
74	665	698	42073	1255	243

TABLE 3. Average genetic merit of cows and transmitting ability of sires in kg: Ayrshires.

Year of fresh-ening	Genetic value AI cows	Transmitting ability AI sires	Genetic value non-AI cows
61	-174	-132	-452
62	-229	-137	-498
63	-225	-163	-518
64	-146	-87	-471
65	-1	56	-492
66	68	122	-403
67	80	144	-366
68	142	171	-338
69	189	197	-235
70	239	220	-209
71	220	193	-118
72	165	107	-130
73	158	91	-126
74	171	85	-110

## RESULTS AND DISCUSSION

The average predicted genetic merit of cows and transmitting ability of sires by year for the Ayrshire, Guernsey, Holstein, Jersey, and Brown Swiss breeds are in Tables 3 to 7. Annual genetic trends (regression of average predicted genetic merit and average predicted transmitting ability on year) are in Table 8.

Estimates of genetic gains per year among Brown Swiss and Ayrshire AI cows were not uniform, since estimates for Brown Swiss were negative from 1961 to 1964 and positive

from 1964 to 1971. Estimates for Ayrshires were positive from 1962 to 1970 and negative from 1970 to 1973. Estimates of genetic gains per year for Guernsey and Jersey cows were also not uniform since estimates for both breeds fluctuated over the 14 yr. Yearly differences in the average genetic merit of AI Holstein cows were relatively uniform.

Estimates of genetic improvement per year for non-AI Brown Swiss and Ayrshire cows paralleled the gains for AI cows. Genetic gains per year among Guernsey and Holstein non-AI cows, 35.4 and 31.0 kg, were greater

TABLE 4. Average genetic merit of cows and transmitting ability of sires in kg: Guernseys.

Year of fresh-ening	Genetic value AI cows	Transmitting ability AI sires	Genetic value non-AI cows
61	-120	-51	-273
62	-71	-21	-296
63	-24	-11	-242
64	-55	-16	-222
65	13	29	-213
66	65	70	-224
67	90	81	-215
68	89	86	-150
69	138	114	-106
70	153	109	-43
71	164	112	38
72	167	106	120
73	192	122	116
74	224	148	131

TABLE 5. Average genetic merit of cows and transmitting ability of sires in kg: Holsteins.

Year of fresh-ening	Genetic value AI cows	Transmitting ability AI sires	Genetic value non-AI cows
61	-161	-110	-370
62	-158	-99	-375
63	-58	-51	-307
64	-42	-30	-306
65	-10	-9	-278
66	29	22	-240
67	39	34	-221
68	55	39	-167
69	53	22	-113
70	80	43	-73
71	118	69	-38
72	152	95	-20
73	171	121	-32
74	205	156	-30

than genetic gains per year among Guernsey and Holstein AI cows, 25.4 and 26.1 kg. Genetic improvement per year for Jersey AI cows, 25.0 kg, was more than for non-AI cows, 12.8 kg. Comparisons of rates of genetic improvement of AI cows and non-AI cows may not be valid, however, because the estimate of genetic improvement in the non-AI cows may include unidentified AI sires classified as natural-service sires. The fraction of unidentified sires which are actually AI sires is unknown. The percentages of non-AI cows without an identified sire were 8.9, 9.9, 44.3,

12.8, and 10.9 among Ayrshire, Guernsey, Holstein, Jersey, and Brown Swiss cows. In 1974, the AI populations had 281, 93, 235, 340, and 224 kg advantages in estimates of average genetic merit for milk production over those classified as non-AI populations.

The rate of genetic improvement for milk has been small but relatively uniform in AI and non-AI populations of Holsteins over the 14 yr. The increase in genetic value for milk for AI and non-AI cows of the other four breeds also has been small but has not been uniform.

TABLE 6. Average genetic merit of cows and transmitting ability of sires in kg: Jerseys.

Year of fresh-ening	Genetic value AI cows	Transmitting ability AI sires	Genetic value non-AI cows
61	-79	-47	-257
62	-89	-51	-278
63	-71	-55	-232
64	-31	-12	-242
65	-13	1	-237
66	-13	1	-239
67	40	47	-249
68	72	69	-215
69	74	58	-187
70	103	73	-166
71	151	113	-124
72	180	133	-141
73	203	164	-97
74	209	171	-131

TABLE 7. Average genetic merit of cows and transmitting ability of sires in kg: Brown Swiss.

Year of fresh-ening	Genetic value AI cows	Transmitting ability AI sires	Genetic value non-AI cows
61	-419	-367	-796
62	-523	-379	-816
63	-556	-406	-744
64	-600	-499	-704
65	-555	-429	-632
66	-489	-372	-608
67	-443	-330	-490
68	-379	-276	-478
69	-320	-197	-531
70	-224	-110	-394
71	-184	-96	-520
72	-185	-79	-378
73	-91	-55	-413
74	-105	-36	-329

Estimates of annual genetic improvement from other studies are listed in Table 9. There are no reported genetic trends for non-AI populations, except for Holsteins, to compare with the genetic trends of non-AI cows that were estimated in this study. For Holsteins, the estimate of 31.0 kg/yr is higher than the only previous report of 23.0 kg/yr.

Trends in the average transmitting ability of AI sires for the Ayrshire, Guernsey, Holstein, Jersey, and Brown Swiss breeds were 23.7, 14.6, 17.9, 18.3, and 34.7 kg, which agree with the range of previous reports of 17 to 28, 16 to 17, 16 to 25, 11 to 17, and 23 to 41 kg. However, annual genetic trends in the AI cow populations for the five breeds were 36.1, 25.4, 26.1, 25.0, and 38.1 kg, consistently less than previously reported estimates of 34 to 55, 32 to 92, -3 to 166,

22 to 83, and 46 to 81 kg. In doubling the trends calculated from average transmitting abilities of sires to estimate genetic trends of cow populations, it is assumed that genetic improvement is constant each generation. This assumption implies that the genetic change due to sires is the same for each generation, which may not be appropriate (20).

### CONCLUSIONS

Genetic progress in dairy cattle has been considerable, although much slower than theoretically possible (Rendel and Robertson, [18]; Van Vleck, [23]). Trends in the genetic contribution of sires of the five breeds were less than previous reports, suggesting that estimating genetic trends of cow populations by doubling the trends in estimated transmitting abilities of sires is biased upwards.

TABLE 8. Regression of average genetic merit and average transmitting ability in kg on year.

Breed	Genetic value AI cows		Transmitting ability AI sires		Genetic value non-AI cows	
	$\bar{X}$	SE	$\bar{X}$	SE	$\bar{X}$	SE
Ayrshire	36.1	5.4	23.7	6.3	36.1	3.4
Guernsey	25.4	1.5	14.6	1.3	35.4	3.2
Holstein	26.1	1.6	17.9	1.2	31.0	1.7
Jersey	25.0	1.0	18.3	1.0	12.8	1.5
Brown Swiss	38.1	5.3	34.7	4.4	36.4	3.5

TABLE 9. Reported estimates of genetic trends in kg.

Refer- ence	Year	Breed	Location	AI or non-AI <sup>a</sup>	Method 1 <sup>b</sup>	Method 2 <sup>c</sup>
5	1959	Hol	New York	AI	...	33
25	1961	Hol	New York	AI	29	...
7	1961	Hol	One herd	Non-AI	23	...
18	1963	Hol	Texas	Both	3	...
1	1964	Jer		Both	-3, 81	...
11	1966	Jer	California	Both	62, 83	...
9	1967	Unknown	California	Both	34	...
2	1967	Hol	New York	Both	14	...
3	1967	Hol	One herd	AI	47	...
4	1968	Hol	Unknown	Both	51	...
14	1969	Hol	One herd	Both	45, 55	...
16	1971	Ayr	One herd	Both	0	...
8	1971	Jer	Unknown	Both	1	...
27	1972	Hol	SE US	AI	48, 18	...
12	1974	Jer	Florida	Both	38	...
17	1974	Hol		Both	53	...
20	1975	Hol		Both	25	...
13	1975	Hol		Both	92	...
15	1976	Hol		Both	33	...
6	1976	Ayr		Both	22	...
23	1977	Guer	Unknown	Both	9, 166	...
		Hol	Mid-west, US	Both	66-133	50
		Jer	Ontario	AI	...	42
		BS	Quebec	AI	...	46
		Ayr	Wisconsin	Both	79	43
		Guer	NE US	AI	...	34
		Hol			...	14
		Jer			...	46
		BS			...	22
		Ayr	NE US	AI	...	46
		Guer			...	55
		Hol			...	32
		Jer			...	39
		BS			...	34
					...	81

<sup>a</sup>If AI or non-AI was not indicated, it was assumed that both AI and non-AI were in the study.

<sup>b</sup>Weighted regression (Smith, 1962) or a modification of the method.

<sup>c</sup>Twice the linear regression of average sire merit on year.

## ACKNOWLEDGMENTS

This research was supported in part by a grant from Eastern Artificial Insemination Cooperative, Ithaca, NY.

## REFERENCES

- 1 Arave, C. W., R. C. Laben, and S. W. Mead. 1964. Measurement of change in twelve California dairy herds. *J. Dairy Sci.* 47:278.
- 2 Branton, C., D. L. Evans, J. R. Steele, and B. R. Farthing. 1967. Estimated genetic progress in milk and milk fat yield in a Louisiana Holstein herd. *J. Dairy Sci.* 50:974.
- 3 Burnside, E. B., and J. E. Legates. 1967. Estimation of genetic trends in dairy cattle populations. *J. Dairy Sci.* 50:1448.
- 4 Burnside, E. B., J. C. Rennie, and G. H. Bowman. 1968. Genetic trends and selection in a dairy cattle herd. *Can. J. Anim. Sci.* 48:243.
- 5 Elston, R. C. 1959. The estimation of genetic gain in milk yield due to sire selection over a period of time. Ph.D. Thesis, Cornell University, Ithaca, NY.
- 6 Everett, R. W., J. F. Keown, and E. E. Clapp. 1976. Production and stayability trends in dairy cattle. *J. Dairy Sci.* 59:1532.
- 7 Gaalaas, R. F., and R. D. Plowman. 1961. Effectiveness of statistical adjustments for yearly fluctuations in production. *J. Dairy Sci.* 44:1188.
- 8 Hargrove, G. L., and J. E. Legates. 1971. Biases in dairy sire evaluation attributable to genetic trend and female selection. *J. Dairy Sci.* 54:1041.
- 9 Harville, D. A., and C. R. Henderson. 1967. Environmental and genetic trends in production and their effects on sire evaluation. *J. Dairy Sci.* 50:870.
- 10 Henderson, C. R. 1975. Use of all relatives in intraherd prediction of breeding values and producing abilities. *J. Dairy Sci.* 58:1910.
- 11 Hillers, J. K., and A. E. Freeman. 1966. Two sources of genetic error in sire proofs. *J. Dairy Sci.* 49:1245.
- 12 Johnson, D. G. 1974. Genetic change in Minnesota dairy herds. *Diss. Abstr. Int.* 34(11):5267B.
- 13 Kennedy, B. W., and J. E. Moxley. 1975. Genetic trends among artificially bred Holsteins in Quebec. *J. Dairy Sci.* 58:1871.
- 14 Miller, P. D., W. E. Lentz, and C. R. Henderson. 1969. Comparison of contemporary daughters of young and progeny tested dairy sires. *J. Dairy Sci.* 52:926.
- 15 Olson, K. E., and E. L. Jensen. 1976. Estimation of genetic trend in Wisconsin Holsteins. *ADSA 71st Ann. Meeting*, p. 74.
- 16 Palmer, J. E., C. J. Wilcox, F. G. Martin, O. G. Verde, and R. E. Barrantes. 1971. Genetic trends in milk production in the Florida Agricultural Experiment Station Jersey herd. *J. Dairy Sci.* 54:455.
- 17 Powell, R. L., and A. E. Freeman. 1974. Genetic trend estimators. *J. Dairy Sci.* 57:1067.
- 18 Qureshi, A. W. 1963. Genetic trends in milk and milk fat production of Texas Dairy Herd Improvement Association cows. *J. Dairy Sci.* 46:629.
- 19 Rendel, J. M., and A. Robertson. 1950. Estimation of genetic gain in milk yield by selection in a closed herd of dairy cattle. *J. Genetics* 50:1.
- 20 Schaeffer, L. R., M. G. Freeman, and E. B. Burnside. 1975. Evaluation of Ontario Holstein dairy sires for milk and fat production. *J. Dairy Sci.* 58:109.
- 21 Slinger, W. D., E. L. Jensen, R. W. Everett, and C. R. Henderson. 1976. Programming cow evaluation. *J. Dairy Sci.* 59:1589.
- 22 Smith, C. 1962. Estimation of genetic change in farm livestock using field records. *Anim. Prod.* 4:239.
- 23 Ufford, G. R. 1977. Dairy sire evaluation using all lactation records in best linear unbiased production procedures. Ph.D. Thesis, Cornell University, Ithaca, NY.
- 24 Van Vleck, L. D. 1976. Theoretical and actual genetic progress in dairy cattle. Page 543 *In* Int. Conf. quantitative genetics. Iowa State University Press, Ames.
- 25 Van Vleck, L. D., and C. R. Henderson. 1961. Measurement of genetic trend. *J. Dairy Sci.* 44:1705.
- 26 Van Vleck, L. D., and C. R. Henderson. 1961. Improvement in production of New York Holsteins due to artificial insemination. *J. Dairy Sci.* 44:1328.
- 27 Verde, O. G., C. J. Wilcox, F. G. Martin, and C. W. Reaves. 1972. Genetic trends in milk production in Florida Dairy Herd Improvement Association herds. *J. Dairy Sci.* 55:1010.