

2018

Mucosal antibody responses to vaccines targeting SIV protease cleavage sites or fulllength Gag and Env proteins in Mauritian cynomolgus macaques

Hongzhao Li
University of Manitoba

Yan Hai
University of Manitoba

So-Yon Lim
Harvard Medical School

Nikki Toledo
University of Manitoba

Jose Crecente-Campo
Universidade de Santiago de Compostela

Follow this and additional works at: <http://digitalcommons.unl.edu/virologypub>



Part of the [Biological Phenomena](#), [Cell Phenomena](#), and [Immunity Commons](#), [Cell and Developmental Biology Commons](#), [Genetics and Genomics Commons](#), [Infectious Disease Commons](#), [Medical Immunology Commons](#), [Medical Pathology Commons](#), and the [Virology Commons](#)

Li, Hongzhao; Hai, Yan; Lim, So-Yon; Toledo, Nikki; Crecente-Campo, Jose; Schalk, Dane; Li, Lin; Omenge, Robert W.; Dacoba, Tamara G.; Liu, Lewis R.; Kashem, Mohammad Abul; Wan, Yanmin; Liang, Binhua; Li, Qingsheng; Rakasz, Eva; Schultz-Darken, Nancy; Alonso, Maria J.; Plummer, Francis A.; Whitney, James B.; and Luo, Ma, "Mucosal antibody responses to vaccines targeting SIV protease cleavage sites or fulllength Gag and Env proteins in Mauritian cynomolgus macaques" (2018). *Virology Papers*. 372.
<http://digitalcommons.unl.edu/virologypub/372>

This Article is brought to you for free and open access by the Virology, Nebraska Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Virology Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.


Authors

Hongzhao Li, Yan Hai, So-Yon Lim, Nikki Toledo, Jose Crecente-Campo, Dane Schalk, Lin Li, Robert W. Omange, Tamara G. Dacoba, Lewis R. Liu, Mohammad Abul Kashem, Yanmin Wan, Binhua Liang, Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, Maria J. Alonso, Francis A. Plummer, James B. Whitney, and Ma Luo

RESEARCH ARTICLE

Mucosal antibody responses to vaccines targeting SIV protease cleavage sites or full-length Gag and Env proteins in Mauritian cynomolgus macaques

Hongzhao Li¹, Yan Hai¹, So-Yon Lim², Nikki Toledo¹, Jose Crecente-Campo³, Dane Schalk⁴, Lin Li⁵, Robert W. Omange¹, Tamara G. Dacoba³, Lewis R. Liu¹, Mohammad Abul Kashem¹, Yanmin Wan⁶, Binhua Liang^{5,7}, Qingsheng Li⁶, Eva Rakasz⁸, Nancy Schultz-Darken⁴, Maria J. Alonso³, Francis A. Plummer^{1,5}, James B. Whitney^{2,9}, Ma Luo^{1,5*}



OPEN ACCESS

Citation: Li H, Hai Y, Lim S-Y, Toledo N, Crecente-Campo J, Schalk D, et al. (2018) Mucosal antibody responses to vaccines targeting SIV protease cleavage sites or full-length Gag and Env proteins in Mauritian cynomolgus macaques. PLoS ONE 13 (8): e0202997. <https://doi.org/10.1371/journal.pone.0202997>

Editor: Aftab A. Ansari, Emory University School of Medicine, UNITED STATES

Received: July 11, 2018

Accepted: August 13, 2018

Published: August 28, 2018

Copyright: © 2018 Li et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This study was supported by the following funding awarded to ML: a National Institute of Allergy and Infectious Diseases grant, with grant number: R01AI111805 and URL: <https://www.niaid.nih.gov/>; a Canadian Institutes of Health Research/Canadian HIV Vaccine Initiative Bridging grant (no grant number available), with URL: <http://www.cihr-irsc.gc.ca/e/42458.html>; and funding

1 Department of Medical Microbiology and Infectious Diseases, University of Manitoba, Winnipeg, MB, Canada, **2** Center for Virology and Vaccine Research, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, United States of America, **3** Center for Research in Molecular Medicine and Chronic Diseases (CIMUS), Campus Vida, Universidade de Santiago de Compostela, Santiago de Compostela, Spain, **4** Scientific Protocol Implementation Unit, Wisconsin National Primate Research Center, Madison, WI, United States of America, **5** National Microbiology Laboratory, Public Health Agency of Canada, Winnipeg, MB, Canada, **6** Nebraska Center for Virology, School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, United States of America, **7** Department of Biochemistry and Medical Genetics, University of Manitoba, Winnipeg, MB, Canada, **8** Immunology Services Unit, Wisconsin National Primate Research Center, Madison, WI, United States of America, **9** Ragon Institute of MGH, MIT, and Harvard, Cambridge, MA, United States of America

☯ These authors contributed equally to this work.

* Ma.Luo@phac-aspc.gc.ca, Ma.Luo@umanitoba.ca

Abstract

HIV mutates rapidly and infects CD4⁺ T cells, especially when they are activated. A vaccine targeting conserved, essential viral elements while limiting CD4⁺ T cell activation could be effective. Learning from natural immunity observed in a group of highly HIV-1 exposed seronegative Kenyan female sex workers, we are testing a novel candidate HIV vaccine targeting the 12 viral protease cleavage sites (PCSs) (the PCS vaccine), in comparison with a vaccine targeting full-length Gag and Env (the Gag/Env vaccine) in a Mauritian cynomolgus macaque/SIV model. In this study we evaluated these vaccines for induction of mucosal antibodies to SIV immunogens at the female genital tract. Bio-Plex and Western blot analyses of cervicovaginal lavage samples showed that both the PCS and Gag/Env vaccines can elicit mucosal IgG antibody responses to SIV immunogens. Significantly higher increase of anti-PCS antibodies was induced by the PCS vaccine than by the Gag/Env vaccine ($p < 0.0001$). The effect of the mucosal antibody responses in protection from repeated low dose pathogenic SIVmac251 challenges is being evaluated.

from National Microbiology Laboratory of Canada (no grant number available), with URL: <https://www.nml-inm.gc.ca/index-eng.htm>. Research reported in this publication was supported in part by the Office of the Director, National Institutes of Health under Award Number P51OD011106 to the Wisconsin National Primate Research Center, University of Wisconsin-Madison. This research was conducted in part at a facility constructed with support from Research Facilities Improvement Program grant numbers RR15459-01 and RR020141-01. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Development of an effective vaccine to human immunodeficiency virus type 1 (HIV) has proven to be a daunting task. Of the six HIV vaccine trials, the RV144 trial was the only one that demonstrated a modest efficacy (31.2%) [1, 2]. HIV primarily targets activated CD4⁺ T cells—a major arm of the immune system [1], apart from its capacity to mutate to evade immune recognition and generate extensive sequence variability [3–5]. Thus, a HIV vaccine generating immune response to conserved, functionally essential viral elements [6], and in the meantime limiting the generation of viral target cells [7, 8], could be more effective [1, 9–18].

The HIV protease cleaves Gag, Gag-Pol and Nef precursor proteins at twelve protease cleavage sites (PCSs) during viral maturation [18, 19]. The process is highly specific, temporally regulated and essential for generating infectious viral particles [20–25]. Even subtle disturbance can be sufficient to interrupt this delicately balanced process and drive it toward a non-productive end [20, 23, 24, 26]. Consistent with their critical function, the sequences surrounding the PCSs are highly conserved among major HIV subtypes [27]. Drugs targeting Gag that impair protease-mediated processing at specific Gag cleavage sites have been developed [28]. A HIV vaccine targeting the viral protease cleavage sites (PCSs) has been proposed for its ability to generate antiviral immune responses, disrupt HIV maturation and limit target cell activation [10, 18, 27].

Simian immunodeficiency virus (SIV) infection of nonhuman primates (NHPs) is currently the best animal model to test HIV vaccine strategies or study HIV pathogenesis [29–42]. To evaluate a candidate HIV vaccine targeting the PCSs (PCS vaccine), we use female Mauritian cynomolgus macaques (MCMs) and SIVmac as a model. The PCS vaccine [35] consists of twelve 20-mer peptides overlapping the twelve PCSs of SIVmac239 [43–45]. These peptides were delivered with recombinant vesicular stomatitis virus (rVSV) [46] and nanoparticles (NANO) [47–51]. In parallel, we also evaluate a vaccine targeting full-length Gag and Env (Gag/Env vaccine).

The majority of HIV infections worldwide are acquired through the mucosal routes during sexual contact [52]. Women are especially vulnerable through vaginal exposure to HIV in seminal fluids and constitute more than half of all infections globally [53]. Therefore, inducing a protective immune response at mucosal sites, including the female genital tract, is extremely important in HIV vaccine development [53–56]. In this study, we evaluated the PCS vaccine in comparison with the Gag/Env vaccine in generating mucosal antibody responses to different immunogens, which may impact on the outcome of viral challenge [57].

Materials and methods

Ethics statement

Female Mauritian cynomolgus macaques (MCMs) were pair-housed within the same experimental group during the immunization phase of the study with visual and auditory access to other conspecifics. Paired monkeys lived in two adjacent standard stainless-steel primate cages (27”L x 27”W x 32”H per cage). Rooms were maintained at 65–75°F, 30–70% humidity, and on a 12:12 light-dark cycle (ON: 0600, OFF: 1800). Standard nonhuman primate chow with fruit or vegetables was provided daily. In addition, we provided foraging activities and physical environmental enrichment at least weekly for both activities. All animals were observed at least twice daily for health or welfare issues. Sedation (ketamine alone, or ketamine/dexmedetomidine, atipamezole for reversal) was provided during the experimental procedures. The experiments were approved by the University of Wisconsin IACUC protocol (G005765) in accordance with the US Animal Welfare Act and following the recommendations of the

National Research Council *Guide for the Care and Use of Laboratory Animals*, 8th Edition and the Weatherall report, *The Use of Nonhuman Primates in Research*. The Wisconsin National Primate Research Center is fully accredited by AAALAC under the University of Wisconsin, Division of Vice-Chancellor for Research and Graduate Education.

Production of high titer rVSVpcs

The sequence of Simian immunodeficiency virus strain SIVmac239 was retrieved from the Los Alamos National Laboratory HIV database (<http://www.hiv.lanl.gov>). The nucleotide sequence encoding 20 amino acids (10 amino acids flanking each side of the cleavage site) overlapping each of the 12 PCSs of SIVmac239 was previously cloned in a recombinant vesicular stomatitis virus (rVSV) vector, pATX VSV-G, and packaged into rVSVpcs virus [35]. To generate large viral stocks for macaque immunization, VeroE6 cells were grown to 90% confluence in a T175 flask with 25 ml media, Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS), penicillin (100 U/ml), streptomycin (100 µg/ml) and L glutamine (2mM) (Invitrogen, CA, USA). 20 ml of the old culture media was removed and replaced with the same volume of fresh media 30 min before viral inoculation. Approximately 1×10^6 plaque forming units (pfu) of rVSVpcs virus was then added to the cell culture and allowed to proliferate for 12–24 hours until 90–100% cell death was observed. The culture supernatants were harvested and centrifuged at 180 g/min for 5 min at room temperature to remove cellular debris.

To concentrate and purify rVSVpcs using density gradient ultracentrifugation, 20 ml of the supernatants were gently layered on top of 8 ml of equilibration buffer (20 mM Tris-HCl, 0.1 M NaCl and 0.1 mM EDTA) containing 20% sucrose (Fisher Scientific) in an Ultra-Clear™ centrifugation tube (Beckman Coulter). To make the equilibration buffer, 20 ml of Ultra Pure 1M Tris-HCl pH7.5 (Invitrogen), 20 ml of 5M NaCl (Invitrogen) and 200 µl of 0.5M EDTA pH8.0 (Ambion) were diluted with sterile water to 1L. Sucrose was freshly added to the equilibration buffer on the day of viral purification. Ultracentrifugation was performed in a Beckman Coulter XPN-80 Ultracentrifuge using the SW 32 Ti rotor at 27,000 rpm for 2h at 4°C. After carefully removing the supernatants, viral pellets were resuspended in 1 ml DMEM containing 10% FBS, and finally stored as 100 µl aliquots in a -80°C freezer.

Viral titers were calculated using the TCID₅₀ calculator by Marco Binder (<https://www.researchgate.net/file.PostFileLoader.html?id=58dad730f7b67ea37125593f&assetKey=AS%3A476999471898624%401490736944531>), based on cytopathic effect of serially diluted viral stocks on VeroE6 cells.

Packaging of PCS peptides into nanoparticles (NANOpCs)

Twelve synthetic 20mer peptides overlapping the PCSs of SIVmac239 were associated to a biodegradable nanoparticle system formed by chitosan and dextran sulfate, as previously described [35].

Generation of rVSVGag/Env vaccine

Full-length Gag and Env coding sequences of SIVmac239 were synthesized and cloned in a Blue Heron pUC(-)MCS plasmid (BlueHeron Biotechnology, Bothell, WA, USA). Each gene sequence was flanked by an upstream MluI restriction site (AAACGCGT), Kozak sequence (GCCACC), start codon, and downstream stop codon and AvrII restriction site (CCTAGGTT). Using these restriction sites, the Gag and Env coding fragments were each sub-cloned into the rVSV vector pATX VSV-G, followed by confirmation with sequencing, and packaged into rVSVgag/env viruses, based on the previously described methods for rVSVpcs [35]. Large

stocks of high-titer purified rVSVgag/env viruses were produced with the same methods as described above for rVSVpcs. As a control vaccine vector, wild type virus (rVSV) was similarly produced. To test SIV protein expression, supernatants from VeroE6 cell cultures infected with these rVSVs were analyzed by Western blot.

Generation of NANOgag/env DNA vaccine

Using the above-mentioned Blue Heron pUC(-)MCS-Gag/Env plasmids as templates, full-length Gag and Env genes were PCR amplified with primers introducing an EcoRI restriction site upstream of the Kozak sequence and start codon and an XhoI restriction site downstream of the stop codon. These primers were:

Forward Gag primer: 5' CCGGAATTCGCCACCATGGGCGTGAGAACTCCG3'

Reverse Gag primer: 5' CCGCTCGAGCTACTGGTCTCCTCCAAAGAGAG3'

Forward Env primer: 5' CCGGAATTCGCCACCATGGGATGTCTTGGGAATC-3'

Reverse Env primer: 5' CCGCTCGAGTCACAAGAGAGTGAGCTCAAGC-3'

The PCR products were then sub-cloned into a DNA vaccine vector, pVAX1, between the EcoRI and XhoI sites, followed by confirmation with sequencing.

The resulting full-length Gag and Env-coding DNA constructs, pVAX1-Gag and pVAX1-Env, were each packaged into DNA vaccine nanoparticles (NANOgag and NANOenv, also collectively named NANOgag/env when administered together), according to the ionotropic gelation technique previously published [58]. Chitosan (Heppe Medical Chitosan GmbH, Halle, Germany) and tripolyphosphate (TPP, Sigma-Aldrich, St. Louis, MO, USA) were separately dissolved in ultrapure water at a concentration of 0.625 mg/mL and 2 mg/mL, respectively. Then 0.7 mL of TPP at the concentration of 2 mg/mL was mixed with 2.1 mL of 0.33 mg/mL solution of plasmid. This mixture was slowly added over 11.2 mL of chitosan solution at a concentration of 0.625 mg/mL, under magnetic stirring. Nanoparticles were instantly formed upon the addition, and the mixture was kept under stirring for 10 minutes. For the freeze-drying process, 0.65 mL of a filtered solution of trehalose at 150 mg/mL was added to 6.5 mL of the nanoparticle suspension. Samples were frozen at -80°C and subsequently freeze-dried (Genesis 25 ES, VirTis Model-Wizard 2.0, SP Industries, USA). Prior to animal administration, the freeze-dried particles were resuspended by adding 0.65 mL of water, vortexing for 10 seconds and shaking horizontally for 10 minutes. The physical characteristics of the freeze-dried particles after resuspension included: size (NANOgag = 235 ± 4 nm and NANOenv = 225 ± 18 nm), polydispersity index (both ≈ 0.2) and zeta potential (NANOgag = $+38 \pm 3$ mV and NANOenv = $+39 \pm 4$ mV).

To test SIV protein expression, the Gag and Env DNA vaccine constructs were each used to transfect HEK293T cells using lipofectamine 2000 (Thermo Fisher Scientific, Waltham, MA, USA). HEK293 T cells were cultured to 90% confluence (in the same media as used for VeroE6 described above) in 6-well plates. Before transfection, the media were replaced with 2 ml of antibiotic-free DMEM containing 10% FBS. 2.5 µg DNA and 10 µl lipofectamine 2000 were diluted separately in 250 µl Opti-MEM (Thermo Fisher Scientific). The diluted DNA was mixed with the diluted lipofectamine 2000 and incubated for 5 min at room temperature. The DNA-lipid complex was then added to the cells. Transfected cells were harvested 24 h post transfection. Approximately 1×10^6 cells were lysed with 50 µl RIPA lysis and extraction buffer (Thermo Fisher Scientific), then passed through a QIAshredder column (Qiagen) by centrifugation at 15,000 g for 2 min. The processed lysates were finally analyzed by Western blot.

Vaccination

Three groups of eight female MCMs were used in the study. Group 1, the PCS vaccine group received rVSVpcs (viruses expressing PCS peptides) and NANOpcs (nanoparticles containing

PCS peptides), Group 2, the Gag/Env vaccine group received rVSVgag/env (viruses expressing full-length Gag and Env proteins) and NANOgag/env (nanoparticles containing plasmid DNA encoding full-length Gag and Env), and Group 3, the Control group received vaccine vector controls (empty rVSV virus, and sterile water—the nanoparticle vehicle). One animal from the Gag/Env vaccine group was euthanized early due to severe health issues unrelated to vaccination, leaving seven animals in this group to complete the study. The vaccination procedure consisted of a prime with rVSVs at week 0, the first boost with rVSVs + NANOs at week 6, the second boost with NANOs at week 16, the third boost with rVSVs + NANOs at week 51 and the fourth boost with rVSVs at week 72, respectively. All rVSVs were administered intramuscularly via the quadriceps muscle, alternating L and R for successive vaccinations. The dose was at 1×10^6 pfu of each rVSVpcs per animal of the PCS Vaccine group or 6×10^6 pfu of rVSVgag and 6×10^6 pfu of rVSVenv per animal of the Gag/Env Vaccine group, except that for the 4th boost 1×10^8 pfu/rVSV type/animal were administered. All NANOs were administered intranasally, with each animal receiving NANO-delivered 50 μ g peptide for each of the twelve PCS peptides (the PCS vaccine group) or NANO-delivered plasmid DNA encoding Gag (500 μ g) and Env(500 μ g) (the Gag/Env vaccine group).

Cervicovaginal lavage (CVL) sample collection

The vaginal lumen of sedated animals was rinsed with 2–6 ml of phosphate buffered saline (PBS) non-traumatically using a needleless syringe. Within this volume range, the total amount of PBS used varied from animal to animal because of the varying sizes of their cervical vaults. The PBS was administered until the vault is full and then collected back into the same syringe. The fluid was gently flushed five times using the same syringe and repeated until 2–4 ml of fluid was collected. To take the volume variation into account, mucosal antibody concentrations were all normalized to total protein concentrations of the collected samples. The CVL sample collection was performed non-traumatically with extreme caution to avoid blood contamination from tissue damage. In addition, the animals were examined every day for menses and we have kept a daily menses record during the study. Due to variable dates of menstruation among individual female macaques and fixed dates of sampling schedule, it was practically difficult to avoid menstruation dates in all CVL collections. Indeed a small percentage of samples (11.25%) were collected on menstruation dates. To rule out effect of any potential menses blood contamination, we excluded data of the samples collected during menstruation from final analysis. All CVL sample aliquots were stored at -80°C until use.

Bio-Plex antibody assay

CVL IgG antibodies to SIV antigens were quantified by following a previously published Bio-Plex multiplexed protocol [35]. CVL total protein concentrations were determined using a NanoOrange[®] protein quantitation kit (Thermo Fisher Scientific) according to the manufacturer's protocol. The IgG antibody level was normalized with the total protein concentration.

Western blot

Western blot was used to detect: (1) SIV Gag and Env protein antigens expressed from rVSV and DNA Gag/Env vaccines; and (2) Vaccination-induced antibodies to purified recombinant Gag and Env proteins (NIH AIDS reagent program). Western blot was conducted following a previously published method [35] with slight modifications. For detection of antigens, 26 μ l of supernatants from rVSV-infected VeroE6 cell culture, or lysates of DNA vaccine-transfected HEK293T cells equivalent to 5×10^5 cells, were diluted and denatured in the format of a 40 μ l NuPAGE sample and loaded for SDS-PAGE, followed by blotting. Gag and Env protein

antigens were probed with standard mouse monoclonal antibodies (NIH AIDS Reagent Program) as listed in the published method [35]. For detection of antibodies, 1 µg of purified recombinant SIV proteins (NIH AIDS Reagent Program), SIVmac251 Gag (Catalog 1845) and SIVmac239 Env (Catalog 12797), were used as standard antigens for blotting. CVL samples from the control, PCS vaccine and Gag/Env vaccine groups of animals were 1:10 diluted and used as primary antibodies. IgG antibodies bound to the antigens were then detected using an anti-monkey IgG-HRP secondary antibody [35].

Statistical analysis

Mann Whitney's test was conducted using GraphPad Prism 7.04 to compare antibody responses among the PCS vaccine, Gag/Env vaccine and control groups. A *p* value less than 0.05 was defined as significant.

Results


Generation of the PCS vaccine and the Gag/Env vaccine

The PCS vaccine consists of twelve 20-amino acid peptides each overlapping one of the twelve PCSs of SIVmac239 (PCS1 through PCS12) (Fig 1A and 1B). The PCS immunogens were delivered in recombinant vesicular stomatitis virus (rVSVpcs) and nanoparticles (NANOpCs). rVSV is a non-pathogenic, replication competent viral vector. It can induce robust humoral and cellular immune responses, and unlike adenoviral vectors, lacks pre-existing human immunity [46]. The success of the vector was demonstrated by the safety and protective efficacy of an Ebola virus vaccine [59, 60]. Biodegradable NANO materials have great capacity for mucosal vaccination [47, 48, 50, 51, 61] as demonstrated by its successful nasal delivery of tetanus toxoid [61, 62] and hepatitis B surface antigen [47, 49, 63] subunit vaccines. For simplicity, the rVSVpcs and NANOpCs were collectively referred to as the PCS vaccine, as both were used in combination to immunize the same group of animals. The PCS vaccine was shown to induce plasma antibodies to the PCS peptides in a pilot study [35]. For comparison, we generated a rVSV vaccine with full-length Gag and Env genes of SIVmac239 (rVSVgag/env). The expression of Gag and Env proteins by rVSVgag/env was confirmed by Western blot using VeroE6 cell cultures infected by these viruses (Fig 1D). Gag and Env-expressing DNA vaccines in pVAX1 were also generated to be delivered with nanomaterial for boosting immune response to Gag and Env (NANOgag/env). The expression of Gag and Env proteins from the DNA vaccine constructs was confirmed by Western blot in transfected HEK293T cells (Fig 1D). The rVSVgag/env and NANOgag/env in combination were named the Gag/Env vaccine, to be compared with the PCS vaccine.

The immunization scheme

We carefully considered the routes of vaccination as they are well known factors to impact on the localized induction of immune responses. In general, systemic delivery of immunogens tends to elicit systemic responses and mucosal delivery of immunogens tends to induce mucosal immune responses [53]. However, systemically delivered viral vectors can also induce mucosal immune responses to HIV or SIV [64, 65]. Mucosal vaccination at one site stimulates immune responses in all mucosal sites, as well as systemic immune responses [53]. Intranasal immunization was reported to be the most effective at eliciting immune responses in the female genital tract [66], and this route induces greater IgG systemic responses than other mucosal routes, oral, rectal or vaginal [67]. It was proposed that systemic prime followed by mucosal boosting may help prevent induction of mucosal tolerance by initial mucosal

A.



B.

PCS peptides


PCS1: APSSGRGGNY/PVQQIGGNYV
 PCS2: GGPGQKARLM/AEALKEALAP
 PCS3: LAPVPIPF AA/AQQRGPRKPI
 PCS4: MAKCPDRQAG/FLGLGPWGKK
 PCS5: GPWGKKPRNF/PMAQVHQGLM
 PCS6: YGQMPRQTGG/FFRPWSMGKE
 PCS7: WSMGKEAPQF/PHGSSASGAD
 PCS8: LQGGDRGF AA/PQFSLWRRPV
 PCS9: LTALGMSLNF/PIAKVEPVKV
 PCS10: KDPIEGEETY/YTDGSCNKQS
 PCS11: LVSQGIRQVL/FLEKIEPAQE
 PCS12: NQGQYMNTPW/RNPAEEREKL

C.

Gag/Env peptides (non-PCS)

SIVgag: VGDHQAAMQIIRDIINEEAADWDL
 SIVenv1: NVTESFDAWNNTVTEQAIEDVWQLFETSIRPCVKLSP
 SIVenv2: RVTAEKYLKDQAQLNAWGCAFRQVCHTTVPWPNA

D.



E.




Fig 1. Vaccines targeting SIV protease cleavage sites (PCSs) or full Gag and Env proteins. (A) Diagram of the twelve protease cleavage sites (PCS1 through PCS12), located on three SIV polyproteins (Pr55Gag, Pr160Gag-Pol and Nef precursor), not drawn to scale. MA: matrix; CA: capsid; NC: nucleocapsid; TFP: transframe protein; PR: protease; RT: reverse transcriptase; and IN: integrase. (B) Peptide sequences of SIV immunogens in a conserved element vaccine targeting the PCSs (the PCS vaccine). Each sequence corresponds to -10 through +10 amino acid positions flanking each cleavage site. Slash (/) indicates the site of protease cleavage. These sequences were confirmed to be specific for SIV by NCBI protein BLAST and conserved among multiple SIV strains. The peptide immunogens were delivered as recombinant vesicular stomatitis viruses (rVSVpcs) and nanoparticles (NANOpCs). Peptide antigens with these sequences were also used in a Bio-Plex multiplexed assay to detect anti-PCS antibodies. (C) Sequences of three Gag or Env (non-PCS) peptides used in Bio-Plex to detect anti-Gag or Env antibodies, including one Gag peptide, named SIVgag, and two Env peptides, named SIVenv1 and SIVenv2. (D) Western blot analyses of protein expression from a full-length Gag and Env-based vaccine (the Gag/Env vaccine). VeroE6 cells were infected with recombinant vesicular stomatitis viruses (rVSVs) carrying full Gag or Env gene of SIVmac239 (rVSVgag/env) and the culture supernatants were analyzed by Western blot to detect Gag or Env protein expression using standard monoclonal antibodies (mAb, NIH AIDS Reagent Program) to Gag or Env. The full Gag and Env genes were also cloned into pVAX1 (a DNA vaccine vector), respectively, followed by NANO packaging (NANOgag/env). HEK293T cells were transfected with these DNA vaccines and analyzed by Western blot. (E) Vaccination scheme. Three groups (Control, PCS vaccine and Gag/Env vaccine) of eight female MCMs per group were primed and boosted on indicated weeks (wk). The Control group received empty rVSV virus and sterile water. One animal from the Gag/Env vaccine group was euthanized early due to severe health issues unrelated to vaccination, leaving seven animals in this group to complete the study. rVSV control vector (rVSV), rVSVpcs or rVSVgag/env was administered intramuscularly. NANO control vector (sterile water), NANOpCs or NANOgag/env was administered intranasally.

<https://doi.org/10.1371/journal.pone.0202997.g001>

vaccination, and should elicit both systemic and mucosal antibodies [53]. Taking all these into considerations, we chose to use rVSVs for systemic prime and boost through the intramuscular route (i.m.) and NANOs for mucosal boost through the intranasal route (i.n.). The resulting vaccination scheme consisted of a prime with rVSVs and four boosts with combinations of rVSVs and NANOs, as illustrated in Fig 1E.

Mucosal SIV-specific IgG antibodies elicited by vaccine modalities targeting different immunogens of SIVmac239

Protective mucosal immune responses to HIV is critical in preventing its mucosal transmission [52, 53]. In several passive and active NHP immunization experiments mucosal IgG antibodies showed protective effect against simian-human immunodeficiency virus (SHIV) acquisition (57–65). Therefore, we analyzed mucosal antigen-specific IgG responses after MCMs were primed and boosted with the PCS vaccine or the Gag/Env vaccine following the vaccination scheme illustrated in Fig 1E. Cervicovaginal lavage (CVL) samples were collected and measured for IgG responses to the twelve PCS peptides and three Gag/Env (non-PCS) peptides, SIVgag, SIVenv1 and SIVenv2 (Fig 1C) by Bio-Plex and to Gag and Env proteins by Western blot.

We first analyzed the dynamics of antibodies to each of the twelve PCS peptides (PCS1 – PCS12) throughout the vaccination procedure. These antibodies showed trend of increase in the PCS vaccine group compared to the Gag/Env vaccine and Control group after prime, the 1st boost and the 4th boost (Fig 2). Similar patterns were observed in the dynamics of antibodies to non-PCS Gag/Env peptides (SIVgag, SIVenv1 and SIVenv2) (Fig 3). We also observed variable antibody responses to PCS peptides among individual animals in the PCS vaccine (Figure A in S1 File), Gag/Env vaccine (Figure B in S1 File) and Control (Figure C in S1 File) groups, by comparing antibody levels between the baseline and one week after the last boost (4th). Six out of seven animals (85.7%) from the PCS vaccine group (Figure A in S1 File) and three out of six animals (50%) from the Gag/Env vaccine group (Figure B in S1 File) showed consistent increase in IgG antibodies to all PCS peptides. In contrast, only one out of eight animals (12.5%) from the control group showed a similar pattern (Figure C in S1 File). Similar results were also seen in antibody responses to Gag/Env peptides (Figures D-F in S1 File).

We quantified the effect of prime, each boost and the full vaccination process on antibody responses to each PCS. The PCS vaccine group showed trends of higher fold increase of




Fig 2. Dynamics of PCS-specific IgG antibodies in cervicovaginal secretions. Cervicovaginal lavage (CVL) samples from the vaccination experiments illustrated in Fig 1E were quantified for levels of IgG antibodies to each PCS peptide (y axis, expressed as ratios of anti-PCS IgG concentration to total protein concentration $\times 10^9$) by a Bio-Plex multiplexed antibody assay, at indicated time points (x axis, weeks post prime). The Control group received empty rVSV virus and sterile water. Data are shown as each value from individual animals with median line, following subtraction of pre-vaccination values. Total animal numbers per group examined are $n = 8$ (Control or PCS group) or $n = 7$ (Gag/Env group). However, for technical stringency, animal samples collected on menstruation dates were excluded from analysis to rule out any potential menses blood contamination. As a result, for some of the data points in the graph, fewer individual values than the total animal number ($n = 8$ or $n = 7$) were available. Grey areas indicate a week interval post prime, 1st boost or 4th boost, with trends of antibody increase in the PCS vaccine group in response to vaccination. The trends did not reach statistical significance.

<https://doi.org/10.1371/journal.pone.0202997.g002>

antibodies (after prime, the 1st boost, the 4th boost, or the full vaccination process) than the other groups (Fig 4).

We then calculated the effect of vaccines on total antibody responses to all PCSs by including antibodies to all PCS sites collectively as anti-PCS antibodies since they all target the viral




Fig 3. Dynamics of Gag/Env-specific IgG antibodies in cervicovaginal secretions. Cervicovaginal lavage (CVL) samples from the vaccination experiments illustrated in Fig 1E were quantified for IgG antibodies to Gag/Env (non-PCS) peptides, SIVgag, SIVenv1 and SIVenv2 (y axis, expressed as ratios of anti-non-PCS IgG concentration to total protein concentration $\times 10^9$) by a Bio-Plex multiplexed antibody assay, at indicated time points (x axis, weeks post prime). The Control group received empty rVSV virus and sterile water. Data are shown as each value from individual animals with median line, following subtraction of pre-vaccination values. Total animal numbers per group examined are $n = 8$ (Control or PCS group) or $n = 7$ (Gag/Env group). However, for technical stringency, animal samples collected on menstruation dates were excluded from analysis to rule out any potential menses blood contamination. As a result, for some of the data points in the graph, fewer individual values than the total animal number ($n = 8$ or $n = 7$) were available. Grey areas indicate a week interval post prime, 1st boost or 4th boost, with trends of antibody increase in the PCS vaccine group in response to vaccination. The trends did not reach statistical significance.

<https://doi.org/10.1371/journal.pone.0202997.g003>




Fig 4. Fold changes of mucosal antibodies to each PCS peptide in response to vaccinations. Graphs show fold changes of mucosal IgG antibodies to each PCS peptide (1 through 12) between the time of a prime/boost and one week after that single prime/boost (Prime, 1st, 2nd, 3rd or 4th), or between the baseline (the start of the full vaccination procedure) and one week after the last boost (the end of the full vaccination procedure) (Full). The Control group received empty rVSV virus and sterile water. Data are shown as each value from individual animals with interquartile range and median line. Total animal numbers per group examined are $n = 8$ (Control or PCS group) or $n = 7$ (Gag/Env group). However, for technical stringency, animal samples collected on menstruation dates were excluded from analysis to rule out any potential menses blood contamination. As a result, for some of the data points in the graph, fewer individual values than the total animal number ($n = 8$ or $n = 7$) were available. Fold changes of antibodies in response to prime, the 1st boost, the 4th boost and the full vaccination procedure showed trends of increase in the PCS vaccine group, compared to the other groups. The trends did not reach statistical significance.

<https://doi.org/10.1371/journal.pone.0202997.g004>




Fig 5. Fold changes of collective anti-PCS antibodies in response to vaccinations. Mucosal IgG antibodies to all individual PCS types (PCS1 through PCS12) were collectively treated as anti-PCS antibodies for calculation. Graph shows fold changes between the time of a prime/boost and one week after that single prime/boost (Prime, 1st, 2nd, 3rd or 4th), or between the baseline (the start of the full vaccination procedure) and one week after the last boost (the end of the full vaccination procedure) (Full). The Control group received empty rVSV virus and sterile water. Total animal numbers per group examined are $n = 8$ (Control or PCS group) or $n = 7$ (Gag/Env group). However, for technical stringency, animal samples collected on menstruation dates were excluded from analysis to rule out any potential menses blood contamination. Bars represent mean \pm SEM. The PCS vaccine group demonstrated significantly higher fold induction of anti-PCS antibodies than the Gag/Env vaccine and Control groups, in response to prime, the 1st boost, the 4th boost and the full vaccination procedure, as determined by Mann Whitney's test: **** $p < 0.0001$.

<https://doi.org/10.1371/journal.pone.0202997.g005>

protease cleavage function. The fold changes of total anti-PCS antibodies, in response to prime, the 1st boost, the 4th boost or the full vaccination process, were significantly higher in the PCS vaccine group than those in the Gag/Env vaccine group and the Control group ($p < 0.0001$) (Fig 5). Among these, the mean fold changes induced by the 4th boost were 12.353 (PCS vaccine), 3.465 (Gag/Env vaccine) and 1.792 (Control), and those induced by the full vaccination process were 25.466 (PCS vaccine), 3.005 (Gag/Env vaccine) and 1.853 (Control) (Fig 5). These results indicated that the PCS vaccine can effectively induce mucosal anti-PCS antibodies at the female genital tract and has a stronger effect than the Gag/Env vaccine.

We also examined the effect of vaccines on antibody responses to Gag/Env. The PCS vaccine group showed trends of higher fold increase in antibodies to two non-PCS Gag/Env peptides, SIVgag and SIVenv1, after prime, the 1st boost, the 4th boost, and the full vaccination process, in comparison to the Control group (Fig 6A). The fold increase of antibodies to the three non-PCS Gag/Env peptides was not apparent in the Gag/Env group or the Control group (Fig 6A). We then tested antibody reactivity by Western blot using Gag and Env proteins, which were expected to detect antibodies that recognize epitopes not limited to the three Gag/Env peptides. Antibodies of vaccinated animals from the PCS and Gag/Env vaccine groups recognized purified recombinant Gag protein (rGag) (Fig 6B). While antibodies from the Gag/Env group showed strong reactivity to purified recombinant Env protein (rEnv), those from the PCS vaccine group demonstrated weak but clear reactivity to rEnv (Fig 6B). These results indicated that the PCS and Gag/Env vaccines can induce mucosal IgG antibodies to Gag and Env.

Cross-reactivity of the vaccine-induced antibodies

Bio-Plex antibody assays suggested that the PCS vaccine, which delivers PCS peptides, induced antibodies that recognize non-PCS Gag/Env peptides (Figs 3 and 6A). The antibody cross-reactivity was supported by Western blot analyses demonstrating that these antibodies reacted





Fig 6. Induction of antibodies to Gag/Env antigens. (A) Graphs show fold changes of mucosal IgG antibodies to Gag/Env (non-PCS) peptides (SIVgag, SIVenv1 and SIVenv2) between the time of a prime/boost and one week after that single prime/boost (Prime, 1st, 2nd, 3rd or 4th), or between the baseline (the start of the full vaccination procedure) and one week after the last boost (the end of the full vaccination procedure) (Full). The Control group received empty rVSV virus and sterile water. Data are shown as each value from individual animals with interquartile range and median line. Total animal numbers per group examined are n = 8 (Control or PCS group) or n = 7 (Gag/Env group). However, for technical stringency, animal samples collected on menstruation dates were excluded from analysis to rule out any potential menses blood contamination. As a result, for some of the data points in the graph, fewer individual values than the total animal number (n = 8 or n = 7) were available. Fold changes of antibodies to SIVgag and SIVenv1 in response to prime, the 1st boost, the 4th boost and the full vaccination procedure showed trends of increase in the PCS vaccine group, compared to the other groups. The trends did not reach statistical significance. (B) Reactivity of mucosal IgG antibodies to Gag and Env proteins. Western blot membranes containing purified recombinant Gag (rGag) or Env (rEnv) protein (NIH AIDS Reagent Program) were used to probe anti-Gag/Env IgG antibodies in CVL samples, collected at one week after the last boost from animals of the PCS vaccine (animal ID: cy0759), Gag/Env vaccine (animal ID: cy0784) and Control (animal ID: cy0779) groups, respectively.

<https://doi.org/10.1371/journal.pone.0202997.g006>

to rGag and rEnv proteins (Fig 6B). The rGag protein contains peptide sequences of PCSs (PCS1 through PCS5) (Fig 1), thus it was expected that the antibodies induced by the PCS vaccine could recognize Gag. However, rEnv does not contain any PCS sequence (Fig 1), therefore the induction of anti-Env antibodies by the PCS vaccine was unexpected. While the underlying mechanism(s) of the cross-reactivity remain to be understood, these data indicated that the PCS vaccine can generate antibodies to both PCS peptides and Env.

Discussion

In this study we examined mucosal antibodies induced by two different modalities of candidate HIV/SIV vaccines, a vaccine targeting short peptide sequences overlapping the 12 protease cleavage sites and a vaccine targeting full Gag and Env of SIVmac239. Since 90% of HIV transmissions occur through the mucosal route [68] and male to female sexual transmissions account for more than half of all HIV infections [53], it is important to test whether a candidate vaccine can induce mucosal immune responses to HIV/SIV antigens. Our study showed that both the PCS vaccine and Gag/Env vaccine can induce cervicovaginal mucosal IgG antibodies to SIV antigens, including PCSs, Gag and Env. The PCS vaccine preferentially generated mucosal IgG antibodies to the PCS peptides, whereas the Gag/Env vaccine generated much stronger mucosal IgG antibodies to Env.

Most of current vaccine studies on anti-HIV antibodies are focused on Env-specific antibodies. However, it was also shown that antibodies to Gag and Pol correlated with natural and post vaccination HIV control [69–74], suggesting that antibodies targeting Gag and Pol could be protective. Not only Env, but also Gag and Pol were included in the vaccine used for the RV144 trial, the only vaccine so far with efficacy against HIV acquisition [2]. A previous pre-clinical study also showed that Gag was required for protection against SIVsmE660 challenge [75]. Potential protective effect of the mucosal antibodies to PCSs, Gag and Env induced by different modalities of vaccines in the current study will be evaluated in repeated low dose SIV-mac251 intravaginal challenges.

We observed that without SIV-specific immunization, antibodies with reactivity to SIV antigens can be present in some MCMs (at variable levels). The mechanism(s) for generating these antibodies remain to be understood. We speculate that endogenous or environmental antigens may induce antibodies with cross-reactivity to SIV antigens and subject to regulation by external stimuli [35, 37]. Our results also showed that the PCS vaccine elicited mucosal antibodies not only to the PCSs but also to non-PCS Gag and Env antigens. While several possibilities may account for this observation [35, 37], one of them may be the cross-reactivity of the anti-PCS antibodies to the non-PCS antigens [76]. Although there is no similarity in primary

sequences between the PCS and non-PCS peptides, they may have similar conformational structures that could contribute to cross-reactivity [76]. It is important to note that the induction of antibodies with reactivity to non-PCS peptides did not distract the PCS vaccine away from its intended targets, the PCSs, since the PCS vaccine was shown to effectively elicit antibodies to these targets. The cross-reactivity of the PCS vaccine-induced antibodies expands the antigen spectrum to additional Gag and Env epitopes. As a result, the PCS vaccine may have the potential to target both PCS-based viral maturation and Env-mediated viral entry. These will need to be further investigated.

In conclusion, our study showed that the PCS vaccine and Gag/Env vaccine can induce mucosal IgG responses to SIV immunogens, including PCS peptides and Gag and Env proteins. The effect of these vaccine-induced mucosal antibodies in protecting macaques from pathogenic SIVmac251 low dose intravaginal challenges will be determined in on-going studies.

Supporting information

S1 File. A single PDF file including Figures A through F.
(PDF)

Acknowledgments

We would like to thank staff at Wisconsin National Primate Research Center Scientific Protocol Implementation Unit and Immunology Services Unit for important technical support. We recognize Dr. Stuart Shapiro, NIH Vaccine Research Program, and Dr. Matthew Gilmour, National Microbiology Laboratory of Canada, for important support and discussion. This work was supported by a NIH grant (R01AI111805), a CIHR/CHVI bridging grant and funding from National Microbiology Laboratory of Canada.

Research reported in this publication was supported in part by the Office of the Director, National Institutes of Health under Award Number P51OD011106 to the Wisconsin National Primate Research Center, University of Wisconsin-Madison. This research was conducted in part at a facility constructed with support from Research Facilities Improvement Program grant numbers RR15459-01 and RR020141-01. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author Contributions

Conceptualization: Hongzhao Li, Yan Hai, Binhua Liang, Qingsheng Li, Eva Rakasz, Maria J. Alonso, Francis A. Plummer, James B. Whitney, Ma Luo.

Data curation: Hongzhao Li, Yan Hai, Jose Crecente-Campo.

Formal analysis: Hongzhao Li, Yan Hai, So-Yon Lim, Nikki Toledo, Jose Crecente-Campo.

Funding acquisition: Binhua Liang, Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, Maria J. Alonso, Francis A. Plummer, James B. Whitney, Ma Luo.

Investigation: Hongzhao Li, Yan Hai, So-Yon Lim, Nikki Toledo, Jose Crecente-Campo, Dane Schalk, Lin Li, Robert W. Omange, Tamara G. Dacoba, Lewis R. Liu, Mohammad Abul Kashem, Yanmin Wan, Binhua Liang, Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, Maria J. Alonso, James B. Whitney, Ma Luo.

Methodology: Hongzhao Li, Yan Hai, Nikki Toledo, Jose Crecente-Campo, Dane Schalk, Lin Li, Robert W. Omenge, Tamara G. Dacoba, Binhua Liang, Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, Maria J. Alonso, James B. Whitney.

Project administration: Binhua Liang, Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, Maria J. Alonso, Francis A. Plummer, James B. Whitney, Ma Luo.

Resources: Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, Maria J. Alonso, Francis A. Plummer, James B. Whitney, Ma Luo.

Supervision: Binhua Liang, Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, Maria J. Alonso, Francis A. Plummer, James B. Whitney, Ma Luo.

Validation: Hongzhao Li, Yan Hai, Nikki Toledo, Jose Crecente-Campo.

Visualization: Hongzhao Li, Yan Hai, So-Yon Lim.

Writing – original draft: Hongzhao Li.

Writing – review & editing: Yan Hai, So-Yon Lim, Jose Crecente-Campo, Binhua Liang, Qingsheng Li, Eva Rakasz, Nancy Schultz-Darken, James B. Whitney, Ma Luo.

References

1. Tomaras GD, Plotkin SA. Complex immune correlates of protection in HIV-1 vaccine efficacy trials. *Immunol Rev.* 2017 Jan; 275(1):245–61. <https://doi.org/10.1111/immr.12514> PMID: 28133811
2. Rerks-Ngarm S, Pitisuttithum P, Nitayaphan S, Kaewkungwal J, Chiu J, Paris R, et al. Vaccination with ALVAC and AIDSVAX to prevent HIV-1 infection in Thailand. *N Engl J Med.* 2009 Dec 3; 361(23):2209–20. <https://doi.org/10.1056/NEJMoa0908492> PMID: 19843557
3. Lynch RM, Wong P, Tran L, O'Dell S, Nason MC, Li Y, et al. HIV-1 fitness cost associated with escape from the VRC01 class of CD4 binding site neutralizing antibodies. *J Virol.* 2015 Apr; 89(8):4201–13. <https://doi.org/10.1128/JVI.03608-14> PMID: 25631091
4. Sheng Z, Schramm CA, Connors M, Morris L, Mascola JR, Kwong PD, et al. Effects of Darwinian Selection and Mutability on Rate of Broadly Neutralizing Antibody Evolution during HIV-1 Infection. *PLoS Comput Biol.* 2016 May; 12(5):e1004940. <https://doi.org/10.1371/journal.pcbi.1004940> PMID: 27191167
5. Sunshine JE, Larsen BB, Maust B, Casey E, Deng W, Chen L, et al. Fitness-Balanced Escape Determines Resolution of Dynamic Founder Virus Escape Processes in HIV-1 Infection. *J Virol.* 2015 Oct; 89(20):10303–18. <https://doi.org/10.1128/JVI.01876-15> PMID: 26223634
6. Rolland M, Nickle DC, Mullins JI. HIV-1 group M conserved elements vaccine. *PLoS Pathog.* 2007 Nov; 3(11):e157. <https://doi.org/10.1371/journal.ppat.0030157> PMID: 18052528
7. Begaud E, Chartier L, Marechal V, Iperio J, Leal J, Versmisse P, et al. Reduced CD4 T cell activation and in vitro susceptibility to HIV-1 infection in exposed uninfected Central Africans. *Retrovirology.* 2006 Jun 22; 3:35. <https://doi.org/10.1186/1742-4690-3-35> PMID: 16792805
8. Hope TJ. Inflammation weakens HIV prevention. *Nat Med.* 2018 Apr 10; 24(4):384–5. <https://doi.org/10.1038/nm.4534> PMID: 29634687
9. Routy JP, Mehraj V. Potential contribution of gut microbiota and systemic inflammation on HIV vaccine effectiveness and vaccine design. *AIDS Res Ther.* 2017 Sep 12; 14(1):48. <https://doi.org/10.1186/s12981-017-0164-9> PMID: 28893288
10. Luo M, Daniuk CA, Diallo TO, Capina RE, Kimani J, Wachihi C, et al. For protection from HIV-1 infection, more might not be better: a systematic analysis of HIV Gag epitopes of two alleles associated with different outcomes of HIV-1 infection. *J Virol.* 2012 Jan; 86(2):1166–80. <https://doi.org/10.1128/JVI.05721-11> PMID: 22072744
11. VaxGen vaccine trial fails the test but may offer insights. *AIDS Alert.* 2003 Apr; 18(4):41, 3–5. PMID: 12710402
12. HIV vaccine failure prompts Merck to halt trial. *Nature.* 2007 Sep 27; 449(7161):390. <https://doi.org/10.1038/449390c> PMID: 17898737
13. McCarthy M. HIV vaccine fails in phase 3 trial. *Lancet.* 2003 Mar 1; 361(9359):755–6. [https://doi.org/10.1016/S0140-6736\(03\)12669-4](https://doi.org/10.1016/S0140-6736(03)12669-4) PMID: 12620743

14. Pal R, Venzon D, Letvin NL, Santra S, Montefiori DC, Miller NR, et al. ALVAC-SIV-gag-pol-env-based vaccination and macaque major histocompatibility complex class I (A*01) delay simian immunodeficiency virus SIVmac-induced immunodeficiency. *J Virol*. 2002 Jan; 76(1):292–302. <https://doi.org/10.1128/JVI.76.1.292-302.2002> PMID: 11739694
15. Plotkin SA. The RV144 Thai HIV vaccine trial. *Hum Vaccin*. 2010 Feb; 6(2):159.
16. Vaccari M, Poonam P, Franchini G. Phase III HIV vaccine trial in Thailand: a step toward a protective vaccine for HIV. *Expert Rev Vaccines*. 2010 Sep; 9(9):997–1005. <https://doi.org/10.1586/erv.10.104> PMID: 20822342
17. Willyard C. Tiny steps towards an HIV vaccine. *Nature*. 2010 Jul 15; 466(7304):S8. <https://doi.org/10.1038/nature09238> PMID: 20631706
18. Li H, Omange RW, Plummer FA, Luo M. A novel HIV vaccine targeting the protease cleavage sites. *AIDS Res Ther*. 2017 Sep 12; 14(1):51. <https://doi.org/10.1186/s12981-017-0174-7> PMID: 28893268
19. de Oliveira T, Engelbrecht S, Janse van Rensburg E, Gordon M, Bishop K, zur Megede J, et al. Variability at human immunodeficiency virus type 1 subtype C protease cleavage sites: an indication of viral fitness? *J Virol*. 2003 Sep; 77(17):9422–30. <https://doi.org/10.1128/JVI.77.17.9422-9430.2003> PMID: 12915557
20. Muller B, Anders M, Akiyama H, Welsch S, Glass B, Nikovics K, et al. HIV-1 Gag processing intermediates trans-dominantly interfere with HIV-1 infectivity. *J Biol Chem*. 2009 Oct 23; 284(43):29692–703. <https://doi.org/10.1074/jbc.M109.027144> PMID: 19666477
21. Pettit SC, Moody MD, Wehbie RS, Kaplan AH, Nantermet PV, Klein CA, et al. The p2 domain of human immunodeficiency virus type 1 Gag regulates sequential proteolytic processing and is required to produce fully infectious virions. *J Virol*. 1994 Dec; 68(12):8017–27. PMID: 7966591
22. Pettit SC, Henderson GJ, Schiffer CA, Swanstrom R. Replacement of the P1 amino acid of human immunodeficiency virus type 1 Gag processing sites can inhibit or enhance the rate of cleavage by the viral protease. *J Virol*. 2002 Oct; 76(20):10226–33. <https://doi.org/10.1128/JVI.76.20.10226-10233.2002> PMID: 12239298
23. Pettit SC, Clemente JC, Jeung JA, Dunn BM, Kaplan AH. Ordered processing of the human immunodeficiency virus type 1 GagPol precursor is influenced by the context of the embedded viral protease. *J Virol*. 2005 Aug; 79(16):10601–7. <https://doi.org/10.1128/JVI.79.16.10601-10607.2005> PMID: 16051852
24. Pettit SC, Lindquist JN, Kaplan AH, Swanstrom R. Processing sites in the human immunodeficiency virus type 1 (HIV-1) Gag-Pro-Pol precursor are cleaved by the viral protease at different rates. *Retrovirology*. 2005; 2:66. <https://doi.org/10.1186/1742-4690-2-66> PMID: 16262906
25. Pettit SC, Everitt LE, Choudhury S, Dunn BM, Kaplan AH. Initial cleavage of the human immunodeficiency virus type 1 GagPol precursor by its activated protease occurs by an intramolecular mechanism. *J Virol*. 2004 Aug; 78(16):8477–85. <https://doi.org/10.1128/JVI.78.16.8477-8485.2004> PMID: 15280456
26. Kaplan AH, Zack JA, Knigge M, Paul DA, Kempf DJ, Norbeck DW, et al. Partial inhibition of the human immunodeficiency virus type 1 protease results in aberrant virus assembly and the formation of noninfectious particles. *J Virol*. 1993 Jul; 67(7):4050–5. PMID: 8510215
27. Luo M, Capina R, Daniuk C, Tuff J, Peters H, Kimani M, et al. Immunogenicity of sequences around HIV-1 protease cleavage sites: potential targets and population coverage analysis for a HIV vaccine targeting protease cleavage sites. *Vaccine*. 2013 Jun 24; 31(29):3000–8. <https://doi.org/10.1016/j.vaccine.2013.04.057> PMID: 23664989
28. Adamson CS, Salzwedel K, Freed EO. Virus maturation as a new HIV-1 therapeutic target. *Expert Opin Ther Targets*. 2009 Aug; 13(8):895–908. <https://doi.org/10.1517/14728220903039714> PMID: 19534569
29. Lu S, Arthos J, Montefiori DC, Yasutomi Y, Manson K, Mustafa F, et al. Simian immunodeficiency virus DNA vaccine trial in macaques. *J Virol*. 1996 Jun; 70(6):3978–91. PMID: 8648735
30. Lu S, Manson K, Wyand M, Robinson HL. SIV DNA vaccine trial in macaques: post-challenge necropsy in vaccine and control groups. *Vaccine*. 1997 Jun; 15(8):920–3. PMID: 9234548
31. Pal R, Kalyanaraman VS, Nair BC, Whitney S, Keen T, Hocker L, et al. Immunization of rhesus macaques with a polyvalent DNA prime/protein boost human immunodeficiency virus type 1 vaccine elicits protective antibody response against simian human immunodeficiency virus of R5 phenotype. *Virology*. 2006 May 10; 348(2):341–53. <https://doi.org/10.1016/j.viro.2005.12.029> PMID: 16460776
32. Lu S, Grimes Serrano JM, Wang S. Polyvalent AIDS vaccines. *Curr HIV Res*. 2010 Dec; 8(8):622–9. PMID: 21054250
33. Chen Y, Wang S, Lu S. DNA Immunization for HIV Vaccine Development. *Vaccines (Basel)*. 2014 Feb 25; 2(1):138–59.

34. Antony JM, MacDonald KS. A critical analysis of the cynomolgus macaque, *Macaca fascicularis*, as a model to test HIV-1/SIV vaccine efficacy. *Vaccine*. 2015 Jun 17; 33(27):3073–83. <https://doi.org/10.1016/j.vaccine.2014.12.004> PMID: 25510387
35. Li H, Nykoluk M, Li L, Liu LR, Omange RW, Soule G, et al. Natural and cross-inducible anti-SIV antibodies in Mauritian cynomolgus macaques. *PLoS One*. 2017; 12(10):e0186079. <https://doi.org/10.1371/journal.pone.0186079> PMID: 28982126
36. Li H, Omange RW, Czarnecki C, Correia-Pinto JF, Crecente-Campo J, Richmond M, et al. Mauritian cynomolgus macaques with M3M4 MHC genotype control SIVmac251 infection. *J Med Primatol*. 2017 Aug; 46(4):137–43. <https://doi.org/10.1111/jmp.12300> PMID: 28748659
37. Li H, Li L, Liu LR, Omange RW, Toledo N, Kashem MA, et al. Hypothetical endogenous SIV-like antigens in Mauritian cynomolgus macaques. *Bioinformatics*. 2018; 14(2):48–52. <https://doi.org/10.6026/97320630014048> PMID: 29618899
38. Policicchio BB, Sette P, Xu C, Haret-Richter G, Dunsmore T, Pandrea I, et al. Emergence of resistance mutations in simian immunodeficiency virus (SIV)-infected rhesus macaques receiving non-suppressive antiretroviral therapy (ART). *PLoS One*. 2018; 13(2):e0190908. <https://doi.org/10.1371/journal.pone.0190908> PMID: 29466356
39. He T, Brocca-Cofano E, Policicchio BB, Sivanandham R, Gautam R, Raehtz KD, et al. Cutting Edge: T Regulatory Cell Depletion Reactivates Latent Simian Immunodeficiency Virus (SIV) in Controller Macaques While Boosting SIV-Specific T Lymphocytes. *J Immunol*. 2016 Dec 15; 197(12):4535–9. <https://doi.org/10.4049/jimmunol.1601539> PMID: 27837106
40. Policicchio BB, Xu C, Brocca-Cofano E, Raehtz KD, He T, Ma D, et al. Multi-dose Romidepsin Reactivates Replication Competent SIV in Post-antiretroviral Rhesus Macaque Controllers. *PLoS Pathog*. 2016 Sep; 12(9):e1005879. <https://doi.org/10.1371/journal.ppat.1005879> PMID: 27632364
41. Pandrea I, Xu C, Stock JL, Frank DN, Ma D, Policicchio BB, et al. Antibiotic and Antiinflammatory Therapy Transiently Reduces Inflammation and Hypercoagulation in Acutely SIV-Infected Pigtailed Macaques. *PLoS Pathog*. 2016 Jan; 12(1):e1005384. <https://doi.org/10.1371/journal.ppat.1005384> PMID: 26764484
42. Carias AM, Allen SA, Fought AJ, Kotnik Halavaty K, Anderson MR, Jimenez ML, et al. Increases in Endogenous or Exogenous Progestins Promote Virus-Target Cell Interactions within the Non-human Primate Female Reproductive Tract. *PLoS Pathog*. 2016 Sep; 12(9):e1005885. <https://doi.org/10.1371/journal.ppat.1005885> PMID: 27658293
43. Whitney JB, Oliveira M, Detorio M, Guan Y, Wainberg MA. The M184V mutation in reverse transcriptase can delay reversion of attenuated variants of simian immunodeficiency virus. *J Virol*. 2002 Sep; 76(17):8958–62. <https://doi.org/10.1128/JVI.76.17.8958-8962.2002> PMID: 12163615
44. Whitney JB, Wainberg MA. Impaired RNA incorporation and dimerization in live attenuated leader-variants of SIVmac239. *Retrovirology*. 2006; 3:96. <https://doi.org/10.1186/1742-4690-3-96> PMID: 17184529
45. Burwitz BJ, Pendley CJ, Greene JM, Detmer AM, Lhost JJ, Karl JA, et al. Mauritian cynomolgus macaques share two exceptionally common major histocompatibility complex class I alleles that restrict simian immunodeficiency virus-specific CD8+ T cells. *J Virol*. 2009 Jun; 83(12):6011–9. <https://doi.org/10.1128/JVI.00199-09> PMID: 19339351
46. Racine T, Kobinger GP, Arts EJ. Development of an HIV vaccine using a vesicular stomatitis virus vector expressing designer HIV-1 envelope glycoproteins to enhance humoral responses. *AIDS Res Ther*. 2017 Sep 12; 14(1):55. <https://doi.org/10.1186/s12981-017-0179-2> PMID: 28893277
47. Paolicelli P, Prego C, Sanchez A, Alonso MJ. Surface-modified PLGA-based nanoparticles that can efficiently associate and deliver virus-like particles. *Nanomedicine (Lond)*. 2010 Aug; 5(6):843–53.
48. Prego C, Garcia M, Torres D, Alonso MJ. Transmucosal macromolecular drug delivery. *J Control Release*. 2005 Jan 3; 101(1–3):151–62. <https://doi.org/10.1016/j.jconrel.2004.07.030> PMID: 15588901
49. Prego C, Paolicelli P, Diaz B, Vicente S, Sanchez A, Gonzalez-Fernandez A, et al. Chitosan-based nanoparticles for improving immunization against hepatitis B infection. *Vaccine*. 2010 Mar 19; 28(14):2607–14. <https://doi.org/10.1016/j.vaccine.2010.01.011> PMID: 20096389
50. Tobio M, Gref R, Sanchez A, Langer R, Alonso MJ. Stealth PLA-PEG nanoparticles as protein carriers for nasal administration. *Pharm Res*. 1998 Feb; 15(2):270–5. PMID: 9523314
51. Correia-Pinto JF, Csaba N, Alonso MJ. Vaccine delivery carriers: insights and future perspectives. *Int J Pharm*. 2013 Jan 2; 440(1):27–38. <https://doi.org/10.1016/j.ijpharm.2012.04.047> PMID: 22561794
52. Tebit DM, Ndambi N, Weinberg A, Quinones-Mateu ME. Mucosal transmission of human immunodeficiency virus. *Curr HIV Res*. 2012 Jan 1; 10(1):3–8. PMID: 22264040

53. Pavot V, Rochereau N, Lawrence P, Girard MP, Genin C, Verrier B, et al. Recent progress in HIV vaccines inducing mucosal immune responses. *AIDS*. 2014 Jul 31; 28(12):1701–18. <https://doi.org/10.1097/QAD.0000000000000308> PMID: 25009956
54. Stieh DJ, Maric D, Kelley ZL, Anderson MR, Hattaway HZ, Beifuss BA, et al. Vaginal challenge with an SIV-based dual reporter system reveals that infection can occur throughout the upper and lower female reproductive tract. *PLoS Pathog*. 2014 Oct; 10(10):e1004440. <https://doi.org/10.1371/journal.ppat.1004440> PMID: 25299616
55. Stieh DJ, Matias E, Xu H, Fought AJ, Blanchard JL, Marx PA, et al. Th17 Cells Are Preferentially Infected Very Early after Vaginal Transmission of SIV in Macaques. *Cell Host Microbe*. 2016 Apr 13; 19(4):529–40. <https://doi.org/10.1016/j.chom.2016.03.005> PMID: 27078070
56. Joag VR, McKinnon LR, Liu J, Kidane ST, Yudin MH, Nyanga B, et al. Identification of preferential CD4 + T-cell targets for HIV infection in the cervix. *Mucosal Immunol*. 2016 Jan; 9(1):1–12. <https://doi.org/10.1038/mi.2015.28> PMID: 25872482
57. Nguyen M, Pean P, Lopalco L, Nouhin J, Phoung V, Ly N, et al. HIV-specific antibodies but not t-cell responses are associated with protection in seronegative partners of HIV-1-infected individuals in Cambodia. *J Acquir Immune Defic Syndr*. 2006 Aug 1; 42(4):412–9. <https://doi.org/10.1097/01.qai.0000222289.97825.35> PMID: 16837821
58. Calvo P, Remunan-Lopez C, Vila-Jato JL, Alonso MJ. Chitosan and chitosan/ethylene oxide-propylene oxide block copolymer nanoparticles as novel carriers for proteins and vaccines. *Pharm Res*. 1997 Oct; 14(10):1431–6. PMID: 9358557
59. Agnandji ST, Huttner A, Zinser ME, Njuguna P, Dahlke C, Fernandes JF, et al. Phase 1 Trials of rVSV Ebola Vaccine in Africa and Europe. *N Engl J Med*. 2016 Apr 28; 374(17):1647–60. <https://doi.org/10.1056/NEJMoa1502924> PMID: 25830326
60. Henao-Restrepo AM, Camacho A, Longini IM, Watson CH, Edmunds WJ, Egger M, et al. Efficacy and effectiveness of an rVSV-vectored vaccine in preventing Ebola virus disease: final results from the Guinea ring vaccination, open-label, cluster-randomised trial (Ebola Ca Suffitl). *Lancet*. 2017 Feb 4; 389(10068):505–18. [https://doi.org/10.1016/S0140-6736\(16\)32621-6](https://doi.org/10.1016/S0140-6736(16)32621-6) PMID: 28017403
61. Vila A, Sanchez A, Janes K, Behrens I, Kissel T, Vila Jato JL, et al. Low molecular weight chitosan nanoparticles as new carriers for nasal vaccine delivery in mice. *Eur J Pharm Biopharm*. 2004 Jan; 57(1):123–31. PMID: 14729088
62. Vila A, Sanchez A, Evora C, Soriano I, Vila Jato JL, Alonso MJ. PEG-PLA nanoparticles as carriers for nasal vaccine delivery. *J Aerosol Med*. 2004 Summer; 17(2):174–85. <https://doi.org/10.1089/0894268041457183> PMID: 15294069
63. Vicente S, Diaz-Freitas B, Peleteiro M, Sanchez A, Pascual DW, Gonzalez-Fernandez A, et al. A polymer/oil based nanovaccine as a single-dose immunization approach. *PLoS One*. 2013; 8(4):e62500. <https://doi.org/10.1371/journal.pone.0062500> PMID: 23614052
64. Lin SW, Cun AS, Harris-McCoy K, Ertl HC. Intramuscular rather than oral administration of replication-defective adenoviral vaccine vector induces specific CD8+ T cell responses in the gut. *Vaccine*. 2007 Mar 8; 25(12):2187–93. <https://doi.org/10.1016/j.vaccine.2006.11.044> PMID: 17229501
65. Tatsis N, Lin SW, Harris-McCoy K, Garber DA, Feinberg MB, Ertl HC. Multiple immunizations with adenovirus and MVA vectors improve CD8+ T cell functionality and mucosal homing. *Virology*. 2007 Oct 10; 367(1):156–67. <https://doi.org/10.1016/j.virol.2007.05.028> PMID: 17590405
66. Pavot V, Rochereau N, Genin C, Verrier B, Paul S. New insights in mucosal vaccine development. *Vaccine*. 2012 Jan 5; 30(2):142–54. <https://doi.org/10.1016/j.vaccine.2011.11.003> PMID: 22085556
67. Lema D, Garcia A, De Sanctis JB. HIV vaccines: a brief overview. *Scand J Immunol*. 2014 Jul; 80(1):1–11. <https://doi.org/10.1111/sji.12184> PMID: 24813074
68. Ruprecht RM, Lakhashe SK. Antibody-mediated immune exclusion of HIV. *Curr Opin HIV AIDS*. 2017 May; 12(3):222–8. <https://doi.org/10.1097/COH.0000000000000369> PMID: 28422786
69. Tjiam MC, Sariputra L, Armitage JD, Taylor JP, Kelleher AD, Tan DB, et al. Control of early HIV-1 infection associates with plasmacytoid dendritic cell-reactive opsonophagocytic IgG antibodies to HIV-1 p24. *AIDS*. 2016 Nov 28; 30(18):2757–65. <https://doi.org/10.1097/QAD.0000000000001242> PMID: 27603291
70. Tjiam MC, Taylor JP, Morshidi MA, Sariputra L, Burrows S, Martin JN, et al. Viremic HIV Controllers Exhibit High Plasmacytoid Dendritic Cell-Reactive Opsonophagocytic IgG Antibody Responses against HIV-1 p24 Associated with Greater Antibody Isotype Diversification. *J Immunol*. 2015 Jun 1; 194(11):5320–8. <https://doi.org/10.4049/jimmunol.1402918> PMID: 25911748
71. French MA, Tjiam MC, Abudulai LN, Fernandez S. Antiviral Functions of Human Immunodeficiency Virus Type 1 (HIV-1)-Specific IgG Antibodies: Effects of Antiretroviral Therapy and Implications for Therapeutic HIV-1 Vaccine Design. *Front Immunol*. 2017; 8:780. <https://doi.org/10.3389/fimmu.2017.00780> PMID: 28725225

72. French MA, Tanaskovic S, Law MG, Lim A, Fernandez S, Ward LD, et al. Vaccine-induced IgG2 anti-HIV p24 is associated with control of HIV in patients with a 'high-affinity' FcγRIIIa genotype. *AIDS*. 2010 Aug 24; 24(13):1983–90. <https://doi.org/10.1097/QAD.0b013e328333c1ce0> PMID: 20634666
73. French MA, Abdulai LN, Fernandez S. Isotype Diversification of IgG Antibodies to HIV Gag Proteins as a Therapeutic Vaccination Strategy for HIV Infection. *Vaccines (Basel)*. 2013 Aug 9; 1(3):328–42.
74. French MA, Center RJ, Wilson KM, Fleyfel I, Fernandez S, Schorcht A, et al. Isotype-switched immunoglobulin G antibodies to HIV Gag proteins may provide alternative or additional immune responses to 'protective' human leukocyte antigen-B alleles in HIV controllers. *AIDS*. 2013 Feb 20; 27(4):519–28. <https://doi.org/10.1097/QAD.0b013e32835cb720> PMID: 23364441
75. Schell JB, Bahl K, Folta-Stogniew E, Rose N, Buonocore L, Marx PA, et al. Antigenic requirement for Gag in a vaccine that protects against high-dose mucosal challenge with simian immunodeficiency virus. *Virology*. 2015 Feb; 476:405–12. <https://doi.org/10.1016/j.virol.2014.12.027> PMID: 25591175
76. Craig L, Sanschagrin PC, Rozek A, Lackie S, Kuhn LA, Scott JK. The role of structure in antibody cross-reactivity between peptides and folded proteins. *J Mol Biol*. 1998 Aug 07; 281(1):183–201. <https://doi.org/10.1006/jmbi.1998.1907> PMID: 9680484




Figure A. The PCS vaccine group: Mucosal IgG responses to PCSs in each individual animal. Cervicovaginal lavage (CVL) samples of the PCS vaccine group animals from the vaccination experiments illustrated in Figure 1E were analyzed for antibodies to PCS peptides using a Bio-Plex multiplexed antibody assay. Each panel represents one animal, with animal ID and MHC haplotypes (MHC I, MHC II) shown on top. Data from one animal (ID cy0758) were excluded due to sample collection on concurrent menstruation dates. Antibody levels (ratios of anti-PCS IgG concentration to total protein concentration, timed by 10^9) in each animal were compared between the baseline (indicated as “B” on x axis) and after the vaccination procedure (one week after the last boost, indicated as “V” on x axis). Each line represents IgG antibody levels to one PCS type, connecting data points before and after vaccination.




Figure B. The Gag/Env vaccine group: Mucosal IgG responses to PCSs in each individual animal. Antibodies to PCS peptides were analyzed and graphed as in Figure A. Data from two animals were excluded due to the following reasons: cy0782 was euthanized early due to health issues unrelated to vaccination. Sample collection of cy0788 involved concurrent menstruation dates.




Figure C. The Control group: Mucosal IgG responses to PCSs in each individual animal. Antibodies to PCS peptides were analyzed and graphed as in Figure A.




Figure D. PCS vaccine group: Mucosal IgG responses to Gag/Env peptides (non-PCS) in each individual animal. Antibodies to Gag/Env peptides were analyzed and graphed similarly to Figure A.




Figure E. Gag/Env vaccine group: Mucosal IgG responses to Gag/Env peptides (non-PCS) in each individual animal. Antibodies to Gag/Env peptides were analyzed and graphed similarly to Figure B.




Figure F. Control group: Mucosal IgG responses to Gag/Env peptides (non-PCS) in each individual animal. Antibodies to Gag/Env peptides were analyzed and graphed similarly to Figure C.