

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and
Plant Health Inspection Service

February 2007

Blackbird Use and Damage of Wildlife Conservation Sunflower Plots: The Second Year

Jonathan M. Raetzman
North Dakota State University

George M. Linz
USDA/APHIS/WS National Wildlife Research Center, george_m_linz@yahoo.com

William J. Bleier
North Dakota State University

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Environmental Sciences Commons](#)

Raetzman, Jonathan M.; Linz, George M.; and Bleier, William J., "Blackbird Use and Damage of Wildlife Conservation Sunflower Plots: The Second Year" (2007). *USDA National Wildlife Research Center - Staff Publications*. 435.

https://digitalcommons.unl.edu/icwdm_usdanwrc/435

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Published in National Sunflower Association Research Forum Papers 2006.

The annual NSA Research Forum is highly regarded as the premier conference at which private and public researchers present the results of their latest sunflower research. For nearly a quarter of a century, the National Sunflower Association has produced the workshop as a forum for sunflower research to be shared with the public.

Papers from NSA Research Forums 2003 and later are online at:

<http://www.sunflowernsa.com/research/default.asp?contentID=70>

NSA Contact Information

National Sunflower Association
4023 State Street
Bismarck, ND 58503-0690
(701) 328-5100
(888) 718-7033

Blackbird Use and Damage of Wildlife Conservation Sunflower Plots: The Second Year

Jonathan M. Raetzman¹, George M. Linz², William J. Bleier¹

1. Department of Biological Sciences, NDSU Fargo, ND 2. USDA WS NWRC Bismarck, ND

Introduction

Annual blackbird damage to sunflower is valued at \$5 to \$10 million. Perhaps more importantly, the National Sunflower Association has identified blackbirds as the key reason for growers to abandon sunflower in areas of prime production. North Dakota and South Dakota together produce approximately 70% of sunflower grown in the United States. This crop adds \$906 million to the regional economy. To date, cost effective and environmentally safe methods to reduce bird damage to sunflower remain elusive. In the 1980's, scientists from the National Wildlife Research Center (NWRC) showed that "decoy" sunflower plots can greatly reduce bird damage to local commercial sunflower fields. At the time, this idea was abandoned due to lack of grower support. Support has increased, however, over the past 10 years due to conservation interests and new federal farm programs. In 2004, Wildlife Services (WS) along with North Dakota State University (NDSU) started the "Wildlife Conservation Sunflower Plots" (WCSP) study.



Sunflower field with large flock of feeding blackbirds

Methods

From 2004-2006, we plan to cooperate with sunflower growers in North Dakota and South Dakota with historical blackbird problems, who are willing to plant at least one 8-ha sunflower plot near a likely blackbird roost. Farmers received \$370.50/ha up to maximum of approximately 8 ha. In 2004, 17 WCSP were planted in North Dakota (Figure 1). In 2006 the number of planted WCSP will increase as funds are available.

From 20 August to 18 October 2004 and from 23 August to 20 October 2005, weekly damage assessment surveys were conducted in the plots. These surveys were done by randomly choosing 24 damage assessment sites within each 8-ha WCSP. Over each field season, the same damage assessment sites were surveyed each time. We recorded head size (cm²) and damage amount (cm²) for each individual head on a weekly basis. We will assume that every seed eaten in a WCSP is a seed saved from a commercial sunflower field. In addition all commercial sunflower fields within 1.5 miles of WCSP were surveyed for damage at the end of the growing season.

WCSP Sites

In 2004 and 2005 the WCSP were located in the Prairie Pothole Region of North Dakota. For 2004 we had 13 WCSP and in 2005 we had 21 WCSP for a total of 34 WCSP (Figure 1).




Figure 1. 2004 and 2005 WCSP locations

Results

We conducted weekly damage surveys on the WCSP in 2004 and 2005 (Figure 2). Damage for 2004 and 2005 was similar, however in 2004 the overall damage increased noticeably around 4 September and then again near 25 September. In 2004, the mean percent damage (Figure 3) was 38.6% (range 1.6% - 90.5%) in 13 WCSP. In 2004, the mean percent damage (Figure 3) was 4.6% (range 0.7%-23.4%) in 24 commercial sunflower fields. The 21 WCSP, in 2005, averaged a mean percent damage (Figure 3) of 31.6% (range 0.7% - 93.7%). The 49 commercial sunflower fields, in 2005, had a mean percent damage (Figure 3) of 3.7% (range 0.0%- 24.1%). These data show that WCSP are being damaged significantly more by blackbirds than commercial sunflower fields in this study.




Figure 2. Percent sunflower damage in individual WCSP over time.




Figure 3. 2004 and 2005, WCSP and commercial sunflower field average damage

Goals

Our goal is to provide baseline data on the use of WCSP for wildlife resource managers to make an informed decision on the environmental effects and efficacy of these plantings for reducing blackbird damage to commercial sunflower and providing habitat for wildlife. We will 1) evaluate the efficacy of small sunflower plots for reducing blackbird damage to commercial fields, 2) identify and quantify avian use of WCSP, 3) identify and quantify habitat factors that influence bird use of WCSP, and 4) estimate the economic costs and benefits of using WCSP to reduce bird damage to sunflower.



Picture left: Red-winged Blackbird



Damage survey in progress

Acknowledgements

We thank SD/ND Wildlife Services, Bismarck, ND, for coordinating and supporting the WCSP. We thank all of the sunflower growers who participated in this study. This project was funded by United States Department of Agriculture and Wildlife Services.