

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Roman L. Hruska U.S. Meat Animal Research  
Center

U.S. Department of Agriculture: Agricultural  
Research Service, Lincoln, Nebraska

---

2013

## Cross-sectional Study Examining *Salmonella enterica* Carriage in Subiliac Lymph Nodes of Cull and Feedlot Cattle at Harvest

Sara E. Gragg  
*Texas Tech University*

Guy H. Loneragan  
*Texas Tech University*

Mindy M. Brashears  
*Texas Tech University*

Terrance M. Arthur  
*USDA-ARS*

Joseph M. Bosilevac  
*USDA-ARS, mick.bosilevac@ars.usda.gov*

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/hruskareports>

---

Gragg, Sara E.; Loneragan, Guy H.; Brashears, Mindy M.; Arthur, Terrance M.; Bosilevac, Joseph M.; Kalchayanand, Norasak; Wang, Rong; Schmidt, John W.; Brooks, J. Chance; Shackelford, Steven D.; Wheeler, Tommy L.; Brown, Tyson R.; Edrington, Thomas S.; and Brichta-Harhay, Dayna M., "Cross-sectional Study Examining *Salmonella enterica* Carriage in Subiliac Lymph Nodes of Cull and Feedlot Cattle at Harvest" (2013). *Roman L. Hruska U.S. Meat Animal Research Center*. 420.  
<https://digitalcommons.unl.edu/hruskareports/420>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Roman L. Hruska U.S. Meat Animal Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Sara E. Gragg, Guy H. Loneragan, Mindy M. Brashears, Terrance M. Arthur, Joseph M. Bosilevac, Norasak Kalchayanand, Rong Wang, John W. Schmidt, J. Chance Brooks, Steven D. Shackelford, Tommy L. Wheeler, Tyson R. Brown, Thomas S. Edrington, and Dayna M. Brichta-Harhay

# Cross-sectional Study Examining *Salmonella enterica* Carriage in Subiliac Lymph Nodes of Cull and Feedlot Cattle at Harvest

Sara E. Gragg,<sup>1</sup> Guy H. Loneragan,<sup>1</sup> Mindy M. Brashears,<sup>1</sup> Terrance M. Arthur,<sup>2</sup> Joseph M. Bosilevac,<sup>2</sup> Norasak Kalchayanand,<sup>2</sup> Rong Wang,<sup>2</sup> John W. Schmidt,<sup>2</sup> J. Chance Brooks,<sup>1</sup> Steven D. Shackelford,<sup>2</sup> Tommy L. Wheeler,<sup>2</sup> Tyson R. Brown,<sup>1</sup> Thomas S. Edrington,<sup>3</sup> and Dayna M. Brichta-Harhay<sup>2</sup>

## Abstract

Bovine peripheral lymph nodes (LNs), including subiliac LNs, have been identified as a potential source of human exposure to *Salmonella enterica*, when adipose trim containing these nodes is incorporated into ground beef. In order to gain a better understanding of the burden of *S. enterica* in peripheral LNs of feedlot and cull cattle, a cross-sectional study was undertaken in which 3327 subiliac LNs were collected from cattle at harvest in seven plants, located in three geographically distinct regions of the United States. Samples were collected in three seasons: Fall 2010, Winter/Spring 2011, and Summer/Fall 2011. A convenience sample of 76 LNs per day, 2 days per season (approximately 1 month apart), was collected per plant, from carcasses held in the cooler for no less than 24 h. Every 10<sup>th</sup> carcass half on a rail was sampled, in an attempt to avoid oversampling any single cohort of cattle. Median point estimates of *S. enterica* contamination were generally low (1.3%); however, median *Salmonella* prevalence was found to be greater in subiliac LNs of feedlot cattle (11.8%) compared to those of cull cattle (0.65%). Enumeration analysis of a subset of 618 feedlot cattle LNs showed that 67% of those harboring *S. enterica* (97 of 144) did so at concentrations ranging from <0.1 to 1.8 log<sub>10</sub> CFU/g, while 33% carried a higher burden of *S. enterica*, with levels ranging from 1.9 to >3.8 log<sub>10</sub> CFU/g. Serotyping of *S. enterica* isolated identified 24 serotypes, with the majority being Montevideo (44.0%) and Anatum (24.8%). Antimicrobial susceptibility phenotypes were determined for all isolates, and the majority (86.1%) were pansusceptible; however, multidrug-resistant isolates (8.3%) were also occasionally observed. As *Salmonella* contained within LNs are protected from carcass interventions, research is needed to define opportunities for mitigating the risk of *Salmonella* contamination in LNs of apparently healthy cattle.

## Introduction

**N**ONTYPHOIDAL *SALMONELLA ENTERICA* is a significant cause of morbidity and mortality in the United States and is estimated to cause over 1 million illnesses each year (Mead *et al.*, 1999; Guo *et al.*, 2011; Scallan *et al.*, 2011). The vast majority of these cases are foodborne and while produce, poultry, and eggs account for most illnesses, beef has been identified as a vehicle of exposure, associated with sporadic cases and outbreaks (Guo *et al.*, 2011). *S. enterica* (hereafter referred to as *Salmonella*) has been frequently recovered from

the hides and feces of healthy cattle (Barkocy-Gallagher *et al.*, 2003; Loneragan and Brashears, 2005; Brichta-Harhay *et al.*, 2008; Kunze *et al.*, 2008; Loneragan *et al.*, 2012), and it is theorized that animal carriage of *Salmonella* ultimately contributes to ground beef contamination (Bosilevac *et al.*, 2009).

During harvest, hides of cattle are likely the primary source of *Salmonella* contamination of carcass surfaces (Barkocy-Gallagher *et al.*, 2001; Barkocy-Gallagher *et al.*, 2003; Arthur *et al.*, 2008; Brichta-Harhay *et al.*, 2008). Accordingly, substantial effort is afforded to preventing contamination of carcasses and removal of contamination through strategies

<sup>1</sup>International Center for Food Industry Excellence (ICFIE), Texas Tech University, Department of Animal and Food Sciences, Lubbock, Texas.

<sup>2</sup>U.S. Department of Agriculture, Agricultural Research Service, U.S. Meat Animal Research Center, Clay Center, Nebraska.

<sup>3</sup>U.S. Department of Agriculture, Agricultural Research Service, Southern Plains Agricultural Research Center, College Station, Texas.

Names are necessary to report factually on available data; however, the U.S. Department of Agriculture (USDA) neither guarantees nor warrants the standard of the product, and the use of the name by USDA implies no approval of the product to the exclusion of others that may also be suitable. The USDA is an equal opportunity provider and employer.

outlined in plant Hazard Analysis Critical Control Point plans. These strategies appear quite effective, as *Salmonella* prevalence after antimicrobial intervention application is typically undetectable or less than 1% (Barkocy-Gallagher *et al.*, 2003; Rivera-Betancourt *et al.*, 2004; Brichta-Harhay *et al.*, 2008). However, despite successful control of surface contamination, it is still possible for *Salmonella* to be recovered from ground beef. In a study of commercial ground beef from seven regions of the United States ( $n = 4136$  samples collected over 2 years), *Salmonella* was recovered from 4.2% of ground beef samples (Bosilevac *et al.*, 2009). Similarly, government testing of ground beef indicates that *Salmonella* contamination averages around 2.1%, and that little improvement in contamination has been achieved over the past decade (FSIS, 2011), even while the prevalence of *Escherichia coli* O157:H7 in ground beef has declined more than 70% (from 0.80% in 2001 to 0.23% in 2010 [FSIS, 2012]).

Research suggests that pathogen contamination of ground beef also can occur by means of contaminated lymph nodes (LNs) (Arthur *et al.*, 2008). Cattle possess many LNs located within fatty tissues that are frequently incorporated into ground beef and thus have the potential to contaminate the final product. Contaminated LNs may explain the difference in *Salmonella* prevalence between postintervention carcasses or trim, and ground beef. When present in LNs, *Salmonella* are protected from chemical and thermal antimicrobial carcass interventions, and as a consequence sanitary harvest procedures may not address this potential source of contamination. Therefore, the objective of this cross-sectional study was to gain a better understanding of the burden of *S. enterica* in peripheral LNs of harvest-ready cattle, by examining the point prevalence, serotypes, and antimicrobial susceptibility phenotypes present in subiliac LNs of cull and fed cattle at harvest, in three regions of the United States.

## Materials and Methods

### Sample collection

Subiliac LNs were collected from carcasses of cull (cows/bulls and dairy) and feedlot (steers/heifers) cattle that had passed federal inspection at commercial abattoirs. Samples were collected during three time periods including September, October, and November 2010 (Fall 2010), February and March 2011 (Winter/Spring 2011), and July, August, and September 2011 (Summer/Fall 2011). A convenience sample of seven commercial processing plants consisting of three plants that primarily harvest feedlot cattle and four that primarily harvest cull cows was initially enrolled. An eighth plant, harvesting primarily feedlot cattle, contributed two sets of samples during the Summer/Fall 2011 sample period. Participating abattoirs were located in regions 2, 3, and 5 of the microbiological monitoring regions defined by the Beef Industry Food Safety Council (Fig. 1). A convenience sample of 76 LNs per day, 2 days per season (approximately 1 month apart), was collected per plant, from postintervention carcasses that had been held in the cooler for no less than 24 h. Approximately every 10<sup>th</sup> carcass half on a rail was sampled, resulting in sample collection from cattle harvested over 1–2 production days, in an attempt to avoid oversampling any single cohort of cattle. However, the possibility remains that multiple nodes from a single cohort reside in a given sample set.




FIG. 1. Map of the Beef Industry Food Safety Council microbiological monitoring regions from which fed and cull cattle lymph nodes were obtained.

Samples were shipped to Texas Tech University in Lubbock, Texas or the U. S. Meat Animal Research Center in Clay Center, Nebraska for processing.

### LN sample processing and *Salmonella* detection

LNs were processed as previously described (Brichta-Harhay *et al.*, 2012). Briefly, surrounding fat and fascia were trimmed from LN samples, which were weighed, surface sterilized by submersion in a boiling water bath, placed into individual filtered sample bags (Nasco, Atlanta, GA), pulverized using a rubber mallet, and then enriched in 80 mL of tryptic soy broth (Becton Dickinson, Sparks, MD) by incubating at 25°C for 2 h and then 42°C for 12 h. Enrichments were subjected to immunomagnetic separation using anti-*Salmonella* beads (Dynabeads; Invitrogen, Carlsbad, CA). Recovered beads were transferred to 3 mL of Rappaport-Vassiliadis (Remel, St. Louis, MO) broth, incubated at 42°C for 18–20 h, then streaked onto xylose lysine desoxycholate (XLD; Remel, St. Louis, MO) and brilliant green sulfa (Becton Dickinson, Franklin Lakes, NJ) agar plates prior to incubation at 37°C for 18–20 h.

Presumptive *Salmonella* isolates were confirmed using *invA* polymerase chain reaction (Rahn *et al.*, 1992; Nucera *et al.*, 2006). Isolates were subjected to molecular serotyping methods (Herrera-León *et al.*, 2004). Resulting phenotypes were further confirmed by traditional slide agglutination (O typing) and tube agglutination (flagellar H typing) methods, using commercial antisera (Difco, BD Diagnostic Systems, Sparks, MD) following manufacturer's guidelines.

### Antimicrobial susceptibility testing

Susceptibility to 15 antimicrobial agents was determined using broth microdilution (Sensititre CMV1AGNF test plates; TREK Diagnostic Systems, Inc., Cleveland, OH) according to manufacturer's guidelines. Isolates were classified as susceptible, intermediate, or resistant for each agent using breakpoints established by the Clinical and Laboratory Standards Institute (CLSI, 2010). Isolates resistant to two or more classes of antimicrobials, as defined by the National Antimicrobial Resistance Monitoring System (FDA, 2011), were considered multi-drug resistant (MDR).

### Salmonella enumeration

*Salmonella* prevalence values observed in the Fall 2010 sample period for feedlot cattle LNs indicated that enumeration analysis of this population could yield data on levels of *Salmonella* present in contaminated tissues. Conversely, enumeration of cull cattle LNs was not attempted because low prevalence suggested the analysis would yield little data. For enumeration, all feedlot cattle LNs collected in the Summer/Fall 2011 sample period ( $n=618$ ) were processed as described above and immediately following homogenization, 1 mL of tryptic soy broth/LN homogenate was plated onto Petrifilm™ Enterobacteriaceae Count Plates (EB; 3M Microbiology, St. Paul, MN) in quadruplicate, and incubated at 37°C for 22–26 h. Petrifilm plates were then held at 4°C until presumptive culture results were obtained. Colonies were counted and 10–100% (depending on number of colonies present) were streaked to XLD for further confirmation. Colonies indicative of *Salmonella* on XLD plates were counted and the proportion of *Salmonella*-positive colonies were applied to the EB Petrifilm counts, in order to estimate the *Salmonella* count for each enumerated sample. LNs observed to harbor *Salmonella*, but at concentrations below the limit of detection ( $\sim 1$  CFU/g) were designated in the  $<0.1 \log_{10}$  CFU/g category.

### Statistical analysis

*Salmonella* point prevalence was calculated for each set of LNs collected ( $n \approx 76$  per set) in a given processing day ( $n=44$  sample sets). Observed prevalence values were grouped by outcome and the frequency distribution was depicted as a histogram (Fig. 2). Median and mean point prevalence values were determined based on animal type, season, and region, and univariate analyses of results were performed using a one-way analysis of variance and Bonferroni's multiple comparison post-test, the Kruskal-Wallis test for nonparametric data and Dunn's multiple comparison post-test, or the Mann-Whitney  $t$ -test for nonparametric data, as indicated in the footnotes of Table 1. Enumeration data also




FIG. 2. Histogram of *Salmonella* prevalence outcomes for the 44 lymph node (LN) sample sets ( $n \approx 76$  LN per set) collected in this study.

were plotted as total estimated CFU/LN versus LN weight in grams (Fig. 3), and the geometric mean  $\log_{10}$  CFU/g for enumerable samples (114 of 618) was determined. Comparisons of prevalence estimates and construction of data plots were performed using Prism 5.0d, GraphPad Software, Inc. ([www.graphpad.com](http://www.graphpad.com), San Diego, CA) and  $p$  values  $<0.05$  were considered significantly different.

### Results

#### *Salmonella* point prevalence and enumeration

In this cross-sectional study, a total of 3327 subiliac LNs were collected from fed and cull cattle carcasses, in three regions of the United States, over 12 months (Table 1). Examination of the frequency distribution of observed prevalence values showed that the majority of sample sets (74.4%) had few-to-no *Salmonella* positives (Fig. 2). As a consequence, the median point estimate for *Salmonella* contamination was found to be 1.3%. However, some sample sets (25.6%) yielded considerably higher prevalence values. These sample sets skewed the resulting distribution such that arithmetic mean prevalence of *Salmonella* contamination was 7.5%. *Salmonella* point prevalence in feedlot cattle LNs was significantly greater than that observed in cull cattle ( $p=0.0006$ ) and appeared to be affected by region and season (Table 1), with levels significantly higher in region 3 as compared with region 5 ( $p=0.0198$ ), and in Summer/Fall 2011 as compared with Winter/Spring 2011 ( $p=0.0304$ ), for samples collected in region 3. Conversely, *Salmonella* prevalence in cull cattle LNs did not appear to be affected by region or season, as levels in this population tended to be low with a median overall point prevalence of 0.65%, although an exception to that trend was observed in region 5 in the Summer/Fall 2011 sample period (Table 1). It should be noted however, that further investigation revealed that the cull cows contributing to this outlier data point originated in region 3 but were transported to region 5 for harvest.

Enumeration analysis of 618 feedlot cattle LNs collected in Summer/Fall 2011 showed that 23.3% ( $n=144$ ) harbored *Salmonella* and that 18.4% ( $n=114$ ) contained levels detectable with the enumeration methods employed (limit of detection  $\sim 1$  CFU/g). The geometric mean concentration of *Salmonella* for enumerable samples was  $1.75 \log_{10}$  CFU/g; however, as shown in Figure 3, a wide range of values were observed. While the majority of quantifiable nodes (58.8%; 67 of 114) contained *Salmonella* at concentrations ranging from 0.1 to  $1.8 \log_{10}$  CFU/g, 41.2% (47 of 114) carried higher levels, ranging from 1.9 to  $3.8 \log_{10}$  CFU/g, or greater (Fig. 3).

#### *Salmonella* serotypes and antimicrobial susceptibility phenotypes

Twenty-four serotypes were identified, with the majority being either Montevideo (44.0%) or Anatum (24.8%; Table 2). Eighteen serotypes were identified among the 33 positive LNs from cull cattle, whereas 14 serotypes were observed among the 233 positive LNs from feedlot cattle. At least two colonies were serotyped for each positive LN, resulting in the isolation of multiple *Salmonella* serotypes from 3.8% ( $n=10$ ) of positive samples. Multiple serotypes were only observed in feedlot cattle LNs, and for prevalence estimates only one of the two serotypes was counted. Serotype combinations in these samples included Montevideo and Anatum ( $n=7$ ), Montevideo

TABLE 1. *SALMONELLA* PERCENT (%) PREVALENCE, AND STANDARD ERROR (SE) IN SUBILIAC LYMPH NODES (LNs) OF FED AND CULL CATTLE AT HARVEST, BY REGION AND SEASON

Sample period	Cull cattle subiliac lymph nodes <sup>a</sup>				Fed cattle subiliac lymph nodes <sup>a</sup>			Overall by season
	Region 2	Region 3	Region 5	All Cull	Region 3	Region 5	All Fed	
Fall 2010								
Sample sets collected <sup>b</sup>	2	4	2	8	4	2	6	14
Number LNs tested	152	304	152	608	279	152	431	1039
Mean % (SE)	0.65 (0.65)	0.97 (0.62)	0	0.65 (0.35)	29.6 (1.5)	0	19.7 <sup>EF</sup> (6.3)	8.8 (3.7)
Median %	0.65	0.65	0	0 <sup>D</sup>	28.3	0	27.6	0.65
Minimum %	0	0	0	0	27.6	0	0	0
Maximum %	1.3	2.6	0	2.6	34.1	0	34.1	34.1
Winter/Spring 2011								
Sample sets collected <sup>b</sup>	2	4	2	8	4	2	6	14
Number LNs tested	151	305	152	608	305	147	452	1060
Mean % (SE)	3.3 (0.7)	0.32 (0.32)	0	0.98 (0.54)	2.3 (0.34)	0.7 (0.7)	1.8 <sup>E</sup> (0.44)	1.3 (0.37)
Median %	3.3	0	0	0 <sup>D</sup>	2.6	0.7	1.9	1.3
Minimum %	2.6	0	0	0	1.3	0	0	0
Maximum %	4.0	1.3	0	4	2.8	1.4	2.8	4.0
Summer/Fall 2011								
Sample sets collected <sup>b</sup>	2	4	2	8	6	2	8	16
Number LNs tested	152	306	152	610	466	152	618	1228
Mean % (SE)	1.3 (0)	0.65 (0.38)	12.5 (8.6)	3.8 (2.5)	24.7 (8.0)	13.2 (1.4)	21.4 <sup>F</sup> (5.9)	12.0 (3.8)
Median %	1.3	0.65	12.5	1.3 <sup>D</sup>	20.0	13.2	17.7	3.9
Minimum %	1.3	0	3.9	0	1.3	11.8	1.3	0
Maximum %	1.3	1.3	21.1	21.1	47.4	14.5	47.4	47.4
Overall by region or type								Lymph nodes overall
Sample sets collected <sup>b</sup>	6	12	6	24	14	6	20	44
Number LNs tested	455	915	456	1826	1050	451	1501	3327
Mean % (SE)	1.75 (0.56)	0.65 (0.25)	4.2 (3.4)	1.8 (0.88)	19.3 (4.4)	4.6 (2.7)	14.7 (3.5)	7.5 (1.9)
Median %	1.3 <sup>A</sup>	0 <sup>A</sup>	0 <sup>A</sup>	0.65 <sup>G</sup>	20.0 <sup>B</sup>	0.7 <sup>C</sup>	11.8 <sup>H</sup>	1.3
Minimum %	0	0	0	0	1.3	0	0	0
Maximum %	4.0	2.6	21.1	21.1	47.4	14.5	47.4	47.4

<sup>a</sup>Common uppercase superscripts indicate values that are not significantly different ( $p > 0.05$ ).

<sup>b</sup>Approximately 76 LNs collected per set, two sets collected per plant, approximately 1 month apart in each season.

<sup>A</sup>Median cull cattle prevalence by region was not significantly different ( $p = 0.2356$ ) one-way analysis of variance (ANOVA) Kruskal–Wallis nonparametric test; Dunn's multiple comparison post-test.

<sup>B/C</sup>Median fed cattle prevalence by region was significantly different ( $p = 0.0198$ ) Mann–Whitney nonparametric two-tailed  $t$ -test.

<sup>D</sup>Median cull cattle prevalence by season was not significantly different ( $p = 0.3007$ ) one-way ANOVA Kruskal–Wallis nonparametric test; Dunn's multiple comparison post-test.

<sup>E/F</sup>Mean fed cattle prevalence by season was significantly different ( $p = 0.0304$ ) one-way ANOVA and Bonferroni's post-test.

<sup>G/H</sup>Median lymph node prevalence overall for cull and fed cattle was significantly different ( $p = 0.0006$ ) Mann–Whitney nonparametric two-tailed  $t$ -test.


and Infantis ( $n = 1$ ), Montevideo and Muenchen ( $n = 1$ ), and Anatum and Kentucky ( $n = 1$ ).

The majority of *Salmonella* isolates (229 of 266) were susceptible to all antimicrobial agents tested. Multidrug resistance (MDR, defined as resistance to two or more antimicrobial drug classes; [FDA, 2011]) was observed in 8.3% ( $n = 266$ ) of isolates (Table 3). Seventeen isolates (6.4%) exhibited the MDR-AmpC phenotype (co-resistance to at least ampicillin, chloramphenicol, streptomycin, sulfisoxazole, tetracycline, amoxicillin/clavulanic acid, ceftiofur, and ceftriaxone) (Gupta *et al.*, 2003; Kunze *et al.*, 2008). Serotypes

demonstrating MDR-AmpC resistance included Reading ( $n = 13$ ), Newport ( $n = 3$ ), and Typhimurium ( $n = 1$ ), although it should be noted that 11 of the 13 Reading isolated were from a single set of 76 LNs collected from feedlot cattle in the Summer/Fall 2011 sample period, and likely represent a cluster originating from a single lot of cattle at harvest (Table 3).

## Discussion

*Salmonella* are versatile enteric pathogens noted for their ability to invade and survive within host lymphoid tissues



**FIG. 3.** Concentration of *Salmonella* in contaminated lymph nodes (LN) of feedlot cattle at harvest. Enumeration data were collected from 618 LN from carcasses of feedlot cattle in Summer/Fall 2011, using the methods described. Of these, 144 LN were positive for *Salmonella* contamination and 114 LN contained *Salmonella* at enumerable levels. Total estimated *Salmonella* CFU/LN was plotted versus LN weight (grams). Estimated total levels of *Salmonella* ( $\log_{10}$  CFU/LN) are indicated by symbol color intensity (as depicted in the key), with white being the lowest (1.3–1.9  $\log_{10}$  CFU/LN) and dark gray being the highest (4.0–4.8 CFU/LN).

(Stevens *et al.*, 2009). Previous research has shown that cattle peripheral LNs can serve as a vehicle for *Salmonella* contamination, if fat trim containing these nodes is incorporated into ground beef (Arthur *et al.*, 2008; Samuel *et al.*, 1980). In this cross-sectional study, we also observed that *Salmonella* could be recovered from subiliac LNs, but additionally found that point estimates of prevalence in feedlot cattle populations appear to be greater than those in cull cattle populations, and that *Salmonella* harborage may be affected by season and region. While further study is needed to confirm the observed trends, these data nevertheless raise intriguing questions regarding the mechanism of *Salmonella* entry into bovine peripheral LNs, and the factors influencing this phenomenon. Possible explanations for the differences observed include diet effects and animal age, as well as management of cattle prior to harvest and differences in the prevalence of *Salmonella* in cattle environments.

Numerous studies have described a seasonal effect on *Salmonella* prevalence in cattle environments, with prevalence peaking in warmer months (summer and fall), and dipping in colder months (winter and spring) (APHIS, 2001; Barkocy-Gallagher *et al.*, 2003; Edrington *et al.*, 2004). Furthermore, mounting evidence from surveys of cattle feces, hides, and environments suggests a regional variation in *Salmonella* prevalence in North America, where the burden of *Salmonella* broadly increases across a southerly gradient. In a large study that included fecal samples of cattle housed in Canadian feedlots, *Salmonella* was recovered from 0.2% of cattle ready for harvest (Sorensen *et al.*, 2002). In the Midwest, Barkocy-Gallagher *et al.* (Barkocy-Gallagher *et al.*, 2003) reported peak

**TABLE 2.** *SALMONELLA* SEROTYPES RECOVERED FROM SUBILIAC LYMPH NODES (LNs) OF FEEDLOT CATTLE AND CULL COWS

Serotype	All LNs (n=266) overall %	Fed cattle LNs (n=233) relative %	Cull cattle LNs (n=33) relative %
Montevideo	44.0	48.5	12.1
Anatum	24.8	27.5	6.1
Reading	4.9	5.2	3.0
Thompson <sup>a</sup>	3.8	3.9	3.0
Meleagridis	3.0	3.4	
Kentucky	3.0	1.7	12.1
C07 NT	2.3	2.6	
Mbandaka	2.3	1.3	9.1
Muenchen <sup>a</sup>	1.5	1.7	
Bredeney	1.1		9.1
Infantis	1.1	1.3	
Newport	1.1	0.9	3.0
Braenderup	0.8		6.1
Brandenburg	0.8		6.1
Cerro	0.8	0.9	
Dublin	0.8		6.1
Muenster	0.8	0.9	
Panama <sup>a</sup>	0.8		6.1
Saint Paul	0.8	0.4	3.0
Cubana	0.4		3.0
Give	0.4		3.0
Kiambu <sup>a</sup>	0.4		3.0
Typhimurium	0.4		3.0
Uganda	0.4		3.0

<sup>a</sup>Indicates predicted serotype—serotype data incomplete due to the presence of R H antigens that do not react with H antisera. Isolates designated as Thompson<sup>a</sup> may be Thompson (6,7,14: k: 1,5) or Ohio (6,7,14: b: 1,w); isolates designated as Panama<sup>a</sup> may be Panama (1,9,12: 1,v: 1,5) or Javiana (1,9,12: 1,z<sub>28</sub>: 1,5). CO7 NT – Nontypeable *Salmonella* that reacts with O-group 6,7 antisera, but H-antigens are nonreactive.

fecal prevalence of 9.1% during the summer and fall months, while in Texas, *Salmonella* was recovered from 32.0% and 25.5% of fecal samples collected from healthy cattle housed in six feedlots and 22 dairies, respectively (Kunze *et al.*, 2008; Farrow *et al.*, 2009). The observed similarity in seasonal and regional prevalence between *Salmonella* in LNs and in cattle environments suggests the potential for an environmental component to the mechanism of how *Salmonella* gains entry to peripheral nodes. It is known that subiliac LNs receive afferent lymph from the skin of the abdominal wall, pelvis, prepuce, and hind limbs; thus, it is possible that *Salmonella* recovered from subiliac LNs may have entered via a transdermal route, through abrasions or biting insects. This idea has been suggested previously (Samuel *et al.*, 1980), and given that cattle hides are a common reservoir for *Salmonella* (Loneragan and Brashears, 2005; Brichta-Harhay *et al.*, 2008; Kunze *et al.*, 2008), the observed correlation between *Salmonella* prevalence on cattle hides, in cattle environments, and in peripheral LNs is perhaps not surprising. The observed difference in prevalence between feedlot and cull cattle in region 3 was unexpected, however, and may reflect differences in hygiene or mitigation practices (i.e., *Salmonella* vaccine use or differences in pest management) because of a greater perceived risk of *Salmonella* as an animal health issue in dairy cattle populations.

TABLE 3. *SALMONELLA* SEROTYPES AND OBSERVED ANTIMICROBIAL SUSCEPTIBILITY PHENOTYPES RECOVERED FROM SUBILIAC LYMPH NODES OF FEEDLOT CATTLE AND CULL COWS

Animal type	Serotype	Antimicrobial resistance phenotype <sup>a</sup>	Number of lymph nodes
Cull	Mbandaka	S	1
	Kentucky	S	1
	Dublin	Ap, C, K, G, Su	1
	Dublin	Ap, C, K, G, Su, Te	1
	Newport	Am, Ap, F, T, C, S, Su, Te	1
	Reading	Am, Ap, F, T, Ax, C, S, Su, Te	1
	Typhimurium	Am, Ap, F, T, Ax, C, S, Su, Te	1
Fed	Montevideo	Te	11
	Thompson	Te	1
	Montevideo	S	1
	Kentucky	Su, Te	1
	C O7 NT <sup>b</sup>	Su, TS	2
	Reading	Am, Ap, F, T, C, S, Su, Te	11 <sup>c</sup>
	Newport	Am, Ap, F, T, Ax, C, S, Su, Te	2
	Reading	Am, Ap, F, T, Ax, C, S, Su, Te	1

<sup>a</sup>Am, amoxicillin-clavulanic acid; Ap, ampicillin; F, cefoxitin; K, Kanamycin; G, Gentamicin; T, ceftiofur; Ax, ceftriaxone; C, chloramphenicol; S, streptomycin; Su, sulfisoxazole; Te, tetracycline; TS, trimethoprim/sulphamethoxazole.

<sup>b</sup>CO7 NT, nontypeable *Salmonella* that reacts with O-group 6,7 antisera, but H-antigens nonreactive.

<sup>c</sup>Detected as a cluster of isolates originating from one set of 76 lymph nodes and thus may originate from a single lot of cattle at harvest.

Having confirmed that subiliac LNs may be a source of *Salmonella* contamination in ground beef, it was important to examine the serotypes and antimicrobial susceptibility profiles of *Salmonella* occupying this niche, because depending on host status, transmission vehicle, and inoculum level, certain *Salmonella* serotypes appear to be more relevant to causing human disease (Jones *et al.*, 2008). Results showed a diverse set of serotypes were isolated; however, two serotypes—Montevideo (44%) and Anatum (24.8%)—represented the majority of isolates (Table 2). These serotypes are frequently isolated from feces and hides of healthy feedlot cattle, especially in region 3 (Fluckey *et al.*, 2007; Kunze *et al.*, 2008), and as the majority of positive LN samples (79.7%) were collected from feedlot cattle in this region, their predominance is not unexpected. It is noteworthy, however, that Montevideo and Anatum are also the most commonly recovered *Salmonella* serotypes from ground beef in both federal testing programs (FSIS, 2011) and national surveys (Bosilevac *et al.*, 2009). Considering the enumeration data presented here, demonstrating that 33% of contaminated LNs tested (47 of 144) harbored *Salmonella* at levels in the range of 1.9 to >3.8 log<sub>10</sub> CFU/g, it is tempting to suggest that these data identify the mechanism by which the majority of *Salmonella* may be entering ground beef. To our knowledge, this is the first report to document the range in *Salmonella* contamination present in cattle peripheral LNs.

Antimicrobial susceptibility phenotyping showed that the majority of *Salmonella* isolated (86%) were susceptible to all antimicrobials tested; however, MDR *Salmonella* (8.3%) were observed (Table 3). Notably, 15.2% of isolates from cull cattle LNs ( $n=33$ ) were MDR while 7.3% of isolates from fed cattle ( $n=233$ ) had MDR phenotypes. When considering *Salmonella* that are potentially more relevant to human disease, Typhimurium and Newport are two of the leading serotypes isolated in cases of human illness in the United States and have been associated with outbreaks attributed to contaminated ground beef (Gupta *et al.*, 2003). Conversely, serotypes Montevideo and Anatum have been implicated in fewer laboratory-confirmed human salmonellosis cases (CDC, 2011), especially from ground beef sources, and lack medical relevance in comparison. These observations highlight the need for investigation into the virulence factors, or adaptive mechanisms that may be associated with increased human illness among medically relevant serotypes. In this study, serotypes Typhimurium and Newport were observed in 6.1% and 0.9% of cull and fed cattle LNs, respectively (Table 3). Quantification of pathogen load in peripheral LNs containing these medically relevant serotypes will aid in modeling the potential quantitative risk imposed by the addition of contaminated LNs to ground beef.

## Conclusions

The data presented show that subiliac LNs can be a significant source of *Salmonella*, if incorporated in ground beef, and that prevalence appears to be affected by season, region, and animal type. Furthermore, we show that contaminated nodes can carry substantial levels of *Salmonella* (1.9 to >3.8 log<sub>10</sub> CFU/g). As LN harborage protects *Salmonella* from carcass interventions, research is needed to define opportunities for mitigating the risk of *Salmonella* contamination in LNs of apparently healthy cattle.

## Acknowledgments

Funding for this research was provided in part by the Beef Checkoff Program. The authors gratefully acknowledge the assistance of the cooperating abattoirs. We also thank Kim Kucera, Frank Reno, Bruce Jasch, Julie Dyer, Sydney Brodrick, Greg Smith, Rebecca McCarthy, Amy Parks, Max Wolf, Kathleen Fermin, Tanya Jackson, Martha Maradiaga, Ashley Hartzog-Hawkins, and Alexandra Calle for their excellent technical support, and Jan Watts and Jody Gallagher for administrative assistance.

## Disclosure Statement

No competing financial interests exist.

## References

- APHIS. *Salmonella* in United States feedlots. 2001. Available at: [http://www.aphis.usda.gov/animal\\_health/nahms/feedlot/downloads/feedlot99/Feedlot99\\_is\\_Salmonella.pdf](http://www.aphis.usda.gov/animal_health/nahms/feedlot/downloads/feedlot99/Feedlot99_is_Salmonella.pdf), accessed March 14, 2012.
- Arthur TM, Brichta-Harhay DM, Bosilevac JM, et al. Prevalence and characterization of *Salmonella* in bovine lymph nodes potentially destined for use in ground beef. *J Food Prot* 2008;71:1685–1688.
- Barkocy-Gallagher GA, Arthur TM, Rivera-Betancourt M, et al. Seasonal prevalence of shiga toxin-producing *Escherichia coli*, including O157:H7 and non-O157 serotypes, and *Salmonella* in



- commercial beef processing plants. *J Food Prot* 2003;66:1978–1986.
- Barkocy-Gallagher GA, Arthur TM, Siragusa GR, *et al.* Genotypic analyses of *Escherichia coli* O157:H7 and O157 nonmotile isolates recovered from beef cattle and carcasses at processing plants in the midwestern states of the United States. *Appl Environ Microbiol* 2001;67:3810–3818.
- Bosilevac JM, Guerini MN, Kalchayanand N, Koohmaraie M. Prevalence and characterization of *Salmonella* in commercial ground beef in the United States. *Appl Environ Microbiol* 2009;75:1892–1900.
- Brichta-Harhay DM, Arthur TM, Bosilevac JM, *et al.* Microbiological analysis of bovine lymph nodes for the detection of *Salmonella enterica*. *J Food Prot* 2012;75:854–858.
- Brichta-Harhay DM, Guerini MN, Arthur TM, *et al.* *Salmonella* and *Escherichia coli* O157:H7 contamination on hides and carcasses of cull cattle presented for slaughter in the United States: An evaluation of prevalence and bacterial loads by immunomagnetic separation and direct plating methods. *Appl Environ Microbiol* 2008;74:6289–6297.
- [CDC] Centers for Disease Control and Prevention. National *Salmonella* surveillance annual data summary, 2009. Atlanta, GA: Centers for Disease Control and Prevention, 2011.
- [CLSI] Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing; Twentieth Informational Supplement*. CLSI document M100-S20. Wayne, PA: Clinical and Laboratory Standards Institute, 2010.
- Edrington TS, Hume ME, Looper ML, *et al.* Variation in the faecal shedding of *Salmonella* and *E. coli* O157:H7 in lactating dairy cattle and examination of *Salmonella* genotypes using pulsed-field gel electrophoresis. *Lett Appl Microbiol* 2004;38:366–372.
- Farrow RL, Loneragan GH, Edrington TS, *et al.* Quantitative herd-level evaluation of the potential for a commercially available vaccine to effectively control *Salmonella* in dairy cattle. *Annual Meeting of the Conference of Research Workers in Animal Diseases*. Chicago, IL, 2009.
- [FDA] Food and Drug Administration. *National Antimicrobial Resistance Monitoring System—Enteric Bacteria (NARMS): 2009 Executive Report*. Rockville, MD: U.S. Department of Health and Human Services, Food and Drug Administration, 2011.
- Fluckey WM, Loneragan GH, Warner R, Brashears MM. Antimicrobial drug resistance of *Salmonella* and *Escherichia coli* isolates from cattle feces, hides, and carcasses. *J Food Prot* 2007;70:551–556.
- [FSIS] Food Safety and Inspection Service. Serotypes profile of *Salmonella* isolates from meat and poultry products: January 1998 through December 2010. 2011. Available at: [http://www.fsis.usda.gov/PDF/Serotypes\\_Profile\\_Salmonella\\_2010.pdf](http://www.fsis.usda.gov/PDF/Serotypes_Profile_Salmonella_2010.pdf), accessed November 10, 2011.
- FSIS. Microbiological results of raw ground beef products analyzed for *Escherichia coli* O157:H7, summarized by calendar year. 2012. Available at: [http://www.fsis.usda.gov/science/Ground\\_Beef\\_E.Coli\\_Testing\\_Results/index.asp](http://www.fsis.usda.gov/science/Ground_Beef_E.Coli_Testing_Results/index.asp), accessed February 14, 2012. (Online)
- Guo C, Hoekstra RM, Schroeder CM, *et al.* Application of bayesian techniques to model the burden of human salmonellosis attributable to U.S. food commodities at the point of processing: Adaptation of a Danish model. *Foodborne Pathog Dis* 2011;8:509–516.
- Gupta A, Fontana J, Crowe C, *et al.* Emergence of multidrug-resistant *Salmonella enterica* serotype Newport infections resistant to expanded-spectrum cephalosporins in the United States. *J Infect Dis* 2003;188:1707–1716.
- Herrera-León S, McQuiston JR, Usera MA, Fields PI, Garaizar J, Echeita MA. Multiplex PCR for distinguishing the most common phase-1 flagellar antigens of *Salmonella* spp. *J Clin Microbiol* 2004;42:2581–2586.
- Jones TF, Ingram LA, Cieslak PR, *et al.* Salmonellosis outcomes differ substantially by serotype. *J Infect Dis* 2008;198:109–114.
- Kunze DJ, Loneragan GH, Platt TM, *et al.* *Salmonella enterica* burden in harvest-ready cattle populations from the southern high plains of the United States. *Appl Environ Microbiol* 2008;74:345–351.
- Loneragan GH, Brashears MM. Effects of using retention-pond water for dust abatement on performance of feedlot steers and carriage of *Escherichia coli* O157 and *Salmonella* spp. *J Am Vet Med Assoc* 2005;226:1378–1383.
- Loneragan GH, Thomson DU, McCarthy RM, *et al.* *Salmonella* diversity and burden in cows on and culled from dairy farms in the Texas high plains. *Foodborne Pathog Dis* 2012;9:1–7.
- Mead PS, Slutsker L, Dietz V, *et al.* Food-related illness and death in the United States. *Emerg Infect Dis* 1999;5:607–625.
- Nucera DM, Maddox CW, Hoiem-Dalen P, Weigel RM. Comparison of API 20E and *invA* PCR for identification of *Salmonella enterica* isolates from swine production units. *J Clin Microbiol* 2006;44:3388–3390.
- Rahn K, De Grandis SA, Clarke RC, *et al.* Amplification of an *invA* gene sequence of *Salmonella typhimurium* by polymerase chain reaction as a specific method of detection of *Salmonella*. *Mol Cell Probes* 1992;6:271–279.
- Rivera-Betancourt M, Shackelford SD, Arthur TM, *et al.* Prevalence of *Escherichia coli* O157:H7, *Listeria monocytogenes*, and *Salmonella* in two geographically distant commercial beef processing plants in the United States. *J Food Prot* 2004;67:295–302.
- Samuel JL, O’Boyle DA, Mathers WJ, Frost AJ. Isolation of *Salmonella* from mesenteric lymph nodes of healthy cattle at slaughter. *Res Vet Sci* 1980;28:238–241.
- Scallan E, Hoekstra RM, Angulo FJ, *et al.* Foodborne illness acquired in the United States—major pathogens. *Emerg Infect Dis* 2011;17:7–15.
- Sorensen O, Van Donkersgoed J, McFall M, Manninen K, Gensler G, Ollis G. *Salmonella* spp. shedding by Alberta beef cattle and the detection of *Salmonella* spp. in ground beef. *J Food Prot* 2002;65:484–491.
- Stevens MP, Humphrey TJ, Maskell DJ. Molecular insights into farm animal and zoonotic *Salmonella* infections. *Philos Trans R Soc Lond B Biol Sci* 2009;364:2709–2723.

Address correspondence to:  
Dayna M. Brichta-Harhay, PhD  
U.S. Department of Agriculture  
Agricultural Research Service  
U.S. Meat Animal Research Center  
Clay Center, NE 68933

E-mail: Dayna.Harhay@ars.usda.gov

**This article has been cited by:**

1. Michael S. Williams, Eric D. Ebel. 2017. Estimating correlation of prevalence at two locations in the farm-to-table continuum using qualitative test data. *International Journal of Food Microbiology* **245**, 29-37. [[CrossRef](#)]
2. Kristi L. Helke, M.A. McCrackin, Ashley M. Galloway, Ann Z. Poole, Cassandra D. Salgado, Bernadette P. Marriott. 2017. Effects of antimicrobial use in agricultural animals on drug-resistant foodborne salmonellosis in humans: A systematic literature review. *Critical Reviews in Food Science and Nutrition* **57**:3, 472-488. [[CrossRef](#)]
3. Michael S. Williams, Eric D. Ebel, Neal J. Golden. 2017. Using indicator organisms in performance standards for reducing pathogen occurrence on beef carcasses in the United States. *Microbial Risk Analysis* . [[CrossRef](#)]
4. IAN YOUNG, BARBARA J. WILHELM, SARAH CAHILL, REI NAKAGAWA, PATRICIA DESMARCHELIER, ANDRIJANA RAJIĆ. 2016. A Rapid Systematic Review and Meta-Analysis of the Efficacy of Slaughter and Processing Interventions To Control Nontyphoidal Salmonella in Beef and Pork. *Journal of Food Protection* **79**:12, 2196-2210. [[CrossRef](#)]
5. KEELYN E. HANLON, MARKUS F. MILLER, LACEY M. GUILLEN, MINDY M. BRASHEARS. 2016. Salmonella Presence in Mandibular, Mesenteric, and Subiliac Lymph Nodes Collected from Sheep and Goats in the United States. *Journal of Food Protection* **79**:11, 1977-1981. [[CrossRef](#)]
6. Devendra H. Shah, Narayan C. Paul, Willium C. Sischo, Rocio Crespo, Jean Guard. 2016. Population dynamics and antimicrobial resistance of the most prevalent poultry-associated Salmonella serotypes. *Poultry Science* **95**:3, 342-352. [[CrossRef](#)]
7. Cernicchiaro Natalia, Ives Samuel E., Edrington Thomas S., Nagaraja Tiruvoor G., Renter David G.. 2016. Efficacy of a Salmonella Siderophore Receptor Protein Vaccine on Fecal Shedding and Lymph Node Carriage of Salmonella in Commercial Feedlot Cattle. *Foodborne Pathogens and Disease* **13**:9, 517-525. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
8. YICHENG XIE, JEFFREY W. SAVELL, ASHLEY N. ARNOLD, KERRI B. GEHRING, JASON J. GILL, T. MATTHEW TAYLOR. 2016. Prevalence and Characterization of Salmonella enterica and Salmonella Bacteriophages Recovered from Beef Cattle Feedlots in South Texas. *Journal of Food Protection* **79**:8, 1332-1340. [[CrossRef](#)]
9. PIA UNTALAN OLAFSON, TYSON R. BROWN, KIMBERLY H. LOHMEYER, ROGER B. HARVEY, DAVID J. NISBET, GUY H. LONERAGAN, THOMAS S. EDRINGTON. 2016. Assessing Transmission of Salmonella to Bovine Peripheral Lymph Nodes upon Horn Fly Feeding. *Journal of Food Protection* **79**:7, 1135-1142. [[CrossRef](#)]
10. Smith Ashley B., Renter David G., Cernicchiaro Natalia, Shi Xiaorong, Nagaraja Tiruvoor G.. 2016. Prevalence and Quinolone Susceptibilities of Salmonella Isolated from the Feces of Preharvest Cattle Within Feedlots that Used a Fluoroquinolone to Treat Bovine Respiratory Disease. *Foodborne Pathogens and Disease* **13**:6, 303-308. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
11. T. S. EDRINGTON, G. H. LONERAGAN, K. J. GENOVESE, D. L. HANSON, D. J. NISBET. 2016. Salmonella Persistence within the Peripheral Lymph Nodes of Cattle following Experimental Inoculation. *Journal of Food Protection* **79**:6, 1032-1035. [[CrossRef](#)]
12. Rodriguez-Rivera Lorraine D., Cummings Kevin J., Loneragan Guy H., Rankin Shelley C., Hanson Devin L., Leone William M., Edrington Thomas S.. 2016. Salmonella Prevalence and Antimicrobial Susceptibility Among Dairy Farm Environmental Samples Collected in Texas. *Foodborne Pathogens and Disease* **13**:4, 205-211. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
13. D. L. HANSON, G. H. LONERAGAN, T. R. BROWN, D. J. NISBET, M. E. HUME, T. S. EDRINGTON. 2016. Evidence supporting vertical transmission of Salmonella in dairy cattle. *Epidemiology and Infection* **144**:05, 962-967. [[CrossRef](#)]
14. GETAHUN E. AGGA, TERRANCE M. ARTHUR, JOHN W. SCHMIDT, RONG WANG, DAYNA M. BRICHTA-HARHAY. 2016. Diagnostic Accuracy of Rectoanal Mucosal Swab of Feedlot Cattle for Detection and Enumeration of Salmonella enterica. *Journal of Food Protection* **79**:4, 531-537. [[CrossRef](#)]
15. Daniel A. Unruh, Justin J. Kastner, Jacob R. Jenott, Sara E. Gragg. Handling of hamburgers and cooking practices 107-122. [[CrossRef](#)]
16. J. L. Vipham, G. H. Loneragan, L. M. Guillen, J. C. Brooks, B. J. Johnson, A. Pond, N. Pond, M. M. Brashears. 2015. Reduced Burden of Salmonella enterica in Bovine Subiliac Lymph Nodes Associated with Administration of a Direct-fed Microbial. *Zoonoses and Public Health* **62**:8, 599-608. [[CrossRef](#)]
17. T. R. BROWN, T. S. EDRINGTON, G. H. LONERAGAN, D. L. HANSON, K. MALIN, J. J. ISON, D. J. NISBET. 2015. Investigation into Possible Differences in Salmonella Prevalence in the Peripheral Lymph Nodes of Cattle Derived from Distinct Production Systems and of Different Breed Types. *Journal of Food Protection* **78**:11, 2081-2084. [[CrossRef](#)]

18. DARIN R. DOERSCHER, TERRY L. LUTZ, STEPHEN J. WHISENANT, KERRY R. SMITH, CRAIG A. MORRIS, CARL M. SCHROEDER. 2015. Microbiological Testing Results of Boneless and Ground Beef Purchased for the National School Lunch Program, 2011 to 2014. *Journal of Food Protection* **78**:9, 1656-1663. [[CrossRef](#)]
19. A. S. LAUFER, J. GRASS, K. HOLT, J. M. WHICHARD, P. M. GRIFFIN, L. H. GOULD. 2015. Outbreaks of Salmonella infections attributed to beef – United States, 1973–2011. *Epidemiology and Infection* **143**:09, 2003-2013. [[CrossRef](#)]
20. Marie Bugarel, Henk C. den Bakker, Kendra K. Nightingale, Dayna M. Brichta-Harhay, Thomas S. Edrington, Guy H. Loneragan. 2015. Two Draft Genome Sequences of a New Serovar of Salmonella enterica , Serovar Lubbock. *Genome Announcements* **3**:2, e00215-15. [[CrossRef](#)]
21. Doyle Marjorie E.. 2015. Multidrug-Resistant Pathogens in the Food Supply. *Foodborne Pathogens and Disease* **12**:4, 261-279. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
22. Min Li, Sasidhar Malladi, H. Scott Hurd, Timothy J. Goldsmith, Dayna M. Brichta-Harhay, Guy H. Loneragan. 2015. Salmonella spp. in lymph nodes of fed and cull cattle: Relative assessment of risk to ground beef. *Food Control* **50**, 423-434. [[CrossRef](#)]
23. T. R. BROWN, T. S. EDRINGTON, K. J. GENOVESE, G. H. LONERAGAN, D. L. HANSON, D. J. NISBET. 2015. Oral Salmonella Challenge and Subsequent Uptake by the Peripheral Lymph Nodes in Calves. *Journal of Food Protection* **78**:3, 573-578. [[CrossRef](#)]
24. T.L. Wheeler, N. Kalchayanand, J.M. Bosilevac. 2014. Pre- and post-harvest interventions to reduce pathogen contamination in the U.S. beef industry. *Meat Science* **98**:3, 372-382. [[CrossRef](#)]
25. Pia Untalan Olafson, Kimberly H. Lohmeyer, Thomas S. Edrington, Guy H. Loneragan. 2014. Survival and Fate of Salmonella enterica serovar Montevideo in Adult Horn Flies (Diptera: Muscidae). *Journal of Medical Entomology* **51**:5, 993-1001. [[CrossRef](#)]
26. 2014. Scientific Opinion on the public health risks related to the maintenance of the cold chain during storage and transport of meat. Part 1 (meat of domestic ungulates). *EFSA Journal* **12**:3. . [[CrossRef](#)]