

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Roman L. Hruska U.S. Meat Animal Research
Center

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

1-10-2019

Complete Genome Sequence of a *Salmonella enterica* subsp. *enterica* Serovar Fresno Isolate Recovered from a Bovine Lymph Node

Bradd J. Haley
USDA ARS Environmental Microbial and Food Safety Laboratory

Timothy P.L. Smith
USDA ARS U.S. Meat Animal Research Center, tim.smith@ars.usda.gov

Gregory P. Harhay
USDA ARS U.S. Meat Animal Research Center

Guy H. Loneragan
Texas Tech University, guy.loneragan@ttu.edu

Hattie E. Webb
Texas Tech University

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/hruskareports>

Haley, Bradd J.; Smith, Timothy P.L.; Harhay, Gregory P.; Loneragan, Guy H.; Webb, Hattie E.; Bugarel, Marie; Kim, Seon Woo; Van Kessel, Jo Ann S.; and Harhay, Dayna M., "Complete Genome Sequence of a *Salmonella enterica* subsp. *enterica* Serovar Fresno Isolate Recovered from a Bovine Lymph Node" (2019). *Roman L. Hruska U.S. Meat Animal Research Center*. 456.
<https://digitalcommons.unl.edu/hruskareports/456>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Roman L. Hruska U.S. Meat Animal Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Bradd J. Haley, Timothy P.L. Smith, Gregory P. Harhay, Guy H. Loneragan, Hattie E. Webb, Marie Bugarel, Seon Woo Kim, Jo Ann S. Van Kessel, and Dayna M. Harhay



Complete Genome Sequence of a *Salmonella enterica* subsp. *enterica* Serovar Fresno Isolate Recovered from a Bovine Lymph Node

Bradd J. Haley,^a Timothy P. L. Smith,^b Gregory P. Harhay,^b Guy H. Loneragan,^c Hattie E. Webb,^{c*} Marie Bugarel,^c Seon Woo Kim,^a Jo Ann S. Van Kessel,^a Dayna M. Harhay^b

^aUSDA ARS Environmental Microbial and Food Safety Laboratory, Beltsville, Maryland, USA

^bUSDA ARS U.S. Meat Animal Research Center, Clay Center, Nebraska, USA

^cInternational Center for Food Safety Excellence, Department of Animal and Food Sciences, Texas Tech University, Lubbock, Texas, USA

ABSTRACT *Salmonella enterica* serovar Fresno is an infrequently isolated serovar whose ecology and genomic characteristics have not yet been described. To further understand the genomic characteristics of this serovar, we sequenced the complete genome of a single isolate recovered from a bovine lymph node at harvest.

Salmonella enterica serovar Fresno is a rare serovar that has been occasionally isolated from humans, livestock, produce, and the environment in at least 10 countries (1). Although this serovar may be broadly distributed on a geographic scale, it is infrequently recovered in routine surveys of foods, food-producing animals, and human clinical cases, indicating that *S. Fresno* may be found at a higher frequency in other reservoirs, such as unmonitored wildlife, in regions where testing for *S. enterica* is not frequently conducted, or that it is relatively rare, in terms of abundance, compared to other serovars. However, *Salmonella* Fresno has been isolated from humans and a bovine lymph node, suggesting that it can cause disease and may cause invasive infections.

S. enterica isolate USMARC-69835 was recovered from a bovine lymph node collected at harvest in 2012 (2). The isolate was cultured on tryptic soy agar at 37°C for 18 to 20 h and subsequently grown overnight at 37°C in tryptic soy broth. Genomic DNA was extracted from this growth using the Genomic-tip 100/G columns and blood and cell culture DNA midi kits (Qiagen, Valencia, CA). Single-molecule real-time (SMRT) sequencing libraries were constructed using C4/P6 chemistry and sequenced on an RS II instrument (Pacific Biosciences [PB], Menlo Park, CA) following the manufacturer's instructions. A total of 78,613 reads with a mean length of 12,213 bp were sequenced. Sequencing reads were filtered for length and quality (minimum subread length, 500; minimum polymerase read quality, 0.80) using the PB SMRT Portal v.2.3.0.140893 analysis pipeline. *De novo* genome assembly was conducted using the Hierarchical Genome Assembly Process (HGAP) v.3.0 with a minimum seed length of 6,000 (3). A dot plot was constructed for every polished contig using Geneious v.11.1.3 (Biomatters Ltd., New Zealand) (4) to identify the overlapping region, which was subsequently trimmed from the 3' end of the contig. The chromosomal origin of replication was identified using Ori-Finder (5). The trimmed and newly oriented sequences were validated using the PB RS_Resequencing pipeline to map the corresponding reads back to the new reference in order to generate closed, circular, consensus concordance assemblies (3). The final assembly yielded two contigs, consisting of one closed chromosome (GenBank accession number CP032444) and one plasmid (GenBank accession number CP032445). Genome sequencing statistics and genomic features are presented in Table 1.

Citation Haley BJ, Smith TPL, Harhay GP, Loneragan GH, Webb HE, Bugarel M, Kim SW, Van Kessel JAS, Harhay DM. 2019. Complete genome sequence of a *Salmonella enterica* subsp. *enterica* serovar Fresno isolate recovered from a bovine lymph node. *Microbiol Resour Announc* 8:e01338-18. <https://doi.org/10.1128/MRA.01338-18>.

Editor Vincent Bruno, University of Maryland School of Medicine

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply. Address correspondence to Dayna M. Harhay, dayna.harhay@ars.usda.gov.

* Present address: Hattie E. Webb, Weems Design Studio, Inc., and Centers for Disease Control and Prevention, Atlanta, Georgia, USA.

Received 27 September 2018

Accepted 11 December 2018

Published 10 January 2019

TABLE 1 Genome sequencing statistics, characteristics, and accession numbers

Strain or plasmid	Serovar	MLST ST	GenBank (SRA) accession no.	No. of reads (fold coverage [×])	Size (bp)	GC content (%)	Inc type ^b	Resistance gene ^a	SPI ^d	Yr of isolation
USMARC-69835	Fresno	649	CP032444 (SRX4742127)	77,242 (165.8)	4,732,430	52.2		<i>aac(6′)-laa^c</i>	1, 2, 3, 4, 5, 6, 8, 9, 11, 13	2012
pSFR1-USMARC-69835			CP032445	3,392 (353)	89,737	50.2	Incl1			

^a Antibiotic resistance genes were determined using ResFinder 3.1 (9).

^b Plasmid Inc types were determined using PlasmidFinder 2.0 (10).

^c May be cryptic and may not confer aminoglycoside resistance (11).

^d SPI, *Salmonella* pathogenicity islands.

After genome sequencing and assembly, the isolate was identified *in silico* as *Salmonella* Fresno and typed as sequence type 649 (ST649) using SeqSero and MLST 2.0, respectively (6, 7). An Incl1 plasmid with a length of 89,737 bp was assembled among the sequencing reads. In a phylogenetic analysis of 457 *S. enterica* genomes, *S. Fresno* USMARC-69835 clustered with members of subclade A1 (8). Compared to 37 other *S. Fresno* genomes, *S. Fresno* USMARC-69835 was more closely related to isolates from livestock and produce in the United States and more distantly related to three other clades of *S. Fresno* isolated outside North America.

Data availability. The complete genome sequence of *S. enterica* USMARC-69835 has been deposited in GenBank under the accession numbers CP032444 (chromosome) and CP032445 (plasmid). The GenBank submission CP032444 represents a single complete genome. The SRA accession number is SRX4742127.

ACKNOWLEDGMENTS

We are grateful for the technical assistance of Kerry Brader, Sandy Bradley, and Kristen Kuhn.

The mention of a trade name, proprietary product, or specific equipment does not constitute a guarantee or warranty by the USDA and does not imply approval to the exclusion of other products that might be suitable.

D.M.H., T.P.L.S., G.P.H., B.J.H., S.W.K., and J.A.S.V.K. are funded by the Agricultural Research Service of the United States Department of Agriculture.

REFERENCES

1. Alikhan NF, Zhou Z, Sergeant MJ, Achtman M. 2018. A genomic overview of the population structure of *Salmonella*. *PLoS Genet* 14:e1007261. <https://doi.org/10.1371/journal.pgen.1007261>.
2. Webb HE, Harhay DM, Brashears MM, Nightingale K, Arthur TM, Bosilevac JM, Kalchayanand N, Schmidt JW, Wang R, Granier SA, Brown TR, Edrington TS, Shackelford SD, Wheeler TL, Loneragan GH. 2017. *Salmonella* in peripheral lymph nodes of healthy cattle at slaughter. *Front Microbiol* 8:2214. <https://doi.org/10.3389/fmicb.2017.02214>.
3. Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
4. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
5. Gao F, Zhang F. 2008. Ori-Finder: a Web-based system for finding *oriCs* in unannotated bacterial genomes. *BMC Bioinformatics* 9:79. <https://doi.org/10.1186/1471-2105-9-79>.
6. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total genome sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>.
7. Zhang S, Yin Y, Jones MB, Zhang Z, Deatherage Kaiser BL, Dinsmore BA, Fitzgerald C, Fields PI, Deng X. 2015. *Salmonella* serotype determination utilizing high-throughput genome sequencing data. *J Clin Microbiol* 53:1685–1692. <https://doi.org/10.1128/JCM.00323-15>.
8. Timme RE, Pettengill JB, Allard MW, Strain E, Barrangou R, Wehnes C, Van Kessel JS, Karns JS, Musser SM, Brown EW. 2013. Phylogenetic diversity of the enteric pathogen *Salmonella enterica* subsp. *enterica* inferred from genome-wide reference-free SNP characters. *Genome Biol Evol* 5:2109–2123. <https://doi.org/10.1093/gbe/evt159>.
9. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
10. Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller Aarestrup F, Hasman H. 2014. *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 58:3895–3903. <https://doi.org/10.1128/AAC.02412-14>.
11. Magnet S, Courvalin P, Lambert T. 1999. Activation of the cryptic *aac(6′)-Iy* aminoglycoside resistance gene of *Salmonella* by a chromosomal deletion generating a transcriptional fusion. *J Bacteriol* 181:6650–6655.