

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Virology Papers

Virology, Nebraska Center for

---

5-13-2021

## Genome Sequence of a Virulent African Swine Fever Virus Isolated in 2020 from a Domestic Pig in Northern Vietnam

Quang Lam Truong

Thi Lan Nguyen

Thi Hoa Nguyen

Jishu Shi

Hiep Lai Xuan Vu

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/virology>



Part of the [Biological Phenomena](#), [Cell Phenomena](#), and [Immunity Commons](#), [Cell and Developmental Biology Commons](#), [Genetics and Genomics Commons](#), [Infectious Disease Commons](#), [Medical Immunology Commons](#), [Medical Pathology Commons](#), and the [Virology Commons](#)

---

This Article is brought to you for free and open access by the Virology, Nebraska Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Virology Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Quang Lam Truong, Thi Lan Nguyen, Thi Hoa Nguyen, Jishu Shi, Hiep Lai Xuan Vu, Thi Lan Huong Lai, and Van Giap Nguyen

---



# Genome Sequence of a Virulent African Swine Fever Virus Isolated in 2020 from a Domestic Pig in Northern Vietnam

Quang Lam Truong,<sup>a</sup> Thi Lan Nguyen,<sup>a</sup> Thi Hoa Nguyen,<sup>a</sup> Jishu Shi,<sup>b</sup> Hiep Lai Xuan Vu,<sup>c</sup> Thi Lan Huong Lai,<sup>a</sup>  Van Giap Nguyen<sup>a</sup>

<sup>a</sup>Key Laboratory of Veterinary Biotechnology, Faculty of Veterinary Medicine, Vietnam National University of Agriculture, Hanoi, Vietnam

<sup>b</sup>College of Veterinary Medicine, Kansas State University, Manhattan, Kansas, USA

<sup>c</sup>Nebraska Center for Virology, University of Nebraska—Lincoln, Lincoln, Nebraska, USA

**ABSTRACT** This study reports the genome sequence of an isolated African swine fever (ASF) virus (VNUA-ASFV-05L1/HaNam) obtained at the fourth passage on pulmonary alveolar macrophages. The virus was isolated during a typical acute ASF outbreak in pigs in a northern province of Vietnam in 2020.

Spreading to nearly all major swine-producing countries, African swine fever (ASF) is currently considered one of the most important transboundary diseases of pigs (1). The etiological agent, ASF virus (ASFV), belongs to the *Asfarviridae* family, *Asfivirus* genus, and possesses an “open” pangenome (2, 3). Of the 24 ASFV genotypes (I to XXIV) known to date (4), only genotypes I and II have been detected outside Africa (5). The genomic complexity of ASFV is reflected by (i) a very large DNA genome (averaging 186,817 bp), (ii) a variable length from 170,101 bp to 193,886 bp, and (iii) the presence of hundreds of open reading frames, which are classified as core genes or accessory genes (2, 3). As a result, continuous genomic characterization of ASFV genomes is essential for diagnostic, epidemiological, and vaccine development purposes.

In this study, we isolated a virulent strain (VNUA-ASFV-05L1/HaNam) from the spleen of a fattening pig that had succumbed to an acute infection. The sample was taken from a small-scale farm of a farrow-to-finish production system located in a northern province of Vietnam (Ly Nhan District, Ha Nam Province). The spleen homogenate was filtered through a 0.45- $\mu$ m membrane and inoculated in pulmonary alveolar macrophages (PAMs). The infected PAM culture showed specific cytopathic effect at 48 h postinoculation, as characterized by hemadsorption dose (HAD). At the fourth passage, the virus titer was 7.14 log<sub>10</sub> 50% HAD (HAD<sub>50</sub>)/ml at 72 h postinfection. The virus was then semipurified and concentrated using an Amicon Ultra-15 centrifugal filter unit (UFC901024; Millipore). Total DNA was extracted with the QIAamp DNA minikit (51304; Qiagen). The extracted DNA was checked for integrity by gel electrophoresis and measured as at least 50 ng/ $\mu$ l.

The next-generation sequencing was conducted by Apical Scientific Sdn Bhd (Selangor, Malaysia). The library was constructed with the NEBNext Ultra DNA library preparation kit, and the sequencing platform was an Illumina NovaSeq 150PE system. Primer sequences were removed from raw Illumina reads using BBDuk of the BBTtools package (<https://jgi.doe.gov/data-and-tools/bbttools>). Quality control reads were assembled *de novo* using SPAdes (6) and polished using Pilon v1.23 (7), implemented in Unicycler (8). All contigs were subjected to BLASTN searches against the NCBI nucleotide database. Open reading frames were predicted using Prodigal (9) and annotated using Prokka v1.14.6 (10). The single contig (VNUA-ASFV-05L1/HaNam) with a BLASTN similarity to an ASFV was aligned with a number of reference genomes using MAFFT v7.450 (11). The pairwise comparison of average nucleotide identity (ANI) between ASFV genomes was performed by ANI Calculator (12) (<https://www.ezbiocloud.net/tools/ani>). Other tools for genomic visualization and classification of

**Citation** Truong QL, Nguyen TL, Nguyen TH, Shi J, Vu HLX, Lai TLH, Nguyen VG. 2021. Genome sequence of a virulent African swine fever virus isolated in 2020 from a domestic pig in northern Vietnam. *Microbiol Resour Announc* 10:e00193-21. <https://doi.org/10.1128/MRA.00193-21>.

**Editor** Simon Roux, DOE Joint Genome Institute

**Copyright** © 2021 Truong et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Thi Lan Nguyen, [nguyenlan@vnua.edu.vn](mailto:nguyenlan@vnua.edu.vn), or Van Giap Nguyen, [nvgiap@vnua.edu.vn](mailto:nvgiap@vnua.edu.vn).

**Received** 15 March 2021

**Accepted** 17 April 2021

**Published** 13 May 2021

multigene family (MGF) proteins in ASFV were geneCo (13) and MGFC (14), respectively. All bioinformatic tools were run with default parameter settings.

The total output of the sequencer was 10,353,104 reads, and the total number of reads that mapped to the reference genome (GenBank accession number [FR682468](#)) was 461,934. The reads were able to yield a longest single linear contig of 186,237 nucleotides, which was confirmed to be ASFV based on the BLAST result (query coverage, 100%; percent identity, 99.89% to 99.99%). The 186,237-nucleotide genome of VNUA-ASFV-05L1/HaNam had a mean GC content of 38.47% and was predicted to contain 158 protein-coding genes. Compared to several recently reported p72 genotype II strains, the genome-wide sequence identity of VNUA-ASFV-05L1/HaNam was 99.95% with respect to Georgia 2007/1 ([FR682468](#)) and 99.98% with respect to InnerMongolia-AES01 ([MK940252](#)). As a molecular marker distinguishing between closely related ASFVs, the tandem repeat sequence (TATATAGGAA) between the I73R and I329L genes of VNUA-ASFV-05L1/HaNam was classified as variant II (15). In terms of genomic organization, VNUA-ASFV-05L1/HaNam had all 86 core genes reported previously (3). VNUA-ASFV-05L1/HaNam contained all five MGFs (MGF-100, MGF-110, MGF-300, MGF-360, and MGF-505). The VNUA-ASFV-05L1/HaNam MGFs were characterized by a typical organization of (i) location at both the 5' and 3' ends and (ii) distribution along the plus and minus strands of the ASFV genome (14). Additionally, among 31 MGF proteins (14), MGF-110E, MGF-110H, and MGF-110I were not detected in the genome of VNUA-ASFV-05L1/HaNam.

**Data availability.** This genome sequence has been deposited in GenBank under the accession number [MW465755](#). The raw reads are available in the SRA under the accession number [SRX10287451](#).

## ACKNOWLEDGMENT

This work was supported by the Ministry of Science and Technology (project DAKH-02/19-DT.02), Socialist Republic of Vietnam.

## REFERENCES

- Bora M, Bora DP, Manu M, Barman NN, Dutta LJ, Kumar PP, Poovathikkal S, Suresh KP, Nimmanapalli R. 2020. Assessment of risk factors of African swine fever in India: perspectives on future outbreaks and control strategies. *Pathogens* 9:1044. <https://doi.org/10.3390/pathogens9121044>.
- Wang Z, Jia L, Li J, Liu H, Liu D. 2020. Pan-genomic analysis of African swine fever virus. *Virology* 535:662–665. <https://doi.org/10.1007/s12250-019-00173-6>.
- Wang L, Luo Y, Zhao Y, Gao GF, Bi Y, Qiu HJ. 2020. Comparative genomic analysis reveals an 'open' pan-genome of African swine fever virus. *Transbound Emerg Dis* 67:1553–1562. <https://doi.org/10.1111/tbed.13489>.
- Quembo CJ, Jori F, Vosloo W, Heath L. 2018. Genetic characterization of African swine fever virus isolates from soft ticks at the wildlife/domestic interface in Mozambique and identification of a novel genotype. *Transbound Emerg Dis* 65:420–431. <https://doi.org/10.1111/tbed.12700>.
- Arias M, de la Torre A, Dixon L, Gallardo C, Jori F, Laddomada A, Martins C, Parkhouse RM, Revilla Y, Rodriguez FAJ, Sanchez-Vizcaino J. 2017. Approaches and perspectives for development of African swine fever virus vaccines. *Vaccines (Basel)* 5:35. <https://doi.org/10.3390/vaccines5040035>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30:772–780. <https://doi.org/10.1093/molbev/mst010>.
- Yoon S-H, Ha S-M, Lim J, Kwon S, Chun J. 2017. A large-scale evaluation of algorithms to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* 110:1281–1286. <https://doi.org/10.1007/s10482-017-0844-4>.
- Jung J, Kim JJ, Yi G. 2019. geneCo: a visualized comparative genomic method to analyze multiple genome structures. *Bioinformatics* 35:5303–5305. <https://doi.org/10.1093/bioinformatics/btz596>.
- Zhu Z, Chen H, Liu L, Cao Y, Jiang T, Zou Y, Peng Y. 2020. Classification and characterization of multigene family proteins of African swine fever viruses. *Brief Bioinform* bbaa380. <https://doi.org/10.1093/bib/bbaa380>.
- Mazur-Panasiuk N, Walczak M, Juskiewicz M, Woźniakowski G. 2020. The spillover of African swine fever in western Poland revealed its estimated origin on the basis of O174L, K145R, MGF 505-5R and IGR I73R/I329L genomic sequences. *Viruses* 12:1094. <https://doi.org/10.3390/v12101094>.