

12-2015

# Substance Use Disorders, Comorbidity, and Arrest among Indigenous Adolescents\*

Kelley J. Sittner Hartshorn

*University of Nebraska-Lincoln*, [kelly.sittner@okstate.edu](mailto:kelly.sittner@okstate.edu)

Les B. Whitbeck

*University of Nebraska-Lincoln*, [lwhitbeck2@unl.edu](mailto:lwhitbeck2@unl.edu)

Patricia Prentice

*University of Nebraska-Lincoln*, [peprentice@hotmail.com](mailto:peprentice@hotmail.com)

Follow this and additional works at: <http://digitalcommons.unl.edu/sociologyfacpub>



Part of the [Family, Life Course, and Society Commons](#), and the [Social Psychology and Interaction Commons](#)

---

Sittner Hartshorn, Kelley J.; Whitbeck, Les B.; and Prentice, Patricia, "Substance Use Disorders, Comorbidity, and Arrest among Indigenous Adolescents\*" (2015). *Sociology Department, Faculty Publications*. 481.

<http://digitalcommons.unl.edu/sociologyfacpub/481>

This Article is brought to you for free and open access by the Sociology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Sociology Department, Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



# HHS Public Access

Author manuscript

*Crime Delinq.* Author manuscript; available in PMC 2016 December 01.

Published in final edited form as:

*Crime Delinq.* 2015 December ; 61(10): 1311–1332. doi:10.1177/0011128712466372.

## Substance Use Disorders, Comorbidity, and Arrest among Indigenous Adolescents\*

**Kelley J. Sittner Hartshorn**

Research Associate Department of Sociology University of Nebraska-Lincoln 217 Benton Hall  
Lincoln NE 68588-0623 khartshorn2@unl.edu Phone: 402-472-2252 Fax: 402-472-4983

**Les B. Whitbeck**

Bruhn Professor of Sociology Department of Sociology University of Nebraska-Lincoln 739  
Oldfather Hall Lincoln, NE 68588-0324 lwhitbeck2@unl.edu Phone: 402-472-5562 Fax:  
402-472-4983

**Patricia Prentice**

University of Nebraska-Lincoln 214 Benton Hall Lincoln NE 68588-0623 peprentice@hotmail.com  
Phone: 402-472-3632 Fax: 402-472-4983

### Abstract

Indigenous adolescents are overrepresented at multiple stages of the justice system, but we know very little about the role that mental health, particularly substance use disorder, plays in Indigenous pathways to arrest. This study examined the association between substance use disorder, its comorbidity with other disorders, and arrest using a longitudinal sample of Indigenous youth from the Northern Midwest and Canada. Of the 16% of youth who reported at least one arrest at Wave 5, half met criteria for substance abuse/dependence, and slightly more for conduct disorder. Substance abuse/dependence and conduct disorder were each associated with an increased risk of arrest, although co-occurring disorders were not. The reciprocal effects of arrest and mental disorder are discussed.

### Keywords

Substance use disorder; Mental disorder; Criminal arrest; Delinquency; American Indian/Alaska Native

---

On any given day, 1 in 25 American Indians/Alaska Natives (AIAN) aged 18 years or older is under the jurisdiction of the criminal justice system. AIAN youths represent about one percent of the U.S. population yet make up two to three percent of those arrested for offenses such as theft and alcohol violations, and, when arrested, they are 18 times more likely to be committed to adult prisons than are European adolescents. Of all youths who are prosecuted federal courts, 32% are placed in a secure facility; of these 74% are AIAN

---

\*This research was funded by the National Institute on Drug Abuse (DA13580) and the National Institute of Mental Health (MH67281), Les B. Whitbeck, Principal Investigator.

Address correspondence to the first author, University of Nebraska-Lincoln, Department of Sociology, 217 Benton Hall, Lincoln, NE 68588. Phone: 402 472-2252. Fax: 402 472-4983. khartshorn2@unl.edu..

(Perry, 2004). Although Indigenous people make up less than one percent of the population in Minnesota, they are disproportionately stopped, arrested, and incarcerated. Indigenous young people accounted for 15.7% of juveniles in the Minnesota Red Wing Correctional Facility (American Indian Policy Center, 2005). Nationally, the average rate of new commitments of Indigenous youth to adult state prisons is almost twice that (1.84 times) of European American youth. These are enormous ethnic disparities, yet we know very little about AIAN young people in the juvenile justice system and their pathways to arrest and incarceration (Cross, 2008).

Linkages between alcohol abuse, arrests, and incarceration of AIAN people have been around for decades (e.g., Levy, Kunitz, and Everett, 1969; Stewart, 1964), and although the mechanisms at work have been refined, the associations largely still hold (Feldstein, Venner, & May, 2006). In a recent Bureau of Indian Affairs study of alcohol-related crime in five Western states and Alaska, alcohol use was involved in 60% of homicides, 58.6% of child abuse, 52.4% of sexual assaults, and 50% of assaults (Leonardson, 2006). Grobsmith (1989) reported that 100% of the AI inmates in a Great Plains prison told her they had a drug or alcohol dependency problem. Substance abuse disorders were the most common psychiatric disorders among AI adolescents in Northern Plains detention centers (Duclos et al., 1998). Although they don't take arrests into account, there have been several recent studies linking early alcohol and drug use to delinquent behaviors among AI adolescents (Barnes, Welte, and Hoffman 2002; Whitbeck, Chen, Hoyt, and Stubben, 2002; Whitbeck, Hoyt, McMorris, Chen, and Stubben, 2001).

Over time, it has become apparent that the linkages between AIAN alcohol and drug use and arrests are more complicated than early research would make them seem. For example, it is imprudent to generalize about substance abuse among AIAN people because of the great diversity among cultures and even among individual communities within cultures (May, 1996). Recent research that indicates higher abstinence rates among some AI communities than in the general U.S. population challenge old stereotypes (Herman-Stahl, Spencer, & Duncan, 2003; Spicer et al., 2003). But other research indicates that in some communities, AIAN adolescents begin drinking earlier, progress more rapidly to regular use (Beauvais, 1998; Robert W. Blum, Harmon, Harris, Bergeisen, & Resnick, 1992), and have earlier-onset of substance use disorders than their counterparts in the general population (Costello, Farmer, Angold, Burns, & Erkanli, 1997; Whitbeck, Yu, Johnson, Hoyt, & Walls, 2008), behaviors that may place AIAN youth at particular risk for negative consequences.

May and Gossage (2001) have pointed out that the majority of adult alcohol-related crimes among Indigenous people are accounted for by a minority of heavy drinking individuals. This is likely the case with juvenile arrests. Young people whose lives are most affected by substance abuse, those that meet diagnostic criteria for SUD, are a minority of heavy users among many abstaining or casual users of alcohol and drugs.

In this paper we investigate the relationships between substance use disorders (SUD) and comorbid SUD and other psychiatric disorders and likelihood of arrest among Indigenous (American Indian and Canadian First Nations) adolescents. We focus on adolescents who meet criteria for SUD in that they represent the habitual users, those most at risk for alcohol

and drug related contacts with the criminal justice system, and those in need of treatment upon entry into the criminal justice system. We use multivariate analyses of longitudinal data to investigate the effects of meeting criteria for SUD, other psychiatric disorders and comorbidity when controlling for other known risk factors to test the relative strength of SUD on risk of entering the criminal justice system.

## Substance Use Disorders, Comorbidity, and Juvenile Arrests

Mental disorders are increasingly common among adolescents in the juvenile justice system. Indeed, the criminal justice system has become the primary referral for mentally disordered adolescents (Grisso, 2004). Of particular concern is the disproportionate number of adolescents with substance use disorders in the juvenile justice system. For example, in their study of adolescents in a Cook County juvenile facility, Teplin and colleagues (2002) found that approximately half of male and female detainees had a substance use disorder.

Psychiatric disorders rarely occur alone during adolescence (Kessler & Walters, 1998) and this is particularly true of SUDs (Hser et al., 2001). A history of conduct disorder doubles the risk of SUD (Sung, Erkanli, Angold, & Costello, 2004), and adolescents with externalizing disorders (e.g., conduct disorder, oppositional defiant disorder, attention deficit hyperactivity disorder (ADHD)) and co-occurring SUDs typically have worse developmental outcomes and poorer treatment prognoses (Clark & Scheid, 2001). Depressive disorders also often co-occur with adolescent-onset SUD (Clark, Kirisci, & Tarter, 1998), a combination that is associated with multiple negative outcomes (Bukstein, Brent, & Kaminer, 1989) and poor treatment outcomes (Grella, Hser, Joshi, & Rounds-Bryant, 2001). There is evidence among the general population that SUDs, co-occurring with any other psychiatric disorder, increase the likelihood that a person will be arrested (Swartz and Lurigio 2007). Additionally, there is some evidence that comorbidity of substance use and psychiatric disorders is more common among the AI children than majority children (Costello et al., 1997; Federman, Costello, Angold, Farmer, & Erkanli, 1997) perhaps putting them at particular risk for negative outcomes such as arrest.

## Theory and Hypotheses

Early onset SUD may be an important factor in understanding the etiology and consequences of serious delinquent and criminal behavior among Indigenous adolescents and, along with comorbid psychiatric disorders, may differentiate those who persist in antisocial behaviors from those who engage in more normative adolescent-limited types of delinquency (Moffitt, 1993). SUDs, particularly in concert with externalizing disorders, increase the likelihood of antisocial behaviors by the disinhibiting effects of substance use on judgment and risk taking (Fromme, Katz, & D'Amico, 1997) and in the activities surrounding procurement and use of illegal substances (Arseneault, Moffitt, Caspi, Taylor, & Silva, 2000). Early onset SUD is strongly associated with later criminality (Loeber, Burke, Lahey, Winters, & Zera, 2000). Cooccurring externalizing disorders and SUDs amplify disruptive, hyperactive and oppositional problem behaviors that are “requisite for sustained and serious violent offending” (Tolan & Gorman-Smith, 1998, p. 75).

Alcohol dependence develops over two to four years (Martin, Langenbucher, Kaczynski, & Chung, 1996) and marijuana dependence may take up to two years or more to emerge (Rosenberg & Anthony, 2001). Such long-standing patterns of use and abuse are difficult to change. This is especially true for adolescents with co-occurring psychiatric disorders who are both less likely to complete treatment programs and more likely to relapse after treatment (Brown, Gleghorn, Schuckit, Myers, & Mott, 1996). SUDs and co-occurring disorders together contribute to the momentum of maladaptive behaviors, establishing trajectories that may become more and more difficult to interrupt. The concept of “cumulative continuity” suggests delinquent adolescents may become ensnared in the consequences of their own behaviors, resulting in lifelong effects (Caspi, Bem, & Elder, 1989; Caspi, Elder, & Bem, 1987). We believe that SUDs and comorbid disorders play an important part of this process.

Based on prior research, we investigate three hypotheses. First, controlling for other risk factors, SUD would increase the odds of arrest among Indigenous adolescents. Second, externalizing disorders (e.g., conduct disorder and ADHD) would increase the likelihood of arrest when controlling for other risk factors. Third, given research findings that comorbidity is common among arrested adolescents, we hypothesize that comorbid disorders, specifically SUD plus other disorders, would increase the likelihood of arrest over and above the direct effects of any single disorder (Dolamanta, Risser, Roberts, & Risser, 2003; Neighbors, Kempton, & Forehand, 1992; Swartz & Lurigio, 2007).

## Methods

### Procedure

These data were collected as part of an ongoing longitudinal study designed in partnership with four U.S. reservations and four Canadian First Nations reserves and a university-based research team. The reservations share a common cultural tradition and language with minor regional variations in dialects. The sample represents one of the most populous Indigenous cultures in the United States and Canada. Prior to the application's funding, the research team was invited to work with the reservations/reserves, and tribal resolutions were obtained. As part of the partnership agreement, the researchers promised that culture and the names of the participating reservations/reserves would be kept anonymous in published reports.

On each participating reservation, an advisory board representing all reservation/reserve districts was appointed by the tribal council. The advisory boards are responsible for handling difficult personnel problems, advising on questionnaire development and reviewing reports for respectful writing and confidentiality. All participating staff on the reservations/reserves were approved by the advisory boards and were either tribal members or, in a few cases, nonmembers who were spouses of tribal members. To ensure quality of data collection, all the interviewers underwent special training at each wave of the study for conducting pencil and paper interviews and computer-assisted personal interviewing for the diagnostic measures. The training included in-class practice, feedback sessions, and homework practice assignments.

Each of the participating reservations and reserves provided a list of families of tribally enrolled children aged 10–12 years who lived on or proximate (within 50 miles) to the reservation or reserve. We attempted to contact all families with a target child within the specified age range. Families were recruited with a personal visit by an Indigenous interviewer at which time the project was explained to them. They were then presented with a traditional gift and invited to participate. The recruitment method was approved by advisory boards on each of the reservation and reserves. If the family members agreed to be interviewed, each family member (the target child and at least one parent or caretaker) received \$40 for their time when the interviews were completed. The recruitment procedure resulted in an overall response rate of 79.4%.

## Sample

These analyses use data from the first five waves of the study. Overall this study had very little attrition, with retention rates ranging from 88.1% to 94.6% across the five waves. We used two strategies to address missing data. For the descriptive analyses (bivariate correlations (Table 1), and prevalence of mental disorder (Table 2) and comorbidity (Table 3) among arrested youth) listwise deletion was used for any cases with missing values on any of the included variables. This resulted in a final sample size of 603 adolescents for the descriptive analyses. For the multivariate analyses predicting arrest at Wave 5 (Table 4), missing data was addressed by multiple imputation in Stata using a freeware command called “ice” (Royston 2005; Royston, Carlin, and White 2009). Multiple imputation allows parameter estimates to be pooled from a series of imputed data sets, using information from all of the covariates included in an analysis, which provides improved parameter estimates and more accurate standard errors than in single imputation methods (Acock, 2005). The use of multiple imputation resulted in a final sample size of 689 adolescents.

## Measures

Not all variables used in the present paper are drawn from the same waves. Although there is considerable consistency of measures used across waves, not all of variables are available at every wave. Namely, only diagnostic information was collected in Wave 4, and thus only the mental disorder variables are drawn from that wave. The prior arrest variable comes from the first three waves for that same reason. In addition, our dependent variable is the only variable drawn from Wave 5.

The dependent variable, *arrest*, was assessed in Wave 5 of the study. The adolescents were asked, “Have you been arrested within the past twelve months?” Those who responded ‘yes’ were coded as 1, ‘no’ was coded as 0. Approximately fifteen percent of the youth responded that they had been arrested at least once in Wave 5.

To assess youth psychiatric disorders, the Diagnostic Interview Schedule for Children-Revised Version II (DISC-R) *conduct disorder*, *attention deficit hyperactivity disorder*, *substance use disorders* (alcohol abuse, alcohol dependence, marijuana abuse, and marijuana dependence), and *major depressive episode* modules were used. The DISC-R is a highly regarded, structured interview intended for use with trained interviewers who are not clinicians. Computer algorithms were used to determine whether each behavior was present

at clinically significant frequency/severity levels. We use the combined youth and parent/caretaker reports, which tend to be more reliable than child reports alone (Schwab-Stone et al., 1996; Shaffer et al., 1996).

Three family risk factors (*parental depression*, parental substance abuse/dependence, and parental rejection) were included in the study. The University of Michigan Composite International Diagnostic Interview (UM-CIDI), based on DSM-III-R criteria, was used to assess parental 12-month major depressive episode (*parental depression*) and parental lifetime substance abuse/dependence (*parental SUD*). Computer algorithms were used to determine whether each behavior was present at clinically significant frequency/severity levels. A single indicator of *parental substance use* was constructed from the individual substance-related diagnoses (alcohol abuse, alcohol dependence, marijuana abuse, marijuana dependence). Diagnoses were coded as “1” if at least one caretaker met diagnostic criteria and “0” if neither caretaker met criteria.

*Parental rejection* is a mean indicator of adolescent-reported rejection over the first three waves of the study. At each wave, respondents were asked how often someone in their family really trusts them, blames them when they don't deserve it, really cares for them, blames them for things that others have done, spends time doing things they enjoy, are unhappy with things they do, and gives them attention for things they do. Positively worded questions were reverse-coded. A high value indicates higher levels of rejection. The measure had acceptable internal reliability, with a Cronbach's alpha of 0.78.

*Prior arrest* was constructed from responses to the question, “Have you ever been arrested?” (Wave 1) and “Have you been arrested within the past twelve months?” (Waves 2–3). Only diagnostic questions were asked at Wave 4, thus no youth report of arrest at that wave is included. Youth who responded that they had been arrested were coded as 1. A single index of prior arrest was constructed by summing the responses, with a range of 0 to 3.

Four control variables were included in the analyses, all drawn from Wave 3. *Gender* is dichotomized and coded as Male = 1. *Age* is a continuous measure of adolescent age in years. *Location* refers to the proximity to towns or other communities. There are two locations identified for this sample. A rural location refers to those reservations/reserves within somewhat close proximity to other towns, and remote refers to those far removed from other communities. The variable is coded so that Remote=1. *Per capita family income* is measured by asking parents/caretakers to indicate their overall household incomes in the past year. Responses were divided by the number of people living within the household, which was then divided by 1,000 to set the metric of this measure in thousands of dollars.

### Analytic Strategy

The analytic strategy of this paper is twofold. First, we used bivariate analyses to explore the associations between arrest and SUD, conduct disorder, ADHD, major depressive episode, and comorbid SUD with these disorders. Second, we employed stepwise logistic regression to test the hypotheses of the associations between mental disorder and arrest.

## Results

### Mental Disorders/Comorbidity and Arrest

Bivariate correlations, means, and standard deviations are presented in Table 1. Arrest was associated with gender, age, per capita family income, parent rejection, prior arrest, substance abuse/dependence (SUD), and conduct disorder (CD). There was no significant association between arrest and major depressive episode (MDE), or arrest and attention deficit hyperactivity disorder (ADHD).

Prior arrest was associated with SUD and CD; it was also associated with meeting criteria for MDE. Parental depression was associated with parent SUD, parent rejection, adolescent SUD, and adolescent CD. Parental rejection was associated with all adolescent mental disorders. Parental SUD was not associated with any adolescent mental disorder.

Table 2 presents the prevalence of the disorders and comorbidity among arrested and nonarrested Indigenous adolescents. Almost two-thirds of youth who were not arrested in Wave 5 (64%) had no mental disorder, compared to only one-third of arrested youth (32.9%). Adolescents without a disorder had the lowest odds of arrest (O.R.=.30). Substance use disorder, although fairly prevalent overall, was much more common among arrested youth: 50.2% of youth arrested in Wave 5 met criteria for SUD, compared to only 21% of nonarrested youth. One-half of adolescents who were arrested met criteria for CD (53.2%). There were no significant differences between arrested and nonarrested youth who met criteria for ADHD or MDE.

In addition to the individual disorders, arrested youth were much more likely to have comorbid disorders (Table 3). Of the 47 adolescents with SUD who were arrested in Wave 5, 74.5% had co-occurring CD, 8.5% had co-occurring MDE, and 12.8% had co-occurring ADHD. But perhaps more importantly, arrested adolescents with either MDE or ADHD had very high rates of co-occurring SUD. Of the five arrested adolescents with MDE, four of them also had comorbid SUD. Of the seven arrested adolescents with ADHD, six of them had comorbid SUD. This suggests that SUD is very common among adolescents with MDE or ADHD who have been arrested, although neither MDE nor ADHD are as prevalent among those with SUD.

Substance use disorder comorbidity was less common but still pervasive among adolescents who were not arrested at Wave 5. Of the 107 adolescents with SUD, 43.9% also had CD, 15% had MDE, and 9.3% had ADHD. Slightly more adolescents meeting criteria for the other disorders had co-occurring SUD. Over half of the 86 adolescents with CD, more than half (54.7%) also had SUD, and fewer adolescents with either MDE (43.2%) or ADHD (25%) had comorbid SUD.

### Logistic Regression Models: Substance Use Disorders, Comorbidity and Arrest

Arrest risk factors and individual mental disorders affecting the probability of arrest were explored in a series of logistic regression models (Table 4). The four control variables (gender, age, per capita family income, and remote location), family risk factors (parental depression, parental SUD, and parental rejection), and prior arrest were entered

simultaneously in the first model. Of the control variables, only income and gender were significant. Males ( $b=1.12$ ,  $\text{Exp}(b)=3.06$ ) had significantly higher odds of arrest than females, but age of adolescent was unrelated to arrest. Of the family risk factors for arrest, only perceived parental rejection was associated with arrest. A one unit increase in parental rejection above the mean was associated with an almost four-fold increase in risk of arrest ( $b=1.35$ ,  $\text{Exp}(b)=3.86$ ). As expected, prior arrest was associated with later arrest. For each additional prior arrest above no arrest, the odds of subsequent arrest increased by 88% ( $b=.63$ ,  $\text{Exp}(b)=1.88$ ).

To address the first two hypotheses, that SUD and externalizing disorders, respectively, would increase the odds of arrest among Indigenous adolescents, each individual mental disorder was added separately into the logistic regression models. Similar to the bivariate analyses, SUD and conduct disorder were each significantly associated with arrest. The odds of arrest for adolescents who met criteria for SUD were three times higher than for adolescents without the disorder (Model 2,  $b=1.13$ ,  $\text{Exp}(b)=3.10$ ). Adolescents with CD had odds of arrest almost four times higher than adolescents without CD (Model 3,  $b=1.38$ ,  $\text{Exp}(b)=3.97$ ). As in the bivariate analyses, ADHD (Model 4) and MDE (Model 5) were not associated with later arrest.

Because both SUD and CD were so strongly associated with arrest, and because there is considerable comorbidity of the two disorders, we tested a model with both disorders entered simultaneously (Model 6). SUD and CD remain significant when both are included, although the odds ratios are smaller than those in Models 2 and 3 with the disorders entered separately. Adolescents with SUD have odds of arrest that are twice as high as adolescents without SUD, controlling for CD ( $b=.69$ ,  $\text{Exp}(b)=1.99$ ). Adolescents with CD have odds of arrest that are more than three times higher than adolescents without CD, controlling for SUD ( $b=1.13$ ,  $\text{Exp}(b)=3.10$ ). To address the third hypothesis, that comorbidity of SUD with other disorders will increase the odds of arrest, we tested an interaction effect of SUD and CD (Model 7). The association was not significant, indicating that there is not a joint effect of the two disorders beyond the additive combinations of their individual effects. Because neither ADHD nor MDE had significant direct effects, we did not test any interactions with those disorders and SUD.

## Discussion/Conclusions

Consistent with prior research (Duclos et al., 1998; Hartney, 2008), Indigenous adolescents with SUD or CD reported high arrest rates. More than half of adolescents who were arrested met criteria for either SUD or CD. Comorbidity, particularly for adolescents who met criteria for SUD and CD, was very common among arrested adolescents. And, importantly, adolescents who met criteria for ADHD and MDE were much more likely to have been arrested if they had concurrent SUD. Indeed, of the seven young people with ADHD who were arrested, six also met criteria for SUD. Similarly, of the five depressed adolescents who were arrested, four met criteria for SUD. Neither of these two disorders, however, increased the odds of arrest.

SUDs also were linked to arrests in the multivariate analyses. Meeting criteria for SUD in Wave 4 increased the likelihood of arrest in Wave 5 three times and meeting criteria for CD at Wave 4 increased the odds of Wave 5 arrest four times. We were surprised to find that the interaction between SUD and CD was not statistically significant. That is, both disorders exerted independent effects but did not have a joint effect on the likelihood of arrest. The lack of association between ADHD and arrest has been documented in other studies (Hirschfeld, Maschi, White, Traub, & Loeber, 2006; Satterfield & Schell, 1997). There is evidence that it is the hyperactivity/impulsivity subtype which is associated with delinquency rather than the attention deficit subtype (Babinski, Hartsough, & Lambert, 1999; Carroll et al., 2006; Lahey & Loeber, 1997), a distinction we could not make in the current study. Similar to ADHD, MDE was not related to Wave 5 arrests regardless of gender of adolescent. This could be due to the small number of adolescents who met criteria for MDE.

There were several interesting findings regarding control variables and family risk factors. One, adolescent age was unrelated to odds of arrest. Two, parental mental health was not associated with adolescent arrest. Of little surprise were the findings regarding gender, parental rejection, and prior arrest. Boys, adolescents who felt rejected by their caretakers prior to Wave 5, and adolescents who had been arrested in previous waves were more likely to be arrested in Wave 5. This is congruent with gendered arrest patterns (Federal Bureau of Investigation, 2009), a considerable body of research that shows family effects on delinquent behaviors (Blumstein, Cohen, Roth, & Visher, 1986; Chatterji & Markowitz, 2001; Farrington, Barnes, & Lambert, 1996; Rowe & Farrington, 1997; Rutter, Giller, & Hagell, 1998), as well as the stability of delinquent behavior across time (Nagin & Farrington, 1992; Nagin & Paternoster, 1991; Sampson & Laub, 2005).

### Limitations

There are several important limitations to this study that deserve consideration. First, it was based upon a single Indigenous culture and may not be generalizable to other Indigenous populations. However, because of the great diversity among cultures and the near impossibility of obtaining a random, representative sample across all of the more than 500 cultures, research such as this must proceed culture by culture and rely on cumulative information as it emerges. Second, the study adolescents lived either on or close to rural reservations, so these results may not reflect the experiences of urban Indigenous young people even if they share the same cultural background. Third, there is always the concern that European diagnostic measures do not take into account cultural differences. Our diagnostic measures, however, are the same as those used in major epidemiological studies of AI people. At this point, these are the best measures available. Fourth, our reliance on self-report data invites the criticism of potential social desirability bias, but we believe the most likely bias would be underreporting arrests and problem behaviors, making our findings conservative.

### Conclusions

Our findings extend previous research linking substance abuse and arrests among Indigenous adolescents through the use of diagnostic measures to identify those most at risk

for serious, perhaps life-long consequences. Only about 5% of adolescents in the general population are likely to be life course persistent antisocial (Moffitt, 1993). Although it is difficult to accurately project the continuation of problem behaviors, we know that these are the young people most at risk for persistent problems with alcohol and drugs and involvement with the criminal justice system into adulthood (Loeber, Stouthamer-Loeber, & White, 1999; Moffitt, Caspi, Harrington, & Milne, 2002). By first arrest, adolescents with this combination of life experiences may already be on a trajectory of negative, self-reinforcing events that will become harder and harder to moderate as time goes by (Sampson & Laub, 2005; Stewart, Simons, Conger, & Scaramella, 2002). They are more likely to be re-arrested (Gendreau, Little, & Goggin, 1996; Wierson & Forehand, 1995), they are less likely to complete treatment programs (Clark & Scheid, 2001; Kaminer, Tarter, Bukstein, & Kabene, 1992), and if treatment is mandated, they are more likely to relapse (Brown et al., 1996; Grella et al., 2001).

The key may be early identification and treatment of related SUD and mental disorders either prior to or at first contact with the criminal justice system. Diversionary programs that require psychological and substance abuse evaluations along with interventions and/or family-based interventions (Mmari, Robert Wm. Blum, & Teufel-Stone, 2010; Pullmann et al., 2006) may reduce early and repeated arrests. Moreover, diversionary programs may be more cost-effective than incarceration for substance abusing offender (Mauser, Van Stelle, & Moberg, 1994). But first arrest already may be too late in a cumulative process. Very early assessments to identify and intervene with high risk children may reduce the arrest rate in the long-term. As we have pointed out, most substance related crimes are committed by a very small group of heavy users (May & Gossage, 2001). These heavy users were likely early onset substance abusers. Genuinely high risk adolescents are a small group of early onset substance abusers almost certainly with pre-existing behavioral problems associated with conduct disorder, oppositional defiant disorder, and/or ADHD. Early identification and intervention with these relatively few children and their families would be extremely cost effective both in monetary and human terms. Without such interventions, we may relegating some of these children to an adolescence ensnared in a sequence of disruptive behaviors, poor academic performance, affiliation with delinquent peers, substance use and eventually involvement in the criminal justice system. If arrest is the point of first intervention, we already may be too late.

## Biographies

**Kelley J. Sittner Hartshorn, Ph. D.** is a post-doctoral research associate for a longitudinal study of Indigenous adolescents in the Department of Sociology at the University of Nebraska-Lincoln. She received her B.S. in Education in 2000, her M.A. in Sociology in 2007, and her Ph.D. in Sociology in 2011, each from the University of Nebraska-Lincoln. Her research focuses on antisocial behavior, mental health, and substance use among adolescents.

**Les B. Whitbeck, Ph.D.** is the John G. Bruhn Professor of Sociology at the University of Nebraska-Lincoln. He received his B.A. in philosophy from Western Washington State University in 1969, a M.A. in Religion from Earlham School of Religion in 1973, a Masters

in Social Work from Indiana University-Purdue University at Indianapolis in 1975 and his Ph.D. in sociology from Washington State University in 1986. He was principal investigator for the Midwest Longitudinal Study of Homeless Adolescents and is currently heading a NIH-funded program of research that focuses on homeless women and children. He is principal investigator for an eight-year diagnostic study of American Indian children aged 10–12 years. Dr. Whitbeck has authored or co-authored two books on homeless adolescents and over 100 refereed journal articles. He received the 2007 Community, Culture, and Prevention Science Award from the Society for Prevention Research.

**Patricia Prentice** is an on-site coordinator for the longitudinal diagnostic study of American Indian children. She is also a grant writer for her reservation.

## References

- Acock AC. Working with missing values. *Journal of Marriage and Family*. 2005; 67:1012–1028.
- American Indian Policy Center. Alcohol/drugs and crime. Searching for Justice. 2005. Retrieved from <http://www.airpi.org/research/SearchingforJustice/sjfindcauses/sjfalco.htm>
- Arseneault L, Moffitt TE, Caspi A, Taylor PJ, Silva PA. Mental disorders and violence in a total birth cohort: Results from the Dunedin Study. *Archives of General Psychiatry*. 2000; 57(10):979–986. [PubMed: 11015816]
- Babinski LM, Hartsough CS, Lambert NM. Childhood conduct problems, hyperactivity-impulsivity, and inattention as predictors of adult criminal activity. *The Journal of Child Psychology and Psychiatry and Allied Disciplines*. 1999; 40(03):347–355.
- Barnes GM, Welte JW, Hoffman JH. Relationship of alcohol use to delinquency and illicit drug use in adolescents: Gender, age, and racial/ethnic differences. *Journal of Drug Issues*. 2002; 32(1):153–178.
- Beauvais F. American Indians and alcohol. *Alcohol Health and Research World*. 1998; 22(4):253–259. [PubMed: 15706751]
- Blum RW, Harmon B, Harris L, Bergeisen L, Resnick MD. American Indian--Alaska Native youth health. *Journal of the American Medical Association*. 1992; 267(12):1637–1644. [PubMed: 1542173]
- Blumstein, A.; Cohen, J.; Roth, JA.; Visher, CA. *Criminal careers and “career criminals”*. National Academy Press; Washington, DC: 1986.
- Brown SA, Gleghorn A, Schuckit MA, Myers MG, Mott MA. Conduct disorder among adolescent alcohol and drug abusers. *Journal of Studies on Alcohol*. 1996; 57(3):314–324. [PubMed: 8709590]
- Bukstein O, Brent D, Kaminer Y. Comorbidity of substance abuse and other psychiatric disorders in adolescents. *American Journal of Psychiatry*. 1989; 146(9):1131–1141. [PubMed: 2669535]
- Carroll A, Hemingway F, Bower J, Ashman A, Houghton S, Durkin K. Impulsivity in juvenile delinquency: Differences among early-onset, late-onset, and non-offenders. *Journal of Youth and Adolescence*. 2006; 35(4):517–527.
- Caspi A, Bem DJ, Elder GH. Continuities and consequences of interactional styles across the life course. *Journal of Personality*. 1989; 57(2):375–406. [PubMed: 2769561]
- Caspi A, Elder GH, Bem DJ. Moving against the world: Life-course patterns of explosive children. *Developmental Psychology*. 1987; 23(2):308–313.
- Chatterji P, Markowitz S. The impact of maternal alcohol and illicit drug use on children's behavior problems: Evidence from the children of the National Longitudinal Survey of Youth. *Journal of Health Economics*. 2001; 20(5):703–731. [PubMed: 11558645]
- Clark, DB.; Scheid, J. Comorbid mental disorders in adolescents with substance use disorders. In: Hubbard, JR.; Martin, PR., editors. *Substance abuse in the mentally and physically disabled*. Marcel Dekker, Inc; New York: 2001. p. 133-168.

- Clark DB, Kirisci L, Tarter RE. Adolescent versus adult onset and the development of substance use disorders in males. *Drug and Alcohol Dependence*. 1998; 49(2):115–121. [PubMed: 9543648]
- Costello EJ, Farmer EMZ, Angold A, Burns BJ, Erkanli A. Psychiatric disorders among American Indian and white youth in Appalachia: The Great Smoky Mountains study. *American Journal of Public Health*. 1997; 87(5):827. [PubMed: 9184514]
- Cross, TL. Native Americans and juvenile justice: A hidden tragedy (No. 11356). Poverty & Race Research Action Council; Washington, DC: 2008.
- Dolamanta DD, Risser WL, Roberts RE, Risser JMH. Prevalence of depression and other psychiatric disorders among incarcerated youths. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2003; 42(4):477–484. [PubMed: 12649635]
- Duclos, Christine Wilson; Beals, Janette; Novins, Douglas K.; Martin, Cheryl; Jewett, Chastity S.; Manson, Spero M. Prevalence of common psychiatric disorders among American Indian adolescent detainees. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1998; 37:866–873. [PubMed: 9695449]
- Farrington DP, Barnes GC, Lambert S. The concentration of offending in families. *Legal and Criminological Psychology*. 1996; (1):47–63.
- Federal Bureau of Investigation. Persons arrested: Crime in the United States 2008. U.S. Department of Justice. 2009. Retrieved March 8, 2009, from <http://www.fbi.gov/ucr/cius2008/arrests/index.html>
- Federman EB, Costello EJ, Angold A, Farmer EMZ, Erkanli A. Development of substance use and psychiatric comorbidity in an epidemiologic study of white and American Indian young adolescents the Great Smoky Mountains Study. *Drug and Alcohol Dependence*. 1997; 44(2–3):69–78. [PubMed: 9088778]
- Feldstein SW, Venner KL, May PA. American Indian/Alaska Native alcohol-related incarceration and treatment. *American Indian and Alaska Native Mental Health Research*. 2006; 13(3):1–22. [PubMed: 17602406]
- Fromme K, Katz E, D'Amico E. Effects of alcohol intoxication on the perceived consequences of risk taking. *Experimental and Clinical Psychopharmacology*. 1997; 5(1):14–23. [PubMed: 9234035]
- Gendreau P, Little T, Goggin C. A meta-analysis of the predictors of adult offender recidivism: What works. *Criminology*. 1996; 34:575.
- Grella CE, Hser Y, Joshi V, Rounds-Bryant J. Drug treatment outcomes for adolescents with comorbid mental and substance use disorders. *The Journal of Nervous and Mental Disease*. 2001; 189(6):384–392. [PubMed: 11434639]
- Grisso, T. *Double jeopardy: Adolescent offenders with mental disorders*. University of Chicago Press; Chicago: 2004.
- Grobsmith ES. Relationship between substance abuse and crime among Native American inmates in the Nebraska Department of Corrections. *Human Organization*. 1989; 48(4):285.
- Hartney, C. Native American youth and the juvenile justice system. National Council on Crime and Delinquency. 2008. Retrieved February 7, 2009, from [http://www.nccd-crc.org/nccd/pubs/2008\\_Focus\\_NativeAmerican.pdf](http://www.nccd-crc.org/nccd/pubs/2008_Focus_NativeAmerican.pdf)
- Herman-Stahl M, Spencer DL, Duncan JE. The implications of cultural orientation for substance use among American Indians. *American Indian and Alaska Native Mental Health Research*. 2003; 11(1):46–66. [PubMed: 12955631]
- Hirschfeld P, Maschi T, White HR, Traub LG, Loeber R. Mental health and juvenile arrests: Criminality, criminalization, or compassion? *Criminology*. 2006; 44(3):593–630.
- Hser Y, Grella CE, Hubbard RL, Hsieh S, Fletcher BW, Brown BS, Anglin MD. An evaluation of drug treatments for adolescents in 4 US cities. *Archives of General Psychiatry*. 2001; 58(7):689–695. [PubMed: 11448377]
- Kaminer Y, Tarter RE, Bukstein OG, Kabene M. Comparison between treatment completers and noncompleters among dually diagnosed substance-abusing adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1992; 31(6):1046–1049. [PubMed: 1429403]
- Kessler RC, Walters EE. Epidemiology of DSM-III-R major depression and minor depression among adolescents and young adults in the National Comorbidity Survey. *Depression and Anxiety*. 1998; 7(1):3–14. [PubMed: 9592628]

- Lahey, BB.; Loeber, R. Attention-deficit/hyperactivity disorder, oppositional defiant disorder, conduct disorder, and adult antisocial behavior: A life span perspective. In: Stoff, DM.; Breiling, J.; Maser, JD., editors. *Handbook of antisocial behavior*. John Wiley & Sons; Hoboken, NJ: 1997. p. 51-59.
- Leonardson, GR. Native American crime in the northwest: 2004–2005 -- BIA information from Alaska, Montana, Wyoming, Idaho, Oregon, and Washington. Montana Board of Crime Control; Dillon, MT: 2006.
- Levy JE, Kunitz SJ, Everett M. Navajo criminal homicide. *Southwestern Journal of Anthropology*. 1969; 25(2):124–152.
- Loeber R, Burke JD, Lahey BB, Winters A, Zera M. Oppositional defiant and conduct disorder: A review of the past 10 years, part I. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2000; 39(12):1468–1484. [PubMed: 11128323]
- Loeber R, Stouthamer-Loeber M, White HR. Developmental aspects of delinquency and internalizing problems and their association with persistent juvenile substance use between ages 7 and 18. *Journal of Clinical Child Psychology*. 1999; 28(3):322. [PubMed: 10446681]
- Martin CS, Langenbucher JW, Kaczynski NA, Chung T. Staging in the onset of DSM-IV alcohol symptoms in adolescents: Survival/hazard analyses. *Journal of Studies on Alcohol*. 1996; 57(5): 549–558. [PubMed: 8858553]
- Mauser E, Van Stelle KR, Moberg DP. The economic impact of diverting substance-abusing offenders into treatment. *Crime & Delinquency*. 1994; 40(4):568–588.
- May, PA. Overview of alcohol abuse epidemiology for American Indian populations. In: Sandefur, GD.; Rindfuss, RR.; Cohen, B., editors. *Changing numbers, changing needs: American Indian demography and public health*. National Academy Press; Washington, DC: 1996. p. 235-256.
- May PA, Gossage J. New data on the epidemiology of adult drinking and substance use among American Indians of the northern states: Male and female data on prevalence, patterns, and consequences. *American Indian and Alaska Native Mental Health Research*. 2001; 10(2):1–26. [PubMed: 11698981]
- Mmari KN, Blum RW, Teufel-Stone N. What increases risk and protection for delinquent behaviors among American Indian youth?: Findings from three tribal communities. *Youth & Society*. 2010; 41(3):382–413.
- Moffitt, Terrie E. Adolescence-limited and life-course-persistent antisocial behavior: A developmental taxonomy. *Psychological Review*. 1993; 100:674–701. [PubMed: 8255953]
- Moffitt TE, Caspi A, Harrington H, Milne BJ. Males on the life-course-persistent and adolescence-limited antisocial pathways: Follow-up at age 26 years. *Development and Psychopathology*. 2002; 14(01):179–207. [PubMed: 11893092]
- Nagin DS, Farrington DP. The stability of criminal potential from childhood to adulthood. *Criminology*. 1992; 30(2):235–260.
- Nagin DS, Paternoster R. On the relationship of past to future participation in delinquency. *Criminology*. 1991; 29(2):163–189.
- Neighbors B, Kempton T, Forehand R. Co-occurrence of substance abuse with conduct, anxiety, and depression disorders in juvenile delinquents. *Addictive Behaviors*. 1992; 17(4):379–386. [PubMed: 1502971]
- Perry, SW. American Indians and crime: A BJS statistical profile, 1992–2002 (No. NCJ 203097). U.S. Department of Justice, Bureau of Justice Statistics; 2004. p. 1-56.
- Pullmann MD, Kerbs J, Koroloff N, Veach-white E, Gaylor R, Sieler D. Juvenile offenders with mental health needs: Reducing recidivism using wraparound. *Crime & Delinquency*. 2006; 52(3): 375–397.
- Reasons, CE.; Kuykendall, JL., editors. *Race, crime, and justice*. Goodyear Publishing Co; Pacific Palisades, CA: 1972.
- Rosenberg MF, Anthony JC. Early clinical manifestations of cannabis dependence in a community sample. *Drug and Alcohol Dependence*. 2001; 64(2):123–131. [PubMed: 11543982]
- Rowe DC, Farrington DP. The familial transmission of criminal convictions. *Criminology*. 1997; 35:177–201.
- Royston P. Multiple imputation of missing values: Update. *The Stata Journal*. 2005; 5(2):188–201.

- Royston P, Carlin JB, White IR. Multiple imputation of missing values: New features for mim. *The Stata Journal*. 2009; 9(2):252–264.
- Rutter, M.; Giller, H.; Hagell, A. *Antisocial behavior by young people: A major new review*. Cambridge University Press; Cambridge, UK: 1998.
- Sampson, R.J.; Laub, J.H. A life-course theory of cumulative disadvantage and the stability of delinquency. In: Thornberry, TP., editor. *Developmental theories of crime and delinquency*. 2nd ed.. Transaction Publishers; New Brunswick, NJ: 2005. p. 133-161.
- Satterfield JH, Schell A. A prospective study of hyperactive boys with conduct problems and normal boys: Adolescent and adult criminality. *Journal of the American Academy of Child and Adolescent Psychiatry*. 1997; 36(12):1726–1735. [PubMed: 9401334]
- Schwab-Stone ME, Shaffer D, Dulcan MK, Jensen PS, Fisher P, Bird HR, Goodman SH, et al. Criterion Validity of the NIMH Diagnostic Interview Schedule for Children Version 2.3 (DISC-2.3). *Journal of the American Academy of Child & Adolescent Psychiatry*. 1996; 35(7): 878–888. doi:97/00004583-199607000-00013. [PubMed: 8768347]
- Shaffer D, Fisher P, Dulcan MK, Davies M, Piacentini J, Schwab-Stone ME, Lahey BB, et al. The NIMH Diagnostic Interview Schedule for Children Version 2.3 (DISC-2.3): Description, Acceptability, Prevalence Rates, and Performance in the MECA Study. *Journal of the American Academy of Child & Adolescent Psychiatry*. 1996; 35(7):865–877. doi: 97/00004583-199607000-00012. [PubMed: 8768346]
- Spicer P, Beals J, Croy CD, Mitchell CM, Novins DK, Moore L, Manson SM. The prevalence of DSM-III-R alcohol dependence in two American Indian populations. *Alcoholism: Clinical and Experimental Research*. 2003; 27(11):1785–1797.
- Stewart EA, Simons RL, Conger RD, Scaramella LV. Beyond the interactional relationship between delinquency and parenting practices: The contribution of legal sanctions. *Journal of Research in Crime and Delinquency*. 2002; 39(1):36–59.
- Stewart O. Questions regarding American Indian criminality. *Human Organization*. 1964; 23(1):61–66.
- Sung M, Erkanli A, Angold A, Costello EJ. Effects of age at first substance use and psychiatric comorbidity on the development of substance use disorders. *Drug and Alcohol Dependence*. 2004; 75(3):287–299. [PubMed: 15283950]
- Swartz JA, Lurigio AJ. Serious mental illness and arrest: The generalized mediating effect of substance use. *Crime & Delinquency*. 2007; 53(4):581–604.
- Teplin LA, Abram KM, McClelland GM, Dulcan MK, Mericle AA. Psychiatric disorders in youth in juvenile detention. *Archives of General Psychiatry*. 2002; 59(12):1133–1143. [PubMed: 12470130]
- Tolan, PH.; Gorman-Smith, D. Development of serious and violent offending careers. In: Loeber, R.; Farrington, D., editors. *Serious & violent juvenile offenders: Risk factors and successful interventions*. Sage Publications, Inc; Thousand Oaks, CA: 1998. p. 68-85.
- Whitbeck LB, Hoyt DR, Chen X, Stubben JD. Predictors of gang involvement among American Indian adolescents. *Journal of Gang Research*. 2002; 10(1):11–26.
- Whitbeck LB, Hoyt DR, McMorris BJ, Chen X, Stubben JD. Perceived discrimination and early substance abuse among American Indian children. *Journal of Health and Social Behavior*. 2001; 42(4):405–424. [PubMed: 11831140]
- Whitbeck LB, Yu M, Johnson KD, Hoyt DR, Walls ML. Diagnostic prevalence rates from early to mid-adolescence among Indigenous Adolescents: First results from a longitudinal study. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2008; 47(8):890–900. [PubMed: 18596558]
- Wierson M, Forehand R. Predicting recidivism in juvenile delinquents: The role of mental health diagnoses and the qualification of conclusions by race. *Behaviour Research and Therapy*. 1995; 33(1):63–7. [PubMed: 7872939]

**Table 1**

Mean, Standard Deviation, and Correlation for All Study Variables (N=603)

	1	2	3	4	5	6	7	8	9	10	11	12	13
1 Arrest	1.00												
2 Male=1	<b>.18</b> ***	1.00											
3 Age	<b>.09</b> *	.03	1.00										
4 Per capita Family Income	<b>-.15</b> ***	.03	.04	1.00									
5 Remote Location	-.01	-.02	-.06	-.04	1.00								
6 Parent Depression	.07	-.04	.02	-.04	-.06	1.00							
7 Parent SUD	.01	.07	.06	.00	<b>-.23</b> ***	<b>.20</b> ***	1.00						
8 Parent Rejection	<b>.12</b> **	.01	.07	-.01	-.06	<b>.08</b> *	.07	1.00					
9 Prior Arrest	<b>.19</b> ***	-.02	<b>.14</b> ***	<b>-.14</b> ***	-.07	.04	.04	<b>.10</b> *	1.00				
10 Substance Abuse/Dependence	<b>.24</b> ***	-.08	<b>.23</b> ***	<b>-.12</b> **	-.02	<b>.09</b> *	.07	<b>.18</b> ***	<b>.27</b> ***	1.00			
11 Conduct Disorder	<b>.32</b> ***	.02	.06	<b>-.18</b> ***	-.05	<b>.11</b> **	.03	<b>.23</b> ***	<b>.20</b> ***	<b>.43</b> ***	1.00		
12 ADHD	-.01	.03	-.01	-.03	-.07	.02	.07	<b>.12</b> **	.05	.06	<b>.14</b> ***	1.00	
13 Depression	-.03	<b>-.10</b> *	.03	.00	-.07	.07	.06	<b>.09</b> *	<b>.13</b> **	<b>.14</b> ***	<b>.13</b> **	<b>.19</b> ***	1.00
<i>M</i>	.16	.49	13.03	\$5,897	.09	.21	.74	.67	.16	.26	.23	.07	.08
<i>SD</i>	.36	.50	.77	\$4,343	.29	.41	.44	.22	.46	.44	.42	.25	.27
$\alpha$													.78

Note: Age, Per capita family income, Parent rejection, and Prior arrest are mean centered.

\* p<0.05,

\*\* p<0.01,

\*\*\* p<0.001

**Table 2**

Prevalence of Mental Disorders among Indigenous Adolescents (n=603)

	Odds Ratios for arrest <sup>a</sup>		Nonarrested youth (n=509)		Arrested youth (n=94)	
			n	%	n	%
No disorder	.30***		326	64.0	31	32.9***
Mental disorder <sup>b</sup>						
Substance Abuse/Dependence	3.84***		107	21.0	47	50.2***
Conduct Disorder	4.98***		86	16.9	50	53.2***
ADHD	.85		40	7.9	7	7.5
Depression	.83		37	7.3	5	5.3

<sup>a</sup>The Odds Ratios are calculated controlling for Gender, Age, Per capita family income and Remote location.

<sup>b</sup>The diagnostic categories of individual mental disorders are not mutually exclusive.

**Table 3**

Prevalence of Substance Abuse/Dependence and Comorbidity among Indigenous Adolescents (n=603)

	Nonarrested youth		Arrested youth	
	n	%	n	%
Substance Abuse/Dependence	107	21.0	47	50.2
with Conduct Disorder	47	43.9	35	74.5
with ADHD	10	9.3	6	12.8
with Depression	16	15.0	4	8.5
Conduct Disorder	86	16.9	50	53.2
with Substance Abuse/Dependence	47	54.7	35	70.0
ADHD	40	7.9	7	7.5
with Substance Abuse/Dependence	10	25.0	6	85.7
Depression	37	7.3	5	5.3
with Substance Abuse/Dependence	16	43.2	4	80.0

Note: The diagnostic categories are not mutually exclusive.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 4**

Logistic Regression Models Predicting Arrest (n=689)

	Model 1		Model 2		Model 3		Model 4		Model 5		Model 6		Model 7	
	b	Exp(b)												
Male=1	1.12 (.24)	<b>3.06</b> ***	1.24 (.25)	<b>3.46</b> ***	1.13 (.26)	<b>3.10</b> ***	1.12 (.24)	<b>3.06</b> ***	1.10 (.24)	<b>3.00</b> ***	1.20 (.26)	<b>3.32</b> ***	1.20 (.14)	<b>3.32</b> ***
Age	.25 (.15)	1.28	.10 (.16)	1.11	.22 (.16)	1.25	.26 (.15)	1.30	.26 (.15)	1.30	.14 (.16)	1.15	.14 (.16)	1.15
Family Income	-.12 (.04)	<b>.89</b> **	-.11 (.04)	<b>.90</b> **	-.09 (.04)	<b>.91</b> **	-.12 (.04)	<b>.89</b> **	-.12 (.04)	<b>.89</b> ***	-.09 (.03)	<b>.91</b> *	-.09 (.04)	<b>.91</b> *
Remote location=1	.09 (.43)	1.09	-.01 (.44)	.99	.12 (.45)	1.13	.07 (.43)	1.07	.08 (.43)	1.08	.05 (.46)	1.05	.05 (.45)	1.05
Parent Depression	.44 (.28)	1.55	.39 (.28)	1.48	.32 (.29)	1.38	.44 (.28)	1.55	.44 (.28)	1.55	.30 (.29)	1.35	.30 (.29)	1.35
Parent SUD	-.15 (.29)	.86	-.22 (.30)	.80	-.11 (.30)	.90	-.13 (.29)	.88	-.13 (.29)	.88	-.16 (.30)	.85	-.16 (.30)	.85
Parent Rejection	1.35 (.55)	<b>3.86</b> *	1.04 (.55)	2.83	.75 (.56)	2.12	1.44 (.56)	<b>4.22</b> *	1.41 (.55)	<b>4.10</b> *	.68 (.57)	1.97	.68 (.57)	<b>1.97</b> *
Prior Arrest	.63 (.20)	<b>1.88</b> **	.41 (.22)	1.51	.48 (.22)	<b>1.62</b> *	.65 (.20)	<b>1.92</b> **	.67 (.20)	<b>1.95</b> **	.37 (.23)	1.45	.37 (.23)	1.45
SUD			1.13 (.28)	<b>3.10</b> ***							.69 (.31)	<b>1.99</b> *	.75 (.41)	2.12
CD					1.38 (.26)	<b>3.97</b> ***					1.13 (.29)	<b>3.10</b> ***	1.18 (.37)	<b>3.25</b> **
ADHD							-.54 (.46)	.58						
MDE														
SUD × CD														
Constant	-2.66	.07 ***	-2.97	.05 ***	-3.05	.05 ***	-2.64	.07 ***	-2.63	.07 ***	-3.17	.04 ***	-3.19	.04 ***

\* p<0.05,  
 \*\* p<0.01,  
 \*\*\* p<0.001