

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

US Fish & Wildlife Publications

US Fish & Wildlife Service

4-12-2011

TRUMPETER SWAN SURVEY of the ROCKY MOUNTAIN POPULATION WINTER 2011

Dave Olson

Division of Migratory Bird Management

Follow this and additional works at: <https://digitalcommons.unl.edu/usfwspubs>

Olson, Dave, "TRUMPETER SWAN SURVEY of the ROCKY MOUNTAIN POPULATION WINTER 2011" (2011).
US Fish & Wildlife Publications. 487.

<https://digitalcommons.unl.edu/usfwspubs/487>


This Article is brought to you for free and open access by the US Fish & Wildlife Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in US Fish & Wildlife Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



U.S. Fish and Wildlife Service

Trumpeter Swan Survey of the Rocky Mountain Population

Winter 2011



Acknowledgements

Personnel who conducted the survey are listed in Appendix C. The survey is a collaborative effort among Red Rock Lakes NWR, Migratory Birds and State Programs -- Mountain-Prairie Region of the U.S. Fish and Wildlife Service, Southeast Idaho National Wildlife Refuge Complex, National Elk Refuge, Harriman State Park, Idaho Department of Fish and Game, Grand Teton National Park, Yellowstone National Park, Wyoming Game and Fish Department, Ruby Lake NWR, Malheur NWR, and the Shoshone-Bannock Tribes. Additionally, R. Cavallaro, M. Wackenhut, B. Waterbury, D. Christopherson, K. Cameron, R. Holman, R. Lonsinger, T. Ferguson, and C. Mitchell assisted with counts in Idaho. M. Fisher, J. Mackay, S. Patla, N. Cadwell, D. Smith, M. St. Louis, and K. Cutting provided information and narratives used to develop this document; conclusions are attributable only to the author.

TRUMPETER SWAN SURVEY
of the
ROCKY MOUNTAIN POPULATION

WINTER 2011

U.S. Fish and Wildlife Service
Migratory Birds and State Programs
Mountain-Prairie Region
Lakewood, Colorado


April 12, 2011


Prepared by:


Dave Olson, Assistant Migratory Game Bird Coordinator
Division of Migratory Bird Management

Approved:


Will Meeks, Acting Regional Chief
Division of Migratory Bird Management


Matt Hogan, Assistant Regional Director
Migratory Birds and State Programs


Richard A. Coleman, Assistant Regional Director
National Wildlife Refuge System

Abstract.— Observers counted 5,712 swans (white birds and cygnets) in the Rocky Mountain Population of trumpeter swans during February 2011, which was a 33% increase from the 4,290 counted during winter 2010. The number of white birds (4,384) and the number of cygnets (1,328) increased 22% and 92% respectively from last year. The cygnet count this year was the highest since 1972. In the tri-state area, the number of total swans increased for Montana (25%), Idaho (27%) and Wyoming (55%) from counts in 2010. The number of birds wintering in areas near restoration flocks increased from last year and was the highest count since 1995. The numbers of birds at Ruby Lake National Wildlife Refuge (NWR) (37), Malheur NWR (10) and Summer Lake Wildlife Management Area (WMA) (78) increased 42%, 11% and 200% respectively from 2010. Reservoir levels in early February were lower than during winter 2010 but 6% above the long term average. Temperatures in the tri-state area during winter 2010-11 were about average or just below average, and precipitation was about 90% of average.

The Rocky Mountain Population (RMP) of trumpeter swans (*Cygnus buccinator*) consists of birds that nest primarily from western Canada southward to Nevada and Wyoming (Fig. 1). The population is comprised of several flocks that nest in different portions of the overall range. The RMP/Canadian Flocks consist of birds that summer primarily in southeastern Yukon Territory, southwestern Northwest Territories, northeastern British Columbia, Alberta, and western Saskatchewan. The RMP/Tri-state Area Flocks summer in areas at the juncture of the boundaries of Montana, Wyoming, and Idaho (hereafter termed the tri-state area) and nearby areas (Fig. 2). The Canadian and Tri-state Area flocks winter sympatrically primarily in the tri-state area. In addition, efforts have been made to establish several RMP restoration flocks, such as those at Ruby Lake National Wildlife Refuge (NWR) in Nevada (i.e., Nevada flock) and those at Malheur NWR and Summer Lake Wildlife Management Area (WMA) and vicinity (i.e., Oregon flock), by translocating adult swans and cygnets from other portions of the RMP. These birds tend to winter in areas near those where they nest. These terms for the various groups of swans are consistent with the Pacific Flyway Management Plan for the RMP of Trumpeter Swans (Subcommittee on the Rocky Mountain Population of Trumpeter Swans 2008).

Although counts of swans wintering in the tri-state area have been conducted since at least the 1950s (Banko 1960), many early efforts were not well-coordinated and were variable. In an attempt to better coordinate the survey, in 1972 the U.S. Fish and Wildlife Service (Service) began the annual Mid-winter Trumpeter Swan Survey in the tri-state region. During the next decade, the area surveyed increased substantially, and by 1981 it was believed all known occupied wintering sites were included (Gale et al. 1988). Recent attempts to expand the wintering range of RMP trumpeter swans have resulted in the inclusion of yet more areas to the survey. Also, some areas may not be surveyed in a particular year due to weather or resource limitations (e.g., staff, money). Such survey modifications make individual counts from year-to-year less comparable, but the data are sufficient to reasonably depict trends in abundance.

The Mid-winter Trumpeter Swan Survey is conducted annually in late January or early February. The survey is conducted cooperatively by several administrative entities and is intended to provide




Fig. 1. Approximate ranges of trumpeter swans during summer (from Moser 2006).




Fig. 2. Map showing the 'core' tri-state area of southeast Idaho, southwest Montana, and northwest Wyoming (provided by the Greater Yellowstone Coalition, Bozeman, Montana).

an annual assessment of the number of RMP trumpeter swans. Only data from 1972 to present, the time frame during which the Service has coordinated the survey, were analyzed for this report.

METHODS

The survey generally is conducted within a relatively short time frame (i.e., 1 week) to reduce the possibility of counting swans more than once due to movements of birds among areas. Aerial cruise surveys generally are used to count numbers of swans in the tri-state area, Nevada, Malheur NWR, and in the Summer Lake WMA and vicinity; ground surveys are used to count the number of swans in isolated pockets of habitat not covered by aerial surveys. During aerial surveys, data are collected by observers seated in a single-engine, fixed-winged aircraft. Flying altitude varies with changes in terrain and surface winds, but generally averages 30-60 m above ground level, and flight speed is between 135-155 kph. One to two observers and the pilot count white (i.e., adults and subadults) and gray (i.e., cygnets) swans in known or suspected habitats. Counts are not adjusted for birds present but not seen by aerial crews, and have an unknown and unmeasured sampling variance associated with them. Ground surveys are used to verify species composition of some swan flocks, because trumpeter and tundra (*C. columbianus*) swans are difficult to differentiate during aerial surveys. Efforts are made to identify and exclude tundra swans from the survey counts. Generally about 30 hours of flight time and additional time spent conducting surveys on the ground are required to complete the survey.

Annual estimates of abundance for Canadian Flocks are determined by subtracting the count of the RMP/U.S. Breeding Segment in the previous fall (e.g., U.S. Fish and Wildlife Service 2008a) from the Mid-winter count. For the estimate of the size of the Canadian Flocks to be accurate, several conditions must be met. First, all swans must be correctly identified to species. Second, the Mid-winter count and the fall count of swans in the RMP/U.S. Breeding Segment must be accurate. Additionally, we must assume that mortality in the RMP/U.S. Breeding Segment between the time of the fall and winter surveys is negligible. Because of problems inherent in surveying biological populations, these conditions probably are seldom met. Thus, this methodology for estimating the size of the RMP/Canadian Flocks likely leads to somewhat biased estimates of the composition of the RMP. However, the historic counts using this methodology generally track those from the quinquennial trumpeter swan survey (U.S. Fish and Wildlife Service 2006), suggesting it produces a useful index to annual abundance for the Canadian Flocks.

To assess production for the RMP, we calculated the percentage of annual total swan counts that were cygnets. However, surveys in Nevada and Oregon did not separate counts into white birds and cygnets until 1992. Therefore, to allow an assessment over a longer time frame with data that are relatively comparable from year-to-year, we used only information from birds counted in the tri-state region. This subset contained a large majority (range = 87%-98%, mean = 95%) of the total RMP counts during 1972-2011. Counts used for analyses in this report are provided in Appendix A.

RESULTS AND DISCUSSION

The 2011 Mid-winter survey was conducted between 31 January and 11 February. Aerial surveys in the tri-state area were completed by 11 February and required about 28 hours to complete. Across most of the areas weather conditions were very favorable for flying with clear skies and calm winds. Ice conditions were consistent over the survey area with low temperatures freezing many of the open water bodies and concentrating swans in the remaining ice-free waters.

Precipitation during December to February was 75% to 125% of average throughout much of the tri-state area (Joint Agricultural Weather Facility 2011). Water levels at 5 reservoirs (American Falls, Island Park, Jackson Lake, Palisades, and Minidoka Dam/Lake Walcott) cumulatively were at 66% of storage capacity on 1 February (data from U.S. Bureau of Reclamation 2011a), 14% below the level of last year but, 6% above the 1972-2010 average (Fig. 3). Together, these reservoirs comprise about 97% of the water-storage capacity for reservoirs listed in the Snake River Basin in eastern Idaho and extreme western Wyoming (U.S. Bureau of Reclamation 2011b). Snowpack as of 1 February throughout much of the tri-state area was generally 100-120% of normal, about 50-100% of normal in south-central Oregon, and about 90-110% of normal in northeastern Nevada (U.S. Department of Agriculture 2011).

The average streamflow on the Henrys Fork near Island Park Reservoir, Idaho during 15 January to 15 February 2011 was 370 cfs, which is near the 1972-2010 average for that recording station (U.S. Bureau of Reclamation 2011a) (Fig. 3). The December-February temperatures were near or just below average throughout much of the primary wintering area in the tri-state region (Fig. 4).

Historical Trends

Methods used to estimate trends in rates of change in RMP abundance were detailed in a previous report (U.S. Fish and Wildlife Service 2003), and will not be reiterated here. Briefly, however, we used least-squares regression on log-transformed counts to assess rates of change in counts of swans over time. Counts from the current Mid-winter survey (2011) were compared to results from 1972-2010, a practice used in Service survey reports for other waterfowl (e.g., Zimpfer et al. 2010, U.S. Fish and Wildlife Service 2010b). Because Nevada and Oregon did not separate total counts of swans into white birds and cygnets prior to 1992 (see above), analyses to assess trends for white birds and cygnets used only counts from the tri-state area.

The counts for total swans of the RMP suggested an increase ($P < 0.01$) of 5.5% per year during 1972-2010 (Table 1, Fig. 5). The number of white birds and cygnets counted in the tri-state region both increased ($P < 0.01$) at 5.7% per year. Counts of birds in Montana (white birds + cygnets) increased slightly (+1.2% per year, $P = 0.06$), whereas average annual rates of growth for birds




Fig. 3. Water storage for 5 reservoirs (see text) in the tri-state region on 1 February, and average streamflow between 15 January and 15 February on the Henrys Fork, 1972-2011.




Fig. 4. Departure of average temperature from normal ($^{\circ}\text{F}$) during December 2010 to February 2011 (Joint Agricultural Weather Facility 2011).

Table 1. Counts of trumpeter swans of the Rocky Mountain Population during winter, 1972-2011.

Year	<u>Tri-state area</u>			<u>Oregon and Nevada^a</u>			<u>Total RMP</u>		
	White	Cygnets	Total	White	Cygnets	Total	White	Cygnets ^b	Total
1972	c	c	616			91			707
1973	c	c	581 ^d			60			641
1974	553	156	709			61			770
1975	595	128	723			40			763
1976	623	102	725			55			780
1977	839	178	1017			46			1063
1978	695	179	874			27			901
1979	743	123	866			62			928
1980	767	172	939			86			1025
1981	1000	247	1247			98			1345
1982	952	266	1218			105			1323
1983	1025	207	1232			90			1322
1984	1128	332	1460			98			1558
1985	1326	190	1516			82			1598
1986	1304	299	1603			59			1662
1987	1196	386	1582			77			1659
1988	1314	408	1722			51			1773
1989	1452	291	1743			54			1797
1990	1591	416	2007			38			2045
1991	1589	342	1931			49			1980
1992	1642	397	2039	99	58	157	1741	455	2196
1993	1659	419	2078	121	36	157	1780	455	2235
1994	1753	543	2296	127	101	228	1880	644	2524
1995	2012	668	2680	93	30	123	2105	698	2803
1996	2129	580	2709	163	64	227	2292	644	2936
1997	2179	407	2586	77	18	95	2256	425	2681
1998 ^e	1756	307	2063	64	29	93	1820	336	2156
1999	2698	772	3470	45 ^f	10 ^f	71	2743 ^f	782 ^f	3541
2000	2694	746	3440	50 ^f	15 ^f	84	2744 ^f	761 ^f	3524
2001	3198	719	3917	47 ^f	11 ^f	90	3245 ^f	730 ^f	4007
2002	3814	546	4360	48 ^f	7 ^f	67	3862 ^f	553 ^f	4427

Table 1. (cont.)

Year	<u>Tri-State Area</u>			<u>Oregon and Nevada^a</u>			<u>Total RMP</u>		
	White	Cygnets	Total	White	Cygnets	Total	White	Cygnets	Total
2003 ^g	3365	532	3897	62	15	77	3427	547	3974
2004 ^g	3785	746	4531	46	7	53	3831	753	4584
2005	4147	1143	5290	59	12	71	4206	1155	5361
2006	4203	1209	5412	58	14	72	4261	1223	5484
2007 ^h	3604	893	4619	56	26	82	3660	919	4701
2008 ^h	3744	790	4545	74	18	92	3818	808	4637
2009	4287	873	5160	90	15	105	4377	888	5265
2010	3553	676	4229	47	14	61	3600	690	4290
2011	4285	1302	5587	99	26	125	4384	1328	5712

^a Total counts not separated into white birds and cygnets prior to 1992.

^b Not calculated prior to 1992 because of no counts for Oregon and Nevada.

^c Not provided because counts for Yellowstone National Park not separated into white birds and cygnets.

^d In Wyoming only Yellowstone National Park surveyed.

^e 1998 counts for the Tri-state area and Total RMP are biased low because aerial survey of Yellowstone National Park not conducted due to hazardous weather; counted by snowmobile with incomplete coverage.

^f Counts biased low because white-bird and cygnet counts for Malheur NWR not available.

^g Oregon/Nevada and Total RMP counts biased low due to incomplete surveys at Summer Lake WMA.

^h White bird and cygnet counts for Tri-state area and Total RMP biased low because 122 birds in 2007 and 11 birds in 2008 in Idaho were not classified as white birds or cygnets.




Fig. 5. Rates of change for counts of swans in the RMP during the Mid-winter Trumpeter Swan Survey, 1972-2011 (dotted and solid lines depict trends for white birds and cygnets, respectively, for swans counted in the tri-state region [see text]; dashed line depicts total RMP swans).




Fig. 6. Rates of change for counts of total swans in states of the tri-state region during the Mid-winter Trumpeter Swan Survey, 1972-2011 (solid, dotted, and dashed lines represent trends for Montana, Idaho, and Wyoming, respectively).

Table 2. Counts of trumpeter swans of the Rocky Mountain Population in individual states during winter, 1972-2011.

	Montana			Idaho			Wyoming			Oregon ^a			Nevada ^a		
	White			White			White			White			White		
Year	birds	Cygnets	Total	birds	Cygnets	Total	birds	Cygnets	Total	birds	Cygnets	Total	birds	Cygnets	Total
1972	209	14	223	303	14	317	b	b	76			50			41
1973	212	28	240	222	58	280	b	b	61 ^c			32			28
1974	233	40	273	282	109	391	38	7	45			36			25
1975	192	32	224	333	94	427	70	2	72			15			25
1976	253	34	287	308	67	375	62	1	63			30			25
1977	315	43	358	395	126	521	129	9	138			17			29
1978	194	68	262	392	96	488	109	15	124			7			20
1979	304	26	330	353	81	434	86	16	102			41			21
1980	374	80	454	250	70	320	143	22	165			65			21
1981	352	36	388	370	110	480	278	101	379			77			21
1982	390	90	480	429	137	566	133	39	172			65			40
1983	363	59	422	493	122	615	169	26	195			52			38
1984	389	109	498	503	162	665	236	61	297			63			35
1985	393	31	424	701	144	845	232	15	247			51			31
1986	380	73	453	744	183	927	180	43	223			33			26
1987	314	63	377	690	255	945	192	68	260			49			28
1988	438	153	591	694	209	903	182	46	228			24			27
1989	342	90	432	817	141	958	293	60	353			36			18
1990	319	38	357	1025	300	1325	247	78	325			23			15
1991	385	70	455	918	211	1129	286	61	347			31			18
1992	438	114	552	892	249	1141	312	34	346	67	56	123	32	2	34
1993	168	70	238	1020	246	1266	471	103	574	91	36	127	30	0	30
1994	199	48	247	1164	397	1561	390	98	488	114	94	208	13	7	20
1995	153	61	214	1391	475	1866	468	132	600	72	27	99	21	3	24
1996	319	82	401	1336	390	1726	474	108	582	140	49	189	23	15	38
1997	204	30	234	1555	272	1827	420	105	525	46	9	55	31	9	40
1998	290	68	358	1200	200	1400	266 ^d	39 ^d	305 ^d	31	7	38	33	22	55
1999	335	153	488	1754	500	2254	609	119	728	16 ^e	2 ^e	34	29	8	37

Table 2. (cont.)

	Montana			Idaho			Wyoming			Oregon ^a			Nevada ^a		
	White			White			White			White			White		
Year	birds	Cygnets	Total	birds	Cygnets	Total	birds	Cygnets	Total	birds	Cygnets	Total	birds	Cygnets	Total
2000	519	155	674	1881	513	2394	294	78	372	15 ^e	6 ^e	40	35	9	44
2001	373	96	469	2404	549	2953	421	74	495	16 ^e	7 ^e	55	31	4	35
2002	600	104	704	2636	357	2993	578	85	663	7 ^e	5 ^e	24	41	2	43
2003	375	58	433	2490	382	2872	500	92	592	28 ^f	8 ^f	36 ^f	34	7	41
2004	583	92	675	2591	563	3154	611	91	702	8 ^f	0 ^f	8 ^f	38	7	45
2005	508	119	627	2954	828	3782	685	196	881	27	10	37	32	2	34
2006	713	211	924	2714	873	3587	776	125	901	36	14	50	22	0	22
2007	466	49	515	2294 ^g	664 ^g	3080	844	180	1024	38	16	54	18	10	28
2008	382	25	407	2694 ^g	616 ^g	3321	668	149	817	49	16	65	25	2	27
2009	168	21	189	3393	740	4133	726	112	838	53	15	68	37	0	37
2010	274	64	338	2631	501	3132	648	111	759	21	14	35	26	0	26
2011	307	121	428	3068	918	3986	910	263	1173	66	22	88	33	4	37

^a Counts for Oregon and Nevada were not separated into white birds and cygnets until 1992.

^b Not provided because counts for Yellowstone National Park not separated into white birds and cygnets.

^c Counts for Yellowstone National Park only; remainder of Wyoming not surveyed.

^d Counts for Wyoming biased low because aerial survey of Yellowstone National Park not conducted due to hazardous weather; counted by snowmobile with incomplete coverage.

^e Counts biased low because white-bird and cygnet counts for Malheur NWR not available.

^f Counts biased low due to incomplete surveys at Summer Lake WMA.

^g Counts biased low because 122 birds in 2007 and 11 birds in 2008 not classified as either white birds or cygnets.

wintering in Idaho (7.0%) and Wyoming (7.0%) ($P < 0.01$) were higher (Table 2, Fig. 6). Although the numbers of birds wintering in each of the 3 states in the tri-state region generally have increased since 1972, the distribution of birds among the states has changed substantially. Whereas during the 1970s and early 1980s about 36% of wintering swans were counted in Montana, only about 15% of the birds wintering in the tri-state area have been counted there during the last decade (Fig. 7). In contrast, the percentage of birds in Idaho has increased from about 53% to about 69% during that same time period. The percentage of birds counted in Wyoming during winter also has increased, from about 11% to 16%.

Counts of total swans wintering in Nevada have fluctuated over time, but suggest an increase ($P = 0.04$) of about 1.0% per year during 1972-2010 (Table 2, Fig. 8). Counts in Nevada during the early 2000s generally were near historic highs. Trumpeter swans in Oregon primarily occur in 2 areas, Malheur NWR and the Summer Lake WMA and vicinity. Introductions of trumpeter swans to Malheur NWR began in the late 1930s; however, birds were not translocated to Summer Lake WMA until the winter of 1992. Analyzing trends for the Oregon Flock as a whole (Table 2) could lead to inappropriate inferences. Therefore, we analyzed data for Malheur NWR (1972-2010) separate from those for Summer Lake WMA. Results suggest a decline (-3.5% per year, $P < 0.01$) for birds wintering at Malheur NWR (Fig. 8, Appendix A). At Summer Lake WMA, most birds were translocated to the area during winter and generally remained in the area for only a few months after being translocated (M. St. Louis, Oregon Department of Fish and Wildlife, personal communication). Thus, in 1997, the winter following the termination of translocations to Summer Lake WMA, the number counted during the survey dropped sharply (Fig. 8). From 1997-2011, an average of about 33 birds has been observed during winter surveys (excluding years with incomplete surveys).

The percentage of the entire RMP estimated to be comprised of Canadian Flocks increased from about 19% during February of 1972 to 92% during February 2005, and then decreased during 2006 to 2008 (Table 3). During 2009 the percentage rose to 91% but in 2010 dropped to 89%. As of 2011 91.5% of the RMP is comprised of Canadian Flocks. The data fit a 2nd-order logarithm model ($P < 0.01$, adjusted $R^2 = 0.94$), suggesting that the percentage may plateau near 90% (Fig. 9). The average percentage of the RMP being comprised of the Canadian Flock during 2001 – 2003 has been 90%. The number of swans estimated to be from Canadian Flocks exhibited a fairly steady increase since the early 1980s, and was nearly 5,000 birds in 2006, but declined to about 4,100 birds in 2008 (Table 3, Fig. 9). During the 2011 winter survey there were 5,228 birds were from the Canadian flock. This is the first time the estimate was above 5,000 birds since the survey began in 1972.

Results from the 2011 survey

During the 2011 winter survey, observers counted 5,712 trumpeter swans in the RMP, which was a 33% increase from the count of last winter (4,290) (Table 1). The number of white birds increased 22%, whereas the count for cygnets increased 92% from that of last year. The total number of swans




Fig. 7. Proportions of total swans counted in each of the states comprising the tri-state region during the Mid-winter Trumpeter Swan Survey, 1972-2011.




Fig. 8. Rates of change in counts of total swans in Nevada (stars and solid line) and Oregon (Malheur NWR [closed circles and dashed line] and Summer Lake WMA [open circles]) during the Mid-winter Trumpeter Swan Survey, 1972-2011. Data for Summer Lake WMA in 2002 and 2003 are from incomplete surveys.

Table 3. Estimates of swan abundance for flocks comprising the Rocky Mountain Population of Trumpeter swans, 1972-2011.

Year	Mid-winter count	U.S. Breeding Flocks ^a	Canadian Flocks	Percent Canadian Flocks
1972	707	572	135	19.1
1975	763	581	182	23.9
1978	901	544	357	39.6
1981	1345	582	763	56.7
1984	1558	547	1011	64.9
1985	1598	563	1035	64.8
1986	1662	575	1087	65.4
1987	1659	452	1207	72.8
1988	1773	611	1162	65.5
1989	1797	659	1138	63.3
1990	2045	598	1447	70.8
1991	1980	626	1354	68.4
1992	2196	555	1641	74.7
1993	2235	563	1672	74.8
1994	2524	354	2170	86.0
1995	2803	454	2349	83.8
1996	2936	427	2509	85.5
1997	2681	458	2223	82.9
1998	2156	427	1729	80.2
1999	3541	469	3072	86.8
2000	3524	417	3107	88.2
2001	4007	481	3526	88.0
2002	4427	487	3940	89.0
2003	3974	371	3603	90.7
2004	4584	417	4167	90.9
2005	5361	417	4944	92.2
2006	5484	510	4974	90.7
2007	4701	507	4194	89.2
2008	4637	527	4110	88.6
2009	5265	459	4806	91.3
2010	4290	473	3817	89.0
2011	5712	484	5228	91.5

^a From U.S. Fish and Wildlife Service 2010a. Counts are from the previous calendar year (e.g., the 2011 value is from the Fall 2010 survey).




Fig. 9. Percent (bars and solid line) and counts (solid dots) of the entire RMP estimated to be comprised of Canadian Flocks during the Mid-winter Trumpeter Swan Survey, 1972-2011.




Fig. 10. Proportion of cygnets counted in the tri-state region during the Mid-winter Trumpeter Swan Survey, 1974-2011. The solid line depicts the 1974-2010 average.

in Montana increased (25%) for the second year in a row. The count in Wyoming increased by 55% from that of winter 2010, and the count of total swans in Idaho increased by 27% over the same time period. Of the birds wintering in the tri-state area during winter 2011, about 8% were in Montana, 71% were in Idaho, and 21% were counted in Wyoming.

The number of swans in Nevada (37) was 42% higher than that of last year (Table 2, Appendix A). A period of cold weather occurred prior to the survey, resulting in normal ice coverage on the marsh at Ruby Lake NWR. Open water was available on spring-fed ponds and spring outflow areas. The total count was near the long-term average (31 swans). The number of swans counted at Malheur NWR (10) was only 1 more than last year (Appendix A). The count at Summer Lake WMA (78) was a 200% increase from last year's count (26). The timing of the survey and weather conditions allowed for a more representative count which could account for the substantial increase (Marty St. Louis pers. comm.).

The estimated number of swans from Canadian Flocks was 5,228 birds, about 1,411 more swans compared to the estimate from winter 2010 and the highest count to date. The estimate indicated about 91.5% of the RMP counted in winter 2011 was comprised of swans from Canadian Flocks (Table 3, Fig. 9). After increasing steadily from the early 1970s to the late 1990s, the proportion has remained near 90% (range = 88.0%-92.2%) for the last 10 years.

The proportion of cygnets for swans counted in the tri-state region during winter 2011 was 0.233. This value was 22% higher than the 1974-2010 average (0.190) (Fig. 10). This marked the first time since 2007 that production was higher than the average.

In summary, RMP trumpeter swans appeared to increase by 5.5% annually between 1972 and 2010. Most of the increase over that time was attributable to increases in the number of birds in the Canadian Flocks, which estimates suggest comprise approximately 91.5% of the population. Although estimates of the size of the Canadian Flocks from the winter RMP surveys typically are greater than those from the quinquennial surveys, the estimates appear to track each other (U.S. Fish and Wildlife Service 2006). The results suggest that annual estimates of the size of the Canadian Flocks from the winter RMP surveys are reasonable, but may slightly overestimate their abundance.

The survey results from the 2011 Mid-winter survey suggest an increase of 1,411 birds from the count of last year. This was the highest count recorded since the survey began in 1972. Additionally the number of swans counted in Yellowstone National Park (167) was a 600% increase from last year's count of 23 and the highest count since 2007. This year, the Hebgen Lake area west of the Park had more ice than last year so the swans that normally winter on the lake may have pushed into the park (Doug Smith, Yellowstone National Park, pers. comm.).

The 5.5% increase in total swans this year, when compared, with the 19% decrease in the winter count between 2009 and 2010 suggest, that factors such as immigration and emigration into and out of the survey area, and potentially changes in annual survival, could influence large change increases and decreases in estimates of annual abundance. However, without additional information regarding

these factors, causes for these large and biologically unlikely annual changes will remain unexplained.

Collectively the restoration flocks (Oregon and Nevada) have had successive increases since 1997, and the count for 2011 represents the highest count since 1996. This increase was probably due to a more complete survey in Oregon which increased their 2011 count by 200% from 2010.

LITERATURE CITED

- Banko, W.E. 1960. The trumpeter swan: Its history, habits, and population in the United States. North American Fauna, No. 63. U.S. Fish and Wildlife Service, Washington, D.C. 214pp.
- Gale, R.S., E.O. Garton, and I.J. Ball. 1988. The history, ecology and management of the Rocky Mountain Population of trumpeter swans. Unpublished report. U.S. Fish and Wildlife Service, Montana Cooperative Wildlife Research Unit, Missoula. 532pp.
- Joint Agricultural Weather Facility. 2011. Weekly Weather and Crop Bulletin. Vol. 98, No. 11. URL:http://www.usda.mannlib.cornell.edu/usda/waob/weather_weekly//2011s/2011/weather_weekly-03-15-2011.pdf.
- Moser, T.J. 2006. The 2005 North American trumpeter swan survey. U.S. Fish and Wildlife Service, Division of Migratory Bird Management, Denver, CO. 14pp.
- Subcommittee on the Rocky Mountain Population of Trumpeter Swans. 2008. Pacific Flyway management plan for the Rocky Mountain Population of Trumpeter Swans. Pacific Flyway Study Committee, c/o U.S. Fish and Wildlife Service, Pacific Flyway Representative, Portland, OR.
- U.S. Bureau of Reclamation. 2011a. Pacific Northwest Region hydromet historical data access. URL:<http://www.usbr.gov/pn/hydromet/arcread.html>.
- U.S. Bureau of Reclamation. 2011b. Pacific Northwest Region end-of-month water conditions. URL:<http://www.usbr.gov/pn/hydromet/denver.html>.
- U.S. Department of Agriculture. 2011. Mountain snowpack as of February 1, 2011. URL:<ftp://ftp.wcc.nrcs.usda.gov/support/water/westwide/snowpack/wy2011/snow0902.gif>.
- U.S. Fish and Wildlife Service. 2003. Trumpeter swan survey of the Rocky Mountain Population, Winter 2003. Division of Migratory Birds and State Programs, Denver, CO. 28pp.
- U.S. Fish and Wildlife Service. 2006. Trumpeter swan survey of the Rocky Mountain Population, Winter 2006. Division of Migratory Birds and State Programs, Denver, CO. 27pp.
- U.S. Fish and Wildlife Service. 2008a. Trumpeter swan survey of the Rocky Mountain Population,

- Winter 2007. Division of Migratory Birds and State Programs, Denver, CO. 27pp.
- U.S. Fish and Wildlife Service. 2010*a*. Trumpeter swan survey of the Rocky Mountain Population, Fall 2010. Division of Migratory Birds and State Programs, Denver, CO. 38pp.
- U.S. Fish and Wildlife Service. 2010*b*. Waterfowl population status, 2010. U.S. Department of the Interior, Washington, D.C. 64pp.
- Zimpfer, N.L., G.S. Zimmerman, E.D. Silverman, and M.D. Koneff. 2010. Trends in duck breeding populations, 1955-2010. U.S. Fish and Wildlife Service Administrative Report, Division of Migratory Bird Management, Laurel, MD. 25pp.

Appendix A. Counts of trumpeter swans of the Rocky Mountain Population during winter, 1972-2011.

Year	Montana			Idaho			Wyoming (outside Yellowstone NP)		
	White birds	Cygnets	Total	White birds	Cygnets	Total	White birds	Cygnets	Total
1972	209	14	223	303	14	317	16	4	20
1973	212	28	240	222	58	280	a	a	a
1974	233	40	273	282	109	391	7	0	7
1975	192	32	224	333	94	427	40	2	42
1976	253	34	287	308	67	375	30	1	31
1977	315	43	358	395	126	521	86	0	86
1978	194	68	262	392	96	488	63	4	67
1979	304	26	330	353	81	434	15	3	18
1980	374	80	454	250	70	320	63	6	69
1981	352	36	388	370	110	480	37	10	47
1982	390	90	480	429	137	566	76	19	95
1983	363	59	422	493	122	615	81	12	93
1984	389	109	498	503	162	665	87	11	98
1985	393	31	424	701	144	845	78	8	86
1986	380	73	453	744	183	927	91	25	116
1987	314	63	377	690	255	945	85	18	103
1988	438	153	591	694	209	903	115	28	143
1989	342	90	432	817	141	958	197	39	236
1990	319	38	357	1025	300	1325	169	46	215
1991	385	70	455	918	211	1129	225	47	272
1992	438	114	552	892	249	1141	204	30	234
1993	168	70	238	1020	246	1266	293	64	357
1994	199	48	247	1164	397	1561	253	74	327
1995	153	61	214	1391	475	1866	327	91	418
1996	319	82	401	1336	390	1726	344	84	428
1997	204	30	234	1555	272	1827	346	102	448
1998	290	68	358	1200	200	1400	109	15	124
1999	335	153	488	1754	500	2254	317	71	388
2000	519	155	674	1881	513	2394	207	65	272
2001	373	96	469	2404	549	2953	368	63	431
2002	600	104	704	2636	357	2993	447	72	519
2003	375	58	433	2490	382	2872	354	58	412
2004	583	92	675	2591	563	3154	462	58	520
2005	508	119	627	2954	828	3782	561	166	727
2006	713	211	924	2714	873	3587	655	111	766
2007	466	49	515	2294 ^h	664 ^h	3080	700	155	855
2008	382	25	407	2694 ^h	616 ^h	3321	603	142	745
2009	168	21	189	3393	740	4133	638	110	748
2010	274	64	338	2631	501	3132	630	106	736
2011	307	121	428	3068	918	3986	785	221	1006

^a Counts not available.

^b Total counts not separated into white birds and cygnets prior to 1992.

^c Swans first translocated to Summer Lake WMA in 1992.

^d Count biased low because aerial survey not conducted due to hazardous weather; snowmobile count with incomplete coverage only.

^e Count biased low due to incomplete survey coverage.

^h Counts biased low because 122 birds in 2007 and 11 birds in 2008 not classified as white birds or cygnets.

Appendix A. (cont.)

Year	Yellowstone NP			Malheur NWR ^b			Summer Lake WMA ^c			Nevada ^b		
	White birds	Cygnets	Total	White birds	Cygnets	Total	White birds	Cygnets	Total	White birds	Cygnets	Total
1972	a	a	56			50						41
1973	a	a	61			32						28
1974	31	7	38			36						25
1975	30	0	30			15						25
1976	32	0	32			30						25
1977	43	9	52			17						29
1978	46	11	57			7						20
1979	71	13	84			41						21
1980	80	16	96			65						21
1981	241	91	332			77						21
1982	57	20	77			65						40
1983	88	14	102			52						38
1984	149	50	199			63						35
1985	154	7	161			51						31
1986	89	18	107			33						26
1987	107	50	157			49						28
1988	67	18	85			24						27
1989	96	21	117			36						18
1990	78	32	110			23						15
1991	61	14	75			31						18
1992	108	4	112	25	13	38	42	43	85	32	2	34
1993	178	39	217	44	15	59	47	21	68	30	0	30
1994	137	24	161	30	7	37	84	87	171	13	7	20
1995	141	41	182	9	1	10	63	26	89	21	3	24
1996	130	24	154	11	3	14	129	46	175	23	15	38
1997	74	3	77	11	5	16	35	4	39	31	9	40
1998	157 ^d	24 ^d	181 ^d	13	6	19	18	1	19	33	22	55
1999	292	48	340	a	a	16	16	2	18	29	8	37
2000	87	13	100	a	a	19	15	6	21	35	9	44
2001	53	11	64	a	a	32	16	7	23	31	4	35
2002	131	13	144	a	a	12	7 ^e	5 ^e	12 ^e	41	2	43
2003	146	34	180	19	5	24	9 ^e	3 ^e	12 ^e	34	7	41
2004	149	33	182	8	0	8	a	a	a	38	7	45
2005	124	30	154	8	0	8	19	10	29	32	2	34
2006	121	14	135	15	5	20	21	9	30	22	0	22
2007	144	25	169	4	0	4	34	16	50	18	10	28
2008	65	7	72	12	2	14	37	14	51	25	2	27
2009	88	2	90	17	3	20	36	12	48	37	0	37
2010	18	5	23	7	2	9	14	12	26	26	0	26
2011	125	42	167	7	3	10	59	19	78	33	4	37

^a Counts not available.

^b Total counts not separated into white birds and cygnets prior to 1992.

^c Swans first translocated to Summer Lake WMA in 1992.

^d Count biased low because aerial survey not conducted due to hazardous weather; snowmobile count with incomplete coverage only.

^e Count biased low due to incomplete survey coverage.

Appendix B. Site-specific counts of trumpeter swans of the Rocky Mountain Population during the Mid-winter Trumpeter Swan Survey, 2011

State or Area	White birds	Cygnets	Total	Pilot/observer/notes
Montana				
<i>Hebgen Lake area</i>				P: N. Caldwell; O: D. Smith (2/2)
Cougar Creek	0	0	0	
Between Quake Lake and Hebgen Lake	0	0	0	
Madison River Arm	48	11	59	
North Spring (Grayling Arm)	11	6	17	
South Fork Arm	77	27	104	
South Fork Madison River/Buttermilk Creek	0	0	0	
Subtotal	136	44	180	
<i>Madison River Valley</i>				P: D. Chapman; O: B. West (1/31)
Odell Creek Area	108	64	172	
Walsh Ponds (south)1	0	0	0	
Walsh Ponds (north)1	0	0	0	
Madison River, south of Ennis	25	12	37	
Madison River, north of Ennis	0	0	0	
Ennis Lake	0	0	0	
Subtotal	133	76	209	
<i>Chain of Lakes</i>				
Cliff Lake	0	0	0	
Wade Lake	0	0	0	
Goose Lake	0	0	0	
Smith Creek (Hidden Lake outlet)	0	0	0	
Subtotal	0	0	0	
<i>Centennial Valley/Red Rock Lakes NWR</i>				
Red Rock River below Lower Lake Dam	0	0	0	
MacDonald Pond	0	0	0	
Culver Pond	11	0	11	
Elk Springs Creek	2	1	3	
Swan Lake	0	0	0	
Shambow Pond	0	0	0	
Red Rock River, Lima	0	0	0	
Subtotal	13	1	14	
<i>Paradise Valley</i>				P: N. Cadwell; O: none (2/2)
Armstrong's Spring Creek	0	0	0	
Bailey's	0	0	0	
Brockway	0	0	0	
DePuys	12	0	12	
Brandis	0	0	0	

Nelson's Spring Creek	0	0	0	
Sacagawea Park	0	0	0	
Yellowstone River 1 mile north of Emigrant	0	0	0	
Beaver Creek	0	0	0	
Yellowstone River - 6 mile	0	0	0	
Yellowstone River - Pray	0	0	0	
Yellowstone River - Pine Creek	13	0	13	
Dana's	0	0	0	
Emigrant Pond	0	0	0	
PMD Ranch	0	0	0	
Subtotal	25	0	25	
MONTANA TOTAL	307	121	428	
Wyoming				
<i>Upper Snake River (Flagg Ranch to Wilson Bridge)</i>				P: D. Stinson; O: S. Patla (2/3 and 2/11)
Polecat Creek	9	8	17	
Flagg Ranch to Jackson Lake	8	3	11	
Jackson Lake	0	3	3	
Jackson Lake to Moran Junction	16	3	19	Dense pancake and shelf ice
Moran Junction to Deadman's	8	2	10	
Deadman's to Moose	32	8	40	
Moose to Gros Ventre Junction	25	17	42	
Gros Ventre Junction area	47	2	49	
Gros Ventre Junction to Wilson Bridge	16	0	16	
Gros Ventre River, Highway 89 to Snake River	3	0	3	
Subtotal	164	46	210	
<i>Gros Ventre River upriver of Kelly</i>				
Kelly Warm Springs, Grand Teton National Park				
Lower Slide Lake				
Upper Gros Ventre				
Subtotal	0	0	0	
<i>Lower Snake River (Wilson Bridge to Alpine)</i>				
Wilson Bridge to South Park Bridge	35	25	60	
Evan's Gravel pit ponds	0	0	0	
South Park Bridge to Hoback	2	4	6	
North Wilson	2	0	2	
Fish Creek, Wilson to Snake River	91	14	105	
Boyles Hill area	40	0	40	
Spring Creek	34	11	45	
Crane Creek	14	8	22	
Lower Flat Creek, Snake River to Jackson	18	6	24	
Rafter J Ponds	5	2	7	
Valley Springs, Captive Swan Pond/Pen Highway 89	8	6	14	
Hoback to Astoria Bridge	0	0	0	

Astoria Bridge-Elbow	11	0	11	
Elbow to Alpine/Palisades Reservoir	0	0	0	
Bailey Lake	2	0	2	
Kelly Swan Facility	3	5	8	
Bondurant pond near Hoback River	2	0	2	
Subtotal	267	81	348	
<i>National Elk Refuge</i>				
Flat Creek main marsh	20	1	21	
Gros Ventre River, Kelly to Highway 89	6	2	8	
Romney pond area				
Lost Spring	10	8	18	
Subtotal	36	11	47	
<i>Salt River (Alpine to Afton)</i>				
Palisades Reservoir, WY Alpine	0	0	0	
Palisades Reservoir to Freedom Road	47	12	59	
Freedom Road to Narrows	9	0	9	
Thayne area	3	0	3	
Narrows to Grover/Auburn Highway	40	19	59	
Grover/Auburn Highway to Swift Creek	51	1	52	
Swift Creek to Headwaters	0	0	0	
Subtotal	150	32	182	
<i>Pinedale</i>				
New Fork Boulder to Pinedale	0	0	0	
Boulder Fish Hatchery				
Daniel Fish Hatchery/Forty Rod Creek	18	9	27	
Warren Bridge to Kendall Bridge, Green River	0	0	0	Frozen
Kendall Bridge to Green River Lakes	0	0	0	Frozen
Subtotal	18	9	27	
<i>Green River (Warren Bridge to Highway 28 Bridge)</i>				
Fontenelle Dam-CCC Bridge	15	3	18	
CCC Bridge to Pilot Farm	120	31	151	
Pilot Farm-Refuge Headquarters	15	8	23	
Refuge to Big Sandy				Not surveyed
Big Sandy to Big Island				Not surveyed
Flaming Gorge Reservoir				Not surveyed
Subtotal	150	42	192	
<i>Dubois area</i>				
Wind River and spring ponds, Dubois	2	2	4	
Dinwoody Lake	23	6	29	
Bull Lake	2	0	2	
Wind River, Dinwoody to Crowheart				

Subtotal	27	8	35	
<i>Yellowstone National Park</i>				
Slough Creek	0	0	0	P: N. Cadwell; O: D. Smith (2/2)
Tern Lake	0	0	0	
Broad Creek, near White Lake	0	0	0	
White Lake	8	4	12	
Beach Springs Lagoon	0	0	0	
Shoshone Geyser Basin	0	0	0	
Lewis River	0	0	0	
Buela Lake	0	0	0	
Yellowstone River	11	0	11	
Lewis - Shoshone Channel	0	0	0	
Lewis Lake	0	0	0	
Falls River	0	0	0	
Shoshone Lake	0	0	0	
Bechler Lake	0	0	0	
Bourndary Creek	0	0	0	
Bechler River	0	0	0	
Firehole River	0	0	0	
Madison River (Madison Jct. to Park boundary)	95	32	127	
Richard's Pond	0	0	0	
Gibbon Meadow	0	0	0	
Nymph Lake	11	6	17	
Elk Park	0	0	0	
North Twin Lake	0	0	0	
Nez Perce Creek near Culex Basin	0	0	0	
Nez Perce Creek near Cowan Creek	0	0	0	
Alum Creek	0	0	0	
Gibbon River north of Madison Junction	0	0	0	
Mud Volcano	0	0	0	
Subtotal	125	42	167	
TOTAL WY outside YNP	785	221	1006	
TOTAL WY including YNP	910	263	1173	
Idaho				P: C. Anderson; O: M. Fisher(2/2-3)
<i>Island Park Area</i>				
Warm Springs (west side of Henrys Lake)				Frozen not flown
Henrys Lake flats				Frozen not flown
Big Springs, North Fork, Mack's Inn Area	5	2	7	
Mack's Inn to Island Park Reservoir	9	2	11	
Island Park Reservoir				Frozen
Island Park Reservoir inlet				Frozen
Trude Ranch Pond				Frozen
Icehouse Reservoir				Frozen
Sheridan Creek, mouth to Sheridan Reservoir	10	17	27	Very little open water
Sheridan Reservoir	0	0	0	

Sheridan Creek cabin and pond	4	5	9	
Subtotal	28	26	54	
<i>Buffalo River Area</i>				
Buffalo River	5	0	5	
Tom's Creek	0	0	0	
Elk Creek/Trudes Siding pond	11	4	15	
Subtotal	16	4	20	
<i>Harriman State Park (HSP) Area</i>				
Island Park Dam through Box Canyon	22	1	23	
Box Canyon - HSP north boundary	58	3	61	
HSP north bounday - Osborne bridge	154	45	199	
Golden Lake	20	4	24	Est 95% frozen
Thurmon Creek	13	1	14	Est 70% frozen
Silver Lake				Frozen
Osborne Bridge - Pinehaven	39	6	45	
Pinehaven	52	15	67	
Fish Pond				frozen
Henrys Fork below Pinehaven - Forest boundary				
Subtotal	358	75	433	
<i>Henrys Fork, HSP to Warm River</i>				
Warm River	0	0	0	
Subtotal	0	0	0	
<i>Lower Henrys Fork Area</i>				
Warm River confluence to Ashton Dam	0	0	0	
Ashton Dam to Chester Dam	214	34	248	
Chester Dam to Highway 33	64	16	80	
Highway 33 - Menan Buttes	0	0	0	
Ashton Ponds				Frozen not flown
Willow Creek Area farmstead ponds	0	0	0	
Mikesell Reservoir 1 & 2				Frozen not flown
Arcadia Reservoir, Upper				Frozen not flown
Arcadia Reservoir, Lower				Frozen not flown
Sand Creek WMA and area	4	0	4	
Singleton Ponds				Frozen not flown
Texas Slough	99	14	113	
Bannock Jim Slough				Frozen not flown
Mud Lake WMA				Frozen not flown
Camas NWR				Frozen not flown
Camas Creek				Frozen not flown
Subtotal	381	64	445	
<i>Teton River Basin</i>				

Teton River to Wilford Dam	6	4	10	
Wilford Dam to Newdale Bridge	30	7	37	
Newdale Bridge to Teton Dam site	71	9	80	
Teton River Canyon	25	3	28	
Teton Basin	186	76	262	
North Fork Teton River				
South Fork Teton River	0	0	0	
Subtotal	318	99	417	
<i>South Fork of the Snake River</i>				
Swan Valley (Palisades Reservoir to Conant Valley)	230	71	301	
Canyon (Conant to Heise)	140	60	200	
Delta (Heise to Menan Buttes)	141	78	219	
Dry bed (Heise to Menan)	0	0	0	
Subtotal	511	209	720	
<i>Main Stem of the Snake River</i>				
Menan Buttes to Idaho Falls	576	163	739	
Dry Bed				
Idaho Falls to Fort Hall (Ferry Butte)	60	14	74	
Blackfoot Marsh	0	0	0	
Subtotal	636	177	813	
<i>Fort Hall Bottoms to American Falls Reservoir</i>				
American Falls Reservoir shoreline	314	75	389	
Kinney Creek	0	0	0	
Portneuf River (Am. Falls Res. to Hwy 86)	56	13	69	
Mouth of Portneuf River				
Spring Creek to American Falls Reservoir	111	28	139	
Jimmie Creek	0	0	0	
Snake River - Tilden Bridge	15	6	21	
Clear Creek and Ross Fork	0	0	0	
Diggie Creek	0	0	0	
Jeff Cabin Creek	6	2	8	
Flying Y oxbows	0	0	0	
Field feeding, Vambaur Road	8	5	13	New 2011, IDFG grd srv 2/2 10:30-4:30
Subtotal	510	129	639	
<i>Snake River below American Falls Dam</i>				
Springfield Reservoir	19	12	31	IDFG grd srv, 2/2 10"30-4:30, 4/5 Trus, 0 TUSW
American Falls Reservoir (except Fort Hall)				
American Falls Dam - Minidoka NWR	5	5	10	Flight data, open only to Mass Rocks SP, frozen below
Minidoka NWR				Not counted, frozen
Minidoka Dam - C.J. Strike Reservoir	13	6	19	IDFG grd srv, 2/3, 5/2 TUSW, 4/2 MUTE (Clear Lk)
Hagerman National Fish Hatchery				Not counted
Bruneau Dunes State Park				Not counted

Bruneau Dunes - C.J. Stike Reservoir				Not counted
Faulkner Pond				Not counted
White Arrow Pond (Bliss)	15	10	25	IDFG grd srv, 2/3, 0 TUSW
Pioneer Reservoir (King Hill)	2	6	8	IDFG grd srv, 2/3, 0 TUSW
Snake River at King Hill				
Silver Creek (Picabo area)	54	19	73	IDFG grd srvy, 2/4
Miracle Hot Springs				Not counted
Dead Horse Lake				Not counted
Butler Pond	25	17	42	2011 addition, IDFG grd srvy, 2/3, 4/0 TUSW
Subtotal	133	75	208	
<i>Grays Lake NWR Area</i>				
Big Springs				Frozen not flown
Shorty's Homestead				Frozen not flown
Blackfoot Reservoir	17	4	21	
Chub Springs, southwest of refuge	7	3	10	
Chesterfield Reservoir				frozen
<i>Chesterfield Reservoir Canal (portneuf R. headwaters)</i>	0	0	0	
<i>Grimm Spring and channel</i>	0	0	0	
<i>U. Portneuf river: Toponce Rd - Pebble Cr Rd</i>	0	0	0	First bridge; includes IDFG's whiskey mike's access
<i>Pebble Cr Rd - Broxon Rd</i>	13	2	15	Second bridge; includes IDFG's U. Portneuf access
<i>Broxon Rd - Symons Rd</i>	0	0	0	Third bridge
<i>Symons Rd - Blazer Hwy. Bridge</i>	3	0	3	Fourth bridge; includes IDFG's Portneuf access
<i>Blazer Hwy. bridge - Hwy 30 Bridge</i>	0	0	0	
Subtotal	40	9	49	
<i>Soda Springs Area</i>				
Woodall Springs	7	1	8	
Alexander Reservoir and Siding				
Miller Ponds	7	2	9	
Government Dam	0	0	0	
Soda Creek	3	1	4	
Soda Canal	0	0	0	
Subtotal	17	4	21	
<i>Bear River Reaches</i>				
Alexander Reservoir	0	0	0	
Alexander Reservoir - Gentile Valley Bridge	65	34	99	
Gentile Valley Bridge - old cheese factory				Frozen, not flown
Gentile Valley Bridge to Oneida Dam	41	7	48	
Montpelier Reservoir (rearing pond)				Frozen, not flown
Oneida Narrows	0	0	0	
Oneida Narrows to Riverdale Bridge	0	0	0	
Riverdale Bridge to Utah border	0	0	0	
Subtotal	106	41	147	

<i>Bear Lake National Wildlife Refuge</i>				
Bear Lake - Alexander Res.	14	6	20	
West Canal Unit				
Rainbow Unit				
Outlet Canal				
Subtotal	14	6	20	
IDAHO TOTAL	3068	918	3986	
Nevada				
Ruby Lake NWR	25	2	27	J. Mackay (2/3)
Franklin Lake				Dry
Oregon				
<i>Malheur NWR</i>				
Refuge total	7	3	10	J. Dastyck (242)
<i>Summer Lake Wildlife Management Area</i>				
Summer Lake WMA	59	19	78	M. St. Louis (2/9)

^aBlank denotes area not surveyed.

Appendix C. Personnel who conducted the 2011 Mid-winter Trumpeter Swan Survey.

Montana (Red Rock Lakes NWR, Centennial Valley, Madison Valley)

Observer: B. West (Red Rock Lakes NWR)
Pilot: D. Chapman (Montana Aircraft, Inc.)

Montana (Hebgen Lake Area and Paradise Valley)

Observer: D. Smith (Yellowstone National Park)
Pilot: N. Cadwell (Elkhorn Aviation)

Idaho

Observer: M. Fisher (Southeast Idaho National Wildlife Refuge Complex)
Pilot: C. Anderson (AvCenter)

Wyoming

Observer: S. Patla (Wyoming Game and Fish Department)
Pilot: D. Stinson (Sky Aviation)

Wyoming (Yellowstone National Park)

Observer: D. Smith (Yellowstone National Park)
Pilot: N. Cadwell (Elkhorn Aviation)

Ruby Lake NWR and vicinity

J. Mackay (Ruby Lake NWR)

Malheur NWR

J. Dastyck (Malheur NWR)

Summer Lake WMA

M. St. Louis (Oregon Department of Fish and Wildlife)
