

2009

Role of *vapBC* toxin–antitoxin loci in the thermal stress response of *Sulfolobus solfataricus*

Charlotte R. Cooper

North Carolina State University

Amanda J. Daugherty

University of Nebraska-Lincoln

Sabrina Tachdjian

North Carolina State University

Paul H. Blum

University of Nebraska - Lincoln, pblum1@unl.edu

Robert M. Kelly

North Carolina State University

Follow this and additional works at: <http://digitalcommons.unl.edu/bioscifacpub>



Part of the [Biology Commons](#)

Cooper, Charlotte R.; Daugherty, Amanda J.; Tachdjian, Sabrina; Blum, Paul H.; and Kelly, Robert M., "Role of *vapBC* toxin–antitoxin loci in the thermal stress response of *Sulfolobus solfataricus*" (2009). *Faculty Publications in the Biological Sciences*. 516.
<http://digitalcommons.unl.edu/bioscifacpub/516>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in the Biological Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Published in final edited form as:

Biochem Soc Trans. 2009 February ; 37(Pt 1): 123–126. doi:10.1042/BST0370123.

Copyright 2009 by the Authors. Journal compilation (c) 2009 Biochemical Society.

Role of *vapBC* toxin–antitoxin loci in the thermal stress response of *Sulfolobus solfataricus*

Charlotte R. Cooper^{*}, Amanda J. Daugherty[†], Sabrina Tachdjian^{*}, Paul H. Blum[†], and Robert M. Kelly^{*,1}

^{*} Department of Chemical and Biomolecular Engineering, North Carolina State University, Raleigh, NC 27695-7905, U.S.A.

[†] Beadle Center for Genetics, University of Nebraska-Lincoln, Lincoln, NE 68588-0666, U.S.A.

Abstract

TA (toxin–antitoxin) loci are ubiquitous in prokaryotic microorganisms, including archaea, yet their physiological function is largely unknown. For example, preliminary reports have suggested that TA loci are microbial stress-response elements, although it was recently shown that knocking out all known chromosomally located TA loci in *Escherichia coli* did not have an impact on survival under certain types of stress. The hyperthermophilic crenarchaeon *Sulfolobus solfataricus* encodes at least 26 *vapBC* (where *vap* is virulence-associated protein) family TA loci in its genome. VapCs are PIN (PilT N-terminus) domain proteins with putative ribonuclease activity, while VapBs are proteolytically labile proteins, which purportedly function to silence VapCs when associated as a cognate pair. Global transcriptional analysis of *S. solfataricus* heat-shock-response dynamics (temperature shift from 80 to 90°C) revealed that several *vapBC* genes were triggered by the thermal shift, suggesting a role in heat-shock-response. Indeed, knocking out a specific *vapBC* locus in *S. solfataricus* substantially changed the transcriptome and, in one case, rendered the crenarchaeon heat-shock-labile. These findings indicate that more work needs to be done to determine the role of VapBCs in *S. solfataricus* and other thermophilic archaea, especially with respect to post-transcriptional regulation.

Keywords

archaeon; heat shock; hyperthermophile; stress response; *Sulfolobus solfataricus*; toxin–antitoxin locus

Introduction

TA (toxin–antitoxin) loci, also known as plasmid addiction or poison–antidote systems, were first identified as a plasmid maintenance mechanism that activated PSK (post-segregational killing) in plasmid-free progeny [1]. These loci encode a cognate protein pair, consisting of a proteolytically labile antitoxin and a toxin [2,3]. Available genome sequence data indicate that TA loci are also chromosomally encoded and ubiquitous in free-living prokaryotes [4]. The widespread occurrence of chromosomally encoded TA loci in the microbial world suggests an important function, although the role of these proteins in either a specific or general sense is largely unknown.

¹To whom correspondence should be addressed (rmkelly@eos.ncsu.edu).

TA loci are typically arranged in operons with the antitoxin gene preceding the toxin gene (except for *higBA* and *hipBA* where the toxin precedes the antitoxin) [5–7]. The TA ORFs (open reading frames) often overlap, making co-expression likely. The functional relationship between toxin and antitoxin proteins appears to be consistent across microbiology, although there is still only limited experimental evidence along these lines. As long as the antitoxin is present, it interacts with the cognate toxin to presumably neutralize toxic activity, which at least in some cases is ribonucleolytic [8–14]. However, when the antitoxins are lost in plasmid-free progeny or proteolytically degraded in chromosomally encoded cases, the free toxins are activated. The details of TA function are unique to specific types. For example, in some cases, TA operons are apparently regulated by the binding of the N-terminus of the antitoxin to the locus promoter region [15,16]. Toxins may also function as co-repressors, since their binding to the antitoxin appears to increase the affinity of the latter for the promoter region [17].

The precise function of chromosomally encoded TA loci remains controversial. TA loci may be bacteriocidal [18] or bacteriostatic [19,20]. In fact, *hipBA* TA loci have been linked to persister cell formation [21]. In *Escherichia coli* K-12, knocking out all five of the known chromosomally encoded TA loci did not have a significant impact on the survival of pH, nutritional or antibiotic stress [22]. Chromosomally encoded TA loci may act as anti-addiction modules that protect cells from PSK by plasmid-encoded TA loci [23]. Whereas TA loci in mesophilic bacteria have been closely examined, this is not the case in the archaea. Many archaeal genomes, particularly thermophilic archaea, encode multiple TA loci. The significance of these TAs under normal or abnormal growth conditions remains to be seen.

***vapBC* TA loci and archaea**

The distribution of members of eight TA families in prokaryotic genomes is widespread. Out of 218 prokaryotic genomes surveyed, over 1472 TA loci have been identified with an additional 63 solitary toxins or antitoxins found [4,24]. To date, the *vapBC* (where *vap* is virulence-associated protein) family is the most abundant TA system among prokaryotic genomes, representing ~40% of all TA loci [4]. It is present in high numbers in the archaea, especially in hyperthermophiles and extreme thermoacidophiles (Figure 1).

Most VapB antitoxins contain an SpoVT/AbrB DNA-binding domain and, as such, belong to the superfamily of transcriptional regulators of the same name. AbrB, which has been studied extensively in *Bacillus subtilis* and *Bacillus anthracis*, is a transition-state regulator [25–27]. SpoVT, an AbrB homologue, was shown to regulate expression of at least 15 genes, probably via DNA-binding interactions with target promoters [28]. VapC toxins are characterized by a PIN (Pil-T N-terminus) domain (a domain homologous with the N-terminal domain of the pilin biogenesis protein Pil-T) [29]. In eukaryotes, PIN domain proteins are ribonucleases involved in nonsense-mediated mRNA decay and RNAi (RNA interference) [30]. PIN domains could provide clues to the cellular targets of VapC toxins, but this connection has yet to be made experimentally. Generally, VapCs are putative ribonucleases, although their precise specificity is not clear. For example, they were found to have endonuclease activity in mycobacteria and exonuclease activity in the hyperthermophilic crenarchaeon *Pyrobaculum aerophilum* [8,9]. Furthermore, a VapC from *Haemophilus influenzae* was determined to be ribonucleolytic, degrading free RNA *in vitro* [11].

Heat-shock-response of hyperthermophilic archaea

Even though extremely thermophilic archaea thrive at extreme temperatures, they also have thermal limits and display a classic heat-shock-response when thermally stressed [31–38].

This response involves the thermosome, or rosettasome, a heat-shock-responsive HSP (heat-shock protein) 60-like molecular chaperone that has been implicated in many cellular roles [39,40]. Examination of several hyperthermophilic archaea undergoing thermal stress has revealed that heat shock has a profound effect on the transcriptome. *Pyrococcus furiosus* response to a temperature shift from 90 to 105°C, included up-regulation of the thermosome (KEGG accession number PF1974) and a HSP20-like sHSP (small heat-shock protein) (accession number PF1883) [35], in addition to several hundred ORFs (K.R. Shockley and R.M. Kelly, unpublished work). Detailed analysis revealed that a novel heat-shock-regulator protein, Phr, in *P. furiosus* prevented synthesis of HSP20, AAA+ (ATPase associated with various cellular activities) (whose function is unclear) and itself by binding to the promoter regions and blocking the RNA polymerase-binding site when under both thermal stress as well as during nutrient-limited stationary growth phase [38]. The heat-shock-response of other hyperthermophilic archaea has also been examined. A temperature shift from 85 to 95°C was lethal for *Methanococcus jannaschii*; after 20 min at 95°C, 76 genes were up-regulated significantly (>2-fold), including an sHSP (KEGG accession number MJ0285) and the thermosome (KEGG accession number MJ0999) [31]. In *Archaeoglobus fulgidus*, 10% of the genome was differentially transcribed after only 5 min at 89°C (up from the normal growth temperature of 78°C); up-regulated ORFs included six of 13 known heat-shock-related genes (KEGG accession numbers AF1296, AF1297, AF1298, AF1451, AF2238 and AF1971). After 1 h at 89°C, 14% of the *A. fulgidus* genome displayed changes in mRNA transcription levels [34]. Among the hyperthermophilic archaea examined for heat-shock-response, *Sulfolobus solfataricus* exhibited the most pronounced change in transcriptome, with approx. one-third of its genome responding to a shift from 80 to 90°C within 5 min of reaching the target temperature; 37% of the up-regulated genes were insertion sequences. Both HSP20 family sHSPs (KEGG accession numbers SSO2427 and SSO2603) were up-regulated, as were many of the *vapBC* TA loci found in *S. solfataricus* [36]. Some TA loci were constitutively expressed at high levels (e.g. *vapBC22*), but thermal stress triggered even higher transcription levels (Figure 2). Other TA loci were significantly up-regulated by thermal stress, such as *vapBC6* and *vapBC8* of *S. solfataricus*. Using *S. solfataricus* strain PBL2025, genetic insertions were made to disrupt the function of individual TA genes [41]. When toxin *vapC22* was disrupted, no obvious phenotype was noted, although approx. 100 ORFs were differentially transcribed 2-fold or more (C.R. Cooper, S. Tachdjain, A.J. Daugherty, P.H. Blum and R.M. Kelly, unpublished work). However, disruption of *vapB6* (and consequently *vapC6*) rendered the organism susceptible to thermal stress (C.R. Cooper, S. Tachdjain, A.J. Daugherty, P.H. Blum and R.M. Kelly, unpublished work). Efforts are now underway to seek to determine the set of essential TA loci required by *S. solfataricus* to survive thermal stress.

Concluding remarks

Because many TA loci are still annotated as 'hypothetical proteins', they are often overlooked as important elements in microbial genomes. Transcriptional data from *S. solfataricus* heat-shock experiments suggested that, although chromosomally encoded TA loci may not play a significant role in mesophilic prokaryotes such as *E. coli*, they are potentially significant stress-response modules in thermophilic archaea. One hypothesis yet to be tested is that TA loci are key components in archaeal RNA management systems. The significance of the large complement of TA loci in thermophilic archaea (as noted, there are 26 *vapBC* loci in *S. solfataricus* alone) remains a mystery. Each locus may play a specific role under normal or stress conditions. Further work is needed to define the targets of specific VapCs in *S. solfataricus* as this may lead to a clearer picture of TA loci in archaeal biology.

Acknowledgments

Funding

This work was supported in part by the National Science Foundation [grant numbers CBET0730091 and CBET0617272]. C.R.C. acknowledges support from a National Institutes of Health T32 Biotechnology Traineeship.

Abbreviations used

HSP	heat-shock protein
ORF	open reading frame
PIN	Pil-T N-terminus
PSK	post-segregational killing
sHSP	small heat-shock protein
TA	toxin–antitoxin
<i>vap</i>	virulence-associated protein

References

- Gerdes K, Rasmussen PB, Molin S. Unique type of plasmid maintenance function: postsegregational killing of plasmid-free cells. *Proc Natl Acad Sci USA* 1986;83:3116–3120. [PubMed: 3517851]
- Jensen RB, Gerdes K. Programmed cell death in bacteria: proteic plasmid stabilization systems. *Mol Microbiol* 1995;17:205–210. [PubMed: 7494469]
- Van Melder L, Bernard P, Couturier M. Lon-dependent proteolysis of CcdA is the key control for activation of CcdB in plasmid-free segregant bacteria. *Mol Microbiol* 1994;11:1151–1157. [PubMed: 8022284]
- Pandey DP, Gerdes K. Toxin–antitoxin loci are highly abundant in free-living but lost from host-associated prokaryotes. *Nucleic Acids Res* 2005;33:966–976. [PubMed: 15718296]
- Black DS, Kelly AJ, Mardis MJ, Moyed HS. Structure and organization of *hip*, an operon that affects lethality due to inhibition of peptidoglycan or DNA synthesis. *J Bacteriol* 1991;173:5732–5739. [PubMed: 1715862]
- Gerdes K. Toxin–antitoxin modules may regulate synthesis of macromolecules during nutritional stress. *J Bacteriol* 2000;182:561–572. [PubMed: 10633087]
- Tian QB, Ohnishi M, Tabuchi A, Terawaki Y. A new plasmid-encoded proteic killer gene system: cloning, sequencing, and analyzing *hig* locus of plasmid Rts1. *Biochem Biophys Res Commun* 1996;220:280–284. [PubMed: 8645296]
- Arcus VL, Backbro K, Roos A, Daniel EL, Baker EN. Distant structural homology leads to the functional characterization of an archaeal PIN domain as an exonuclease. *J Biol Chem* 2004;279:16471–16478. [PubMed: 14734548]
- Arcus VL, Rainey PB, Turner SJ. The PIN-domain toxin–antitoxin array in mycobacteria. *Trends Microbiol* 2005;13:360–365. [PubMed: 15993073]
- Christensen SK, Pedersen K, Hansen FG, Gerdes K. Toxin–antitoxin loci as stress-response-elements: ChpAK/MazF and ChpBK cleave translated RNAs and are counteracted by tmRNA. *J Mol Biol* 2003;332:809–819. [PubMed: 12972253]
- Daines DA, Wu MH, Yuan SY. VapC-1 of nontypeable *Haemophilus influenzae* is a ribonuclease. *J Bacteriol* 2007;189:5041–5048. [PubMed: 17496075]
- Kamada K, Hanaoka F. Conformational change in the catalytic site of the ribonuclease YoeB toxin by YefM antitoxin. *Mol Cell* 2005;19:497–509. [PubMed: 16109374]

13. Zhang Y, Zhang J, Hoeflich KP, Ikura M, Qing G, Inouye M. MazF cleaves cellular mRNAs specifically at ACA to block protein synthesis in *Escherichia coli*. *Mol Cell* 2003;12:913–923. [PubMed: 14580342]
14. Zhang Y, Zhu L, Zhang J, Inouye M. Characterization of ChpBK, an mRNA interferase from *Escherichia coli*. *J Biol Chem* 2005;280:26080–26088. [PubMed: 15901733]
15. Kedzierska B, Lian LY, Hayes F. Toxin–antitoxin regulation: bimodal interaction of YefM–YoeB with paired DNA palindromes exerts transcriptional autorepression. *Nucleic Acids Res* 2007;35:325–339. [PubMed: 17170003]
16. Magnuson P, Lehnher H, Mukhopadhyay G, Yarmolinsky MB. Autoregulation of the plasmid addiction operon of bacteriophage P1. *J Biol Chem* 1996;271:18705–18710. [PubMed: 8702525]
17. Gerdes K, Christensen SK, Lobner-Olesen A. Prokaryotic toxin–antitoxin stress response loci. *Nat Rev Microbiol* 2005;3:371–382. [PubMed: 15864262]
18. Engelberg-Kulka H, Amitai S, Kolodkin-Gal I, Hazan R. Bacterial programmed cell death and multicellular behavior in bacteria. *PLoS Genet* 2006;2:e135. [PubMed: 17069462]
19. Pedersen K, Christensen SK, Gerdes K. Rapid induction and reversal of a bacteriostatic condition by controlled expression of toxins and antitoxins. *Mol Microbiol* 2002;45:501–510. [PubMed: 12123459]
20. Pedersen K, Zavialov AV, Pavlov MY, Elf J, Gerdes K, Ehrenberg M. The bacterial toxin RelE displays codon-specific cleavage of mRNAs in the ribosomal A site. *Cell* 2003;112:131–140. [PubMed: 12526800]
21. Correia FF, D'Onofrio A, Rejtar T, Li LY, Karger BL, Makarova K, Koonin EV, Lewis K. Kinase activity of overexpressed HipA is required for growth arrest and multidrug tolerance in *Escherichia coli*. *J Bacteriol* 2006;188:8360–8367. [PubMed: 17041039]
22. Tsilibaris V, Maenhaut-Michel G, Mine N, Van Melder L. What is the benefit to *Escherichia coli* of having multiple toxin–antitoxin systems in its genome? *J Bacteriol* 2007;189:6101–6108. [PubMed: 17513477]
23. Saavedra De Bast M, Mine N, Van Melder L. Chromosomal toxin–antitoxin systems may act as anti-addiction modules. *J Bacteriol* 2008;190:4603–4609. [PubMed: 18441063]
24. Makarova KS, Grishin NV, Shabalina SA, Wolf YI, Koonin EV. A putative RNA-interference-based immune system in prokaryotes: computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action. *Biol Direct* 2006;1:7. [PubMed: 16545108]
25. Strauch MA, Ballar P, Rowshan AJ, Zoller KL. The DNA-binding specificity of the *Bacillus anthracis* AbrB protein. *Microbiology* 2005;151:1751–1759. [PubMed: 15941984]
26. Vaughn JL, Feher V, Naylor S, Strauch MA, Cavanagh J. Novel DNA binding domain and genetic regulation model of *Bacillus subtilis* transition state regulator *abrB*. *Nat Struct Biol* 2000;7:1139–1146. [PubMed: 11101897]
27. Vaughn JL, Feher VA, Bracken C, Cavanagh J. The DNA-binding domain in the *Bacillus subtilis* transition-state regulator AbrB employs significant motion for promiscuous DNA recognition. *J Mol Biol* 2001;305:429–439. [PubMed: 11152601]
28. Bagyan I, Hobot J, Cutting S. A compartmentalized regulator of developmental gene expression in *Bacillus subtilis*. *J Bacteriol* 1996;178:4500–4507. [PubMed: 8755877]
29. Anantharaman V, Aravind L. New connections in the prokaryotic toxin–antitoxin network: relationship with the eukaryotic nonsense-mediated RNA decay system. *Genome Biol* 2003;4:R81. [PubMed: 14659018]
30. Clissold PM, Ponting CP. PIN domains in nonsense-mediated mRNA decay and RNAi. *Curr Biol* 2000;10:R888–R890. [PubMed: 11137022]
31. Boonyaratankornkit BB, Simpson AJ, Whitehead TA, Fraser CM, El-Sayed NM, Clark DS. Transcriptional profiling of the hyperthermophilic methanarchaeon *Methanococcus jannaschii* in response to lethal heat and non-lethal cold shock. *Environ Microbiol* 2005;7:789–797. [PubMed: 15892698]
32. Han CJ, Park SH, Kelly RM. Acquired thermotolerance and stressed-phase growth of the extremely thermoacidophilic archaeon *Metallosphaera sedula* in continuous culture. *Appl Environ Microbiol* 1997;63:2391–2396. [PubMed: 16535631]

33. Peeples TL, Kelly RM. Bioenergetic response of the extreme thermoacidophile *Metallosphaera sedula* to thermal and nutritional stresses. *Appl Environ Microbiol* 1995;61:2314–2321. [PubMed: 16535051]
34. Rohlin L, Trent JD, Salmon K, Kim U, Gunsalus RP, Liao JC. Heat shock response of *Archaeoglobus fulgidus*. *J Bacteriol* 2005;187:6046–6057. [PubMed: 16109946]
35. Shockley KR, Ward DE, Chhabra SR, Connors SB, Montero CI, Kelly RM. Heat shock response by the hyperthermophilic archaeon *Pyrococcus furiosus*. *Appl Environ Microbiol* 2003;69:2365–2371. [PubMed: 12676722]
36. Tachdjian S, Kelly RM. Dynamic metabolic adjustments and genome plasticity are implicated in the heat shock response of the extremely thermoacidophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* 2006;188:4553–4559. [PubMed: 16740961]
37. Trent JD, Osipiuk J, Pinkau T. Acquired thermotolerance and heat shock in the extremely thermophilic archaeobacterium *Sulfolobus sp.* strain B12. *J Bacteriol* 1990;172:1478–1484. [PubMed: 2106513]
38. Vierke G, Engelmann A, Hebbeln C, Thomm M. A novel archaeal transcriptional regulator of heat shock response. *J Biol Chem* 2003;278:18–26. [PubMed: 12381724]
39. Kagawa HK, Osipiuk J, Maltsev N, Overbeek R, Quait-Randall E, Joachimiak A, Trent JD. The 60 kDa heat shock proteins in the hyperthermophilic archaeon *Sulfolobus shibatae*. *J Mol Biol* 1995;253:712–725. [PubMed: 7473746]
40. Kagawa HK, Yaoi T, Brocchieri L, McMillan RA, Alton T, Trent JD. The composition, structure and stability of a group II chaperonin are temperature regulated in a hyperthermophilic archaeon. *Mol Microbiol* 2003;48:143–156. [PubMed: 12657051]
41. Schelert J, Drozda M, Dixit V, Dillman A, Blum P. Regulation of mercury resistance in the crenarchaeote *Sulfolobus solfataricus*. *J Bacteriol* 2006;188:7141–7150. [PubMed: 17015653]




Figure 1. The number of chromosomally encoded *vapBC* TA loci increases with optimal growth temperature for genome-sequenced archaea

T_{opt} data were obtained from The Prokaryotic Growth Temperature Database (<http://pgtdb.csie.ncu.edu.tw/>).




Figure 2. *Sulfolobus solfataricus* P2 *vapBC* TA loci transcriptome before and after heat shock (temperature shift from 80 to 90°C)

Red indicates up-regulation and blue represents down-regulation, relative to the genome-wide average transcription level. Missing *vapBs* were not annotated at the time the DNA microarray used here was fabricated.